

## Approach to equilibrium for a forced Burgers equation

WERNER KIRSCH AND BARRY SIMON<sup>1</sup>

*Abstract.* We show that approach to equilibrium in certain forced Burgers equations is implied by a decay estimate on a suitable intrinsic semigroup estimate, and we verify this estimate in a variety of cases including a periodic force.

### 1. Introduction

This paper is a contribution to the literature [10, 11, 3] on large time asymptotics of the forced Burgers equation

$$\frac{\partial u_i}{\partial t} + \sum_{j=1}^{\nu} u_j \frac{\partial u_j}{\partial x_i} = \frac{1}{2} \Delta u_i + \frac{\partial V(x)}{\partial x_i}, \quad t \geq 0, x \in \mathbb{R}^{\nu}, i = 1, \dots, \nu, \quad (1.1)$$

where  $u$  is real valued. We will make two assumptions on the initial data  $u_j(x, t = 0) = u_j^{(0)}(x)$ :

$$(i) \quad \frac{\partial u_i^{(0)}}{\partial x_j} = \frac{\partial u_j^{(0)}}{\partial x_i} \quad \text{all } i, j \quad (1.2)$$

$$(ii) \quad \psi_0(x) \equiv \int_0^x \vec{u}^{(0)}(y) \cdot dy \in L^{\infty}. \quad (1.3)$$

(1.2), which is vacuous in the standard  $\nu = 1$  case, implies that the value of  $\psi_0$  given by (1.3) is independent of the path taken from 0 to  $x$  in the line integral. Typical of our results is:

**THEOREM 1.1.** *Let  $V$  be a  $C^1$  periodic function on  $\mathbb{R}^{\nu}$ . Then, there is a unique initial condition  $u_{\infty}^{(0)}(x)$  obeying (1.2), (1.3) for (1.1) whose solution is independent of  $t$ . Moreover, if  $u^{(0)}$  is any other initial data obeying (1.2)/(1.3), then*

$$\lim_{t \rightarrow \infty} \sup_x [|u(x, t) - u_{\infty}^{(0)}(x)|] = 0. \quad (1.4)$$

---

Received February 26, 2001; accepted April 9, 2001.

<sup>1</sup>Supported in part by NSF Grant No. DMS-9707661.

REMARK. (1.3) does not imply that  $u^{(0)}$  is  $L^\infty$ , but our proof shows that for  $t > 0$ ,  $u(\cdot, t) \in L^\infty$ . Thus, for  $t > 0$ , the quantity in the limit in (1.4) is finite.

What is new about our ideas and Theorem 1.1 is that there is no regularity condition on the initial condition  $u$  at infinity other than (1.3). Previous approaches require at least that  $e^{-\psi_0}$  have some kind of average. To understand how we overcome this, we need to begin the proof by reminding the reader of the Cole-Hopf transformation. Define

$$\varphi_0(x) = \exp(-\psi_0(x)) \quad (1.5)$$

so that

$$u_i^{(0)} = -\varphi_0^{-1} \frac{\partial \varphi_0}{\partial x_i}. \quad (1.6)$$

$V$  is bounded so

$$H = -\frac{1}{2}\Delta + V \quad (1.7)$$

is bounded below. Thus we add a constant to  $V$  so that henceforth

$$\inf \text{spec}(H) = 0. \quad (1.8)$$

Define

$$\varphi(x, t) = (e^{-tH} \varphi_0)(x). \quad (1.9)$$

Then direct manipulation shows that

PROPOSITION 1.2. Let  $\psi_0 \in L^\infty$  be  $C^1$  and let  $\varphi_0, \varphi$  obey (1.5), (1.9). Then, for  $t > 0$ ,  $\varphi(x, t) > 0$  and  $\varphi, \nabla \varphi, \Delta \varphi$  are  $C^1$  and

$$u(x, t) = -\varphi(x, t)^{-1} (\nabla \varphi)(x, t) \quad (1.10)$$

obeys (1.1) for  $t > 0$  and  $\lim_{t \downarrow 0} u(\cdot, t) = u^{(0)} \equiv \nabla \psi_0$ .

REMARKS. 1. It follows from [9] with  $V$  a  $C^1$  function with bounded derivatives that  $\varphi$  and  $(-\Delta + V)\varphi$  are  $C^1$ . It follows that  $\Delta \varphi$  is  $C^1$  which by elliptic regularity means  $\nabla \varphi$  is  $C^1$ . The Laplacian here and in (1.1) may be distributional rather than classical. If  $\nabla V$  is assumed Hölder continuous, we can replace these by classical derivatives.

2. As we will discuss below,  $e^{-tH}$  maps  $L^\infty$  to  $L^\infty$  and (1.9) is intended in the sense of the  $L^\infty$  map.

3. This result does not require that  $V$  be periodic;  $V$  need only be  $C^1$  and in  $L^\infty$ . We will use it in this form below.

When  $V$  is periodic, it has a periodic ground state  $\Omega$ , that is, a positive periodic solution of

$$\left(-\frac{1}{2}\Delta + V\right)\Omega = 0. \tag{1.11}$$

Thus

$$u_\infty^{(0)}(x) = -\Omega(x)^{-1}(\nabla\Omega)(x) \tag{1.12}$$

is a stationary solution of (1.1). A natural approach to (1.4) is to prove that

$$\varphi \rightarrow c\Omega \tag{1.13}$$

and

$$\nabla\varphi \rightarrow c\nabla\Omega \tag{1.14}$$

both in  $L^\infty$ . This is essentially what previous works do.

To understand the limitations of this approach and why one can hope to go beyond them, consider the case  $V \equiv 0$ . Then  $u_\infty = 0$ ,  $\Omega \equiv 1$ , and

$$\varphi(x, t) = (2\pi t)^{-\nu/2} \int \exp(-(x - y)^2/2t)\varphi_0(y) dy.$$

If  $(2R)^{-\nu} \int_{\sup_i |y_i| \leq R} \varphi_0(y) d^\nu y \rightarrow c$  as  $R \rightarrow \infty$ , it is not hard to see that  $\varphi(x, t) \rightarrow c$  as  $t \rightarrow \infty$  for each fixed  $x$ , so (1.13) holds, but this is not true in general.

For example, if  $R_n = e^{e^n}$  and

$$\varphi_0(y) = 2 + (-1)^n \quad \text{if} \quad R_n < \sup_i |y_i| < R_{n+1},$$

then for  $t \sim R_n R_{n+1}$ , it is not hard to see that  $\varphi(0, t) \sim 2 + (-1)^n$  and thus  $\varphi(0, t)$  does not have a limit. But in this example, one can see that  $\nabla\varphi$  does go to zero.

Our key observations are that rather than prove (1.13) and (1.14) separately, it suffices to prove that

$$\nabla(\varphi/\Omega) \rightarrow 0 \tag{1.15}$$

and that (1.15) is equivalent to some estimates on the intrinsic semigroup associated to  $H$ . Specifically, let  $K_t(x, y)$  be the integral kernel of  $e^{-tH}$  and let

$$L_t(x, y) = \Omega^{-1}(x)K_t(x, y)\Omega(y)^{-1}. \tag{1.16}$$

$L_t$  is the kernel of a semigroup on  $L^2(\mathbb{R}^\nu, \Omega^2 d^\nu x)$ .

To prove Theorem 1.1, we will prove two estimates:

$$|\partial_x L_t(x, y)| \leq C t^{-\nu/2} [\exp(-D(x - y)^2/t) + \exp(-E|x - y|)] \tag{1.17}$$

for suitable  $C, D$  and all  $t > 1$ , all  $x, y$  and

$$|\partial_x L_t(x, y)| \leq C t^{-(v+1)/2} \quad (1.18)$$

all  $t > 1$ , all  $x, y$ .

We will show that if  $\Omega$  obeys  $0 < a \leq \Omega < b$ , is  $C^3$  and  $V \equiv \frac{1}{2}\Omega^{-1}(\Delta\Omega)$  is bounded and uniformly Hölder continuous, then (even if  $V$  is not periodic)

$$|\partial_x L_t(x, y)| \leq C t^{-\alpha} [\exp(-D(x-y)^2/t)] \quad (1.19)$$

for some  $\alpha > v/2$ , all  $t \geq 1$ . This will lead to the following generalization of Theorem 1.1:

**THEOREM 1.3.** *Let  $V$  be a  $C^1$  function and suppose that  $V$  is bounded and uniformly Hölder continuous. Suppose that  $-\frac{1}{2}\Delta + V$  has a ground state  $\Omega$  obeying  $0 < a \leq \Omega \leq b$  for some  $a, b$  and all  $t$ . Then there is a unique initial condition  $u_\infty^{(0)}(x) (= \nabla\Omega/\Omega)$  obeying (1.2), (1.3) for (1.1) whose solution is independent of  $t$ . Moreover, if  $u^{(0)}$  is any other initial data obeying (1.2)/(1.3), then*

$$\lim_{t \rightarrow \infty} \sup_x [|u(x, t) - u_\infty^{(0)}(x)|] = 0. \quad (1.20)$$

Certain quasiperiodic Schrödinger operators have a quasiperiodic ground state [5]. Thus the above theorem applies to this situation as well; see [11].

In Section 2, we will reduce Theorems 1.1 and 1.3 to (1.19). In Section 3, we will derive (1.17) using ideas due to Davies. In Section 4, we will prove (1.18) in the periodic case and (1.19) in general.

We are dedicating this paper to the memory of Tosio Kato, who taught us so much about Schrödinger operators, about semigroups, and about non-linear equations, areas which come together here.

## 2. Reduction to intrinsic heat kernel estimates

According to Proposition 1.2, the solution  $u$  of (1.1) is related to a solution of (1.9) via (1.16). This leads to

**THEOREM 2.1.** *Suppose there exist  $\Omega$  so*

$$e^{-tH}\Omega = \Omega \quad (2.1)$$

and

$$0 < a \leq \Omega(x) \leq b \quad (2.2)$$

and that

$$\lim_{t \rightarrow \infty} \|\vec{\nabla}(\varphi(\cdot, t)/\Omega(\cdot))\|_{\infty} = 0. \quad (2.3)$$

Then (1.20) holds with  $u_{\infty}^{(0)} = \nabla\Omega/\Omega$ .

*Proof.* By (1.3), we have

$$0 < c_1 \leq \varphi_0 \leq c_2$$

so by (2.2)

$$c_1 b^{-1} \Omega \leq \varphi_0 \leq c_2 a^{-1} \Omega.$$

Since  $e^{-tH}$  is positivity preserving and (2.1) holds,

$$c_1 b^{-1} \Omega(x) \leq \varphi(x, t) \leq c_2 a^{-1} \Omega(x).$$

So by (2.2) again,

$$c_1 b^{-1} a \leq \varphi(x, t) \leq c_2 a^{-1} b. \quad (2.4)$$

Now

$$\begin{aligned} u(\cdot, t) - u_{\infty}^{(0)} &= (\nabla\varphi)(\cdot, t)/\varphi(\cdot, t) - (\nabla\Omega)(\cdot)/\Omega(x) \\ &= (\Omega\nabla\varphi - \varphi\nabla\Omega)/\varphi\Omega \\ &= [\nabla(\varphi(\cdot, t)/\Omega)][\Omega/\varphi]. \end{aligned}$$

Since  $\varphi$  and  $\Omega$  are uniformly in  $t$  and  $x$  bounded above and below, we see that (1.20) is equivalent to (2.3).  $\square$

Now consider the unitary map  $U : L^2(\mathbb{R}^{\nu}) \rightarrow L^2(\mathbb{R}^{\nu}, \Omega^2 dx)$  by  $(Uf)(x) = f(x)\Omega(x)^{-1}$  and let  $M$  be the self-adjoint operator  $UHU^{-1}$  on  $L^2(\mathbb{R}^{\nu}, \Omega^2 dx)$ . Then, as is well-known (and a direct calculation),

$$(f, Mf)_{L^2(\mathbb{R}^{\nu}, \Omega^2 dx)} = \int (\nabla f)^2 \Omega^2 dx$$

or equivalently,

$$Mf = -\Delta f - 2(\vec{\nabla}\Omega)\Omega^{-1} \cdot \vec{\nabla}f. \quad (2.5)$$

Now let  $K_t(x, y)$  be the integral kernel of  $e^{-tH}$ , that is,

$$(e^{-tH} f)(x) = \int K_t(x, y) f(y) d^{\nu}y$$

and let  $L_t(x, y)$  be the integral kernel of  $e^{-tM}$ , that is,

$$(e^{-tM} f)(x) = \int L_t(x, y) f(y) \Omega^2(y) d^v y.$$

Since  $e^{-tM} = U e^{-tH} U^{-1}$ , we see that  $L_t$  and  $K_t$  are related by (1.16).

Now if  $\varphi = e^{-tH} \varphi_0$ , then

$$\begin{aligned} [\varphi(\cdot, t) / \Omega(\cdot)] &= U e^{-tH} \varphi_0 = e^{-tM} U \varphi_0 \\ &= \int L_t(\cdot, y) \varphi_0(y) \Omega(y) d^v y. \end{aligned}$$

Since  $\varphi_0$  and  $\Omega$  are uniformly bounded, we see that

$$|\nabla[\varphi(x, t) / \Omega(x)]| \leq c \int |\partial_x L_t(x, y)| d^v y.$$

Thus:

**PROPOSITION 2.2.** *A sufficient condition for (2.3) to hold for any initial  $\varphi_0$  (coming from a  $u_i^{(0)}$  obeying (1.2)/(1.3)) is that*

$$\sup_x \int |\partial_x L_t(x, y)| d^v y \rightarrow 0 \tag{2.6}$$

as  $t \rightarrow \infty$ .

**THEOREM 2.3.** *If (1.19) holds or if (1.17)/(1.18) hold, then (2.6) holds.*

*Proof.* (1.19) plus scaling implies that

$$\int |\partial_x L_t(x, y)| d^v y \leq C_1 t^{-\alpha} t^{v/2}$$

which goes to zero if  $t \rightarrow \infty$  since  $\alpha > v/2$ . (1.17)/(1.18) imply

$$|\partial_x L_t(x, y)| \leq C t^{-v/2-1/4} [\exp(-D(x-y)^2/2t) + \exp(-\frac{1}{2}E|x-y|)]$$

which implies

$$\int |\partial_x L_t(x, y)| d^v y \leq C_1 t^{-1/4} \pm C_2 t^{-v/2-1/4}$$

which goes to zero as  $t \rightarrow \infty$ . □

### 3. Exponential-Gaussian estimates on $\partial_x L_t$

Our goal in this section is to explain how one can get (1.17) from ideas of Davies [1, 2]. His ideas immediately imply an estimate

$$|K_t(x, y)| \leq C_\varepsilon t^{-v/2} \exp(-(x-y)^2/(4+\varepsilon)t) \tag{3.1}$$

for any  $\varepsilon > 0$ . (3.1) implies (even dropping the Gaussian) that

$$\|e^{-tH}\|_{L^1 \rightarrow L^\infty} \leq C t^{-\nu/2}$$

so by interpolation with boundedness on  $L^\infty$  (see Simon [9] and references therein),

$$\|e^{-tH}\|_{L^1 \rightarrow L^2} = \|e^{-tH}\|_{L^2 \rightarrow L^\infty} \leq C t^{-\nu/4}.$$

Thus

$$\|e^{-(t+is)H}\|_{L^1 \rightarrow L^\infty} \leq \|e^{-tH/2}\|_{L^2 \rightarrow L^\infty} \|e^{-isH}\|_{L^2 \rightarrow L^2} \|e^{-tH/2}\|_{L^1 \rightarrow L^2} \leq C t^{-\nu/2}$$

and thus for any  $|\theta| < \pi/2$ ,

$$\|\exp(-t e^{i\theta} H)\|_{L^1 \rightarrow L^\infty} \leq C_\theta t^{-\nu/2}$$

which yields

$$|K_{t e^{i\theta}}(x, y)| \leq C_\theta t^{-\nu/2}. \quad (3.2)$$

By interpolation between (3.1) and (3.2), we see that for complex  $t$  in a sector  $S_\theta = \{t \mid |\text{Arg}(t)| \leq \theta\}$  we have if  $\theta < \pi/2$ ,

$$|K_t(x, y)| \leq C_\theta |t|^{-\nu/2} \exp(-D_\theta(x-y)^2/|t|) \quad (3.3)$$

and this implies by a Cauchy estimate that in the same sectors

$$|\partial_t K_t(x, y)| \leq C_\theta |t|^{-\nu/2-1} \exp(-D_\theta(x-y)^2/|t|)$$

(where  $C_\theta, D_\theta$  can change value from one equation to the next).

Thus for  $t \geq 1$  and real,

$$|(-\Delta + V)_x K_t(x, y)| \leq C |t|^{-\nu/2-1} \exp(-D(x-y)^2/|t|). \quad (3.4)$$

Since  $V$  is bounded, (3.3) and (3.4) imply that for  $t \geq 1$ ,

$$|(-\Delta_x + 1)K_t(x, y)| \leq C |t|^{-\nu/2} \exp(-D(x-y)^2/|t|).$$

But  $\partial_x(-\Delta_x + 1)^{-1}$  has an explicit convolution integral kernel which is  $L^1$  at short distances and exponentially decaying at large. This implies for  $t \geq 1$

$$|(\partial_x K_t)(x, y)| \leq C t^{-\nu/2} [\exp(-D(x-y)^2/t) + \exp(-E|x-y|)]$$

which, with (3.3) and the formula (1.16), implies (1.17). We summarize:

**THEOREM 3.1.** *The estimate (1.17) holds for any potential  $V$  obeying the hypothesis of Theorem 1.3.*

#### 4. Improved time decay estimates on $\partial_t L_t$

Here the goal is to show that

$$|(\partial_x L_t)(x, y)| \leq C t^{-\alpha} \quad (4.1)$$

for some  $\alpha > \nu/2$ . We believe the estimate holds with  $\alpha = \nu/2 + 1/2$  and have proven this if  $V$  is periodic: One makes a Bloch wave decomposition [8] to write the semigroup as an integral over the Brillouin zone and a Gaussian approximation to control the resulting integral. One uses the fact that the minimum of the bottom band is known to be a unique point with a strictly quadratic minimum [4]. In general, we rely on estimates of Porper-Eidel'man [6] and only get  $\alpha > \nu/2$ . (In a later paper [7], they get  $\alpha = \nu/2 + 1/2$  for the case  $p = 1$  below that does not accommodate our situation.)

Indeed, (4.1) is exactly Corollary 3.4 of their paper which they prove for fundamental solutions of equations of the form

$$p(x)\partial_t u = \nabla \cdot (a(t, x)\nabla u), \quad (4.2)$$

where  $p, a$  and  $\nabla a$  are all Hölder continuous.

But (2.5) can be rewritten

$$Mf = -\Omega^{-2}\nabla \cdot (\Omega^2\nabla f)$$

so

$$\partial_t u = -Mu$$

is of the form (4.2) where

$$p = \Omega^2 \quad \text{and} \quad a_{ij} = \Omega^2 \delta_{ij}.$$

Thus, their result applies so long as  $\Omega$  is  $C^1$  with  $\nabla\Omega$  uniformly Hölder continuous.

Since  $\Delta\Omega = V\Omega$  with  $V \in C^1$  and  $\Omega$  a priori bounded, we see that

$$\nabla\Omega = \nabla(-\Delta + 1)^{-1}(-V + 1)\Omega.$$

Since  $(-V + 1)\Omega$  is bounded, the explicit integral kernel for  $\nabla(-\Delta + 1)^{-1}$  shows  $\nabla\Omega$  is uniformly Hölder continuous of any order less than 1. Thus, their Corollary 3.4 applies and (4.1) holds.

#### Acknowledgments

W. K. would like to thank T. Tombrello for the hospitality of Caltech where this research was done. We would like to thank E. B. Davies, T. Coulhon, A. Grigoryan, Y. Pinchover, and D. Robinson for valuable correspondence.

## REFERENCES

- [1] DAVIES, E. B., *Explicit constants for Gaussian upper bounds on heat kernels*, Amer. J. Math. 109 (1987), 319–334.
- [2] DAVIES, E. B., *Heat Kernels and Spectral Theory*, Cambridge Tracts in Mathematics 92, Cambridge University Press, Cambridge, 1989.
- [3] KIRSCH, W. and KUTZELNIGG, A., *Time asymptotics for solutions of the Burgers equation with a periodic force*, Math. Z. 232 (1999) no. 4, 691–705.
- [4] KIRSCH, W. and SIMON, B., *Comparison theorems for the gap of Schrödinger operators*, J. Funct. Anal. 75 (1987), 396–410.
- [5] KOZLOV, S. M., *Ground states of quasiperiodic operators*, Sov. Math. Dokl. 28 (1983), 114–119. Translated from Dokl. Akad. Nauk SSSR 271 (1983), 532–536.
- [6] PORPER, F. O. and EIDEL'MAN, S. D., *Two-sided estimates of fundamental solutions of second-order parabolic equations, and some applications*, Russian Math. Surveys 39 (1984) no. 3, 119–178. Translated from Usp. Mat. Nauk. 39 (1984) no. 3, 107–156.
- [7] PORPER, F. O. and EIDEL'MAN, S. D., *Theorems related to the asymptotic proximity of fundamental solutions to parabolic equations*, Dokl. Math. 52 (1995) no. 2, 237–240. Translated from Dokl. Akad. Nauk 344 (1995) no. 5, 586–589.
- [8] REED, M. and SIMON, B., *Methods of Modern Mathematical Physics, IV. Analysis of Operators*, Academic Press, New York, 1978.
- [9] SIMON, B., *Schrödinger semigroups*, Bull. Amer. Math. Soc. 7 (1982), 447–526.
- [10] SINAI, YA. G., *Two results concerning asymptotic behavior of solutions of the Burgers equation with force*, J. Stat. Phys. 64 (1991) no. 1–2, 1–12.
- [11] SINAI, YA. G., *Asymptotic behavior of solutions of 1D-Burgers equation with quasi-periodic forcing*, Topol. Methods Nonlinear Anal. 11 (1998) no. 2, 219–226.

Werner Kirsch  
Institut für Mathematik  
Ruhr-Universität Bochum  
44780 Bochum  
Germany  
e-mail: werner@mathphys.ruhr-uni-bochum.de

Barry Simon  
Division of Physics, Mathematics, and Astronomy  
253-37, California Institute of Technology  
Pasadena, CA 91125-3700  
USA  
e-mail: bsimon@caltech.edu



To access this journal online:  
<http://www.birkhauser.ch>

---