

Poisson brackets of orthogonal polynomials

María José Cantero^{a,1}, Barry Simon^{b,*,2}

^aDepartment of Applied Mathematics, University of Zaragoza, 50018 Zaragoza, Spain

^bMathematics 253-37, California Institute of Technology, Pasadena, CA 91125, USA

Received 9 March 2007; received in revised form 4 June 2007; accepted 9 July 2007

Available online 14 September 2007

Communicated by C.K. Chui and H.N. Mhaskar

Dedicated to the memory of G.G. Lorentz

Abstract

For the standard symplectic forms on Jacobi and CMV matrices, we compute Poisson brackets of OPRL and OPUC, and relate these to other basic Poisson brackets and to Jacobians of basic changes of variable. © 2007 Elsevier Inc. All rights reserved.

Keywords: Jacobi matrix; CMV matrix; OPUC; OPRL

1. Introduction

It has been known since the discoveries of Flaschka [15] and Moser [37] concerning the finite Toda lattice as a completely integrable system that finite Jacobi matrices of fixed trace support a natural symplectic form. Explicitly, if

$$J = \begin{pmatrix} b_1 & a_1 & & & \\ a_1 & b_2 & a_2 & & \\ & \ddots & \ddots & \ddots & \\ & & a_{N-1} & b_N & \end{pmatrix} \quad (1.1)$$

* Corresponding author.

E-mail addresses: mjcanter@unizar.es (M.J. Cantero), bsimon@caltech.edu (B. Simon).

¹ Supported in part by a research grant from the Ministry of Education and Science of Spain, MTM2005-08648-C02-01, and by the Aragón government, Project E-64 of DGA.

² Supported in part by NSF Grant DMS-0140592 and U.S.–Israel Binational Science Foundation (BSF) Grant No. 2002068.

on the manifold where $a_j > 0$ and

$$\sum_{j=1}^N b_j = c \quad (1.2)$$

the nonzero Poisson brackets of the a 's and b 's are

$$\{b_k, a_k\} = -\frac{1}{4} a_k, \quad k = 1, \dots, N-1, \quad (1.3)$$

$$\{b_k, a_{k-1}\} = \frac{1}{4} a_{k-1}, \quad k = 2, \dots, N. \quad (1.4)$$

For the basic definitions of symplectic manifolds and Poisson brackets, see Deift [9].

The nonzero Poisson brackets occur for neighboring elements in J . As one would guess, there are differing conventions, so that sometimes the $\frac{1}{4}$ in (1.3) and (1.4) are $\frac{1}{2}$ and sometimes one takes $\{, \}$ to be the negative of our choice.

It is easy to check that $\sum_{j=1}^N b_j$ commutes with all a 's and b 's as is necessary for this to define a bracket on the manifold where (1.2) holds. It is also easy to see that the form defined by (1.3)/(1.4) is nondegenerate and closed so there is an underlying symplectic form; see the end of Section 3.

If one takes

$$H = 2 \operatorname{Tr}(J^2) = 2 \sum_{j=1}^N b_j^2 + 4 \sum_{j=1}^{N-1} a_j^2 \quad (1.5)$$

then the Hamiltonian flow,

$$\dot{b}_k = \{H, b_k\} = 2(a_k^2 - a_{k-1}^2), \quad (1.6)$$

$$\dot{a}_k = \{H, a_k\} = a_k(b_{k+1} - b_k) \quad (1.7)$$

is the Toda flow in Flaschka form [15]. It is completely integrable; indeed, (see (1.11))

$$\{\operatorname{Tr}(J^n), \operatorname{Tr}(J^m)\} = 0 \quad (1.8)$$

for all n, m .

If $d\rho$ is the spectral measure for J and $(1, 0, \dots, 0)^t$, then

$$d\rho = \sum_{j=1}^N \rho_j \delta_{x_j}. \quad (1.9)$$

The $\{\rho_j\}_{j=1}^N$ and $\{x_j\}_{k=1}^N$ are not independent; indeed, by (1.2),

$$\sum_{j=1}^N x_j = c, \quad \sum_{j=1}^N \rho_j = 1. \quad (1.10)$$

Fundamental to developments are the following Poisson brackets:

$$\{x_j, x_k\} = 0 \quad \{x_j, \rho_k\} = \frac{1}{2} [\delta_{jk} \rho_j - \rho_j \rho_k]. \quad (1.11)$$

The analog of $\{x_j, x_k\} = 0$ for periodic Toda chains is due to Flaschka [15] and (1.11) appeared (implicitly) first in Moser [37].

Our original motivation was to understand the analog of this for CMV matrices. Along the way, we realized that Poisson brackets of orthogonal polynomials were not previously studied, even in the Jacobi case, so we decided to discuss both cases.

We also felt that the centrality of (1.11), which is partly known to experts, was not always so clear, so we also decided to say something about that. Indeed, one of our initial goals was to prove the analog of $\{x_i, x_j\} = 0$ for the periodic case of OPUC whose previous proofs [47 (Section 11.11),39,40] were involved. Hence one application of (1.11) we discuss is going from (1.11) to the periodic case.

Since $\sum_{j=1}^N x_j$ and $\sum_{j=1}^N \rho_j$ are constant, $\{x_j, \rho_k\}$ must sum to zero over j or k , as can be checked in (1.11) since $\sum_{j=1}^N \rho_j = 1$.

We note one curiosity of (1.11), namely, $\{x_j, \rho_k\}$ is symmetric in j and k . We will eventually also find

$$\{\rho_j, \rho_k\} = \frac{\rho_j \rho_k}{x_j - x_k} - \sum_{m \neq j} \frac{\rho_j \rho_k \rho_m}{x_j - x_m} + \sum_{m \neq k} \frac{\rho_j \rho_k \rho_m}{x_k - x_m} \tag{1.12}$$

but will make no use of this complicated formula here. It is needed if one wants to find angle variables for the Toda flows (see [32,18]).

Simultaneous to our work, Gekhtman–Nenciu [18] were studying Poisson brackets of Carathéodory functions, which we will see is closely related to our work. Indeed, a key to our progress was learning of this work and the earlier paper of Faybusovich–Gekhtman [14]. We also mention not unrelated earlier papers of Kako–Mugibayashi [29,30] and Kulish [35]. During the refereeing of this paper, we learned of a preprint of Tsiganov [53], clearly independent of ours, that also computes Poisson brackets of transfer matrices which include some of our OP Poisson brackets.

While we focus on a new approach to Poisson brackets, we would emphasize to the reader that there are long-standing powerful algebraic methods (e.g., [33]) with recent developments especially relevant to our work [22,36].

We now turn to methods using orthogonal polynomials on the real line (OPRL) and unit circle (OPUC); for background, recursion relations, etc., see [17,52,6,46,47]. For quick introductions, see Section 1.2 of [46] for OPRL and [45] for OPUC.

For the monic OPRL, $P_n(x)$, our basic result is that for $n = 1, 2, \dots, N$,

$$\{P_n(x), P_n(y)\} = \{P_{n-1}(x), P_{n-1}(y)\} = 0 \tag{1.13}$$

$$2\{P_n(x), P_{n-1}(y)\} = \left[\frac{P_n(x)P_{n-1}(y) - P_n(y)P_{n-1}(x)}{x - y} \right] - P_{n-1}(x)P_{n-1}(y). \tag{1.14}$$

Notice the occurrence of the Bezoutian (see [23]) of P_n and P_{n-1} in the first term on the right of (1.14). (1.13) and (1.14) will follow by a direct induction in n .

There is a symmetry between P_n, P_{n-1} and P_n, Q_n (where Q_n is the second kind polynomial (of degree $n - 1$) for P_n), for P_N is a determinant of $z - J$ and P_{N-1} (resp. Q_N) are the determinants obtained by removing the last and rightmost row and column (resp. first and

leftmost row and column). This will immediately lead to

$$\{P_n(x), P_n(y)\} = \{Q_n(x), Q_n(y)\} = 0, \tag{1.15}$$

$$2\{P_n(x), Q_n(y)\} = - \left[\frac{P_n(x)Q_n(y) - P_n(y)Q_n(x)}{x - y} \right] + Q_n(x)Q_n(y). \tag{1.16}$$

The sign changes because inverting order changes the sign of $\{ , \}$. All these calculations appear in Section 2. In [14], several different Poisson brackets on polynomials are considered, including one that is essentially equivalent to (1.16).

In Section 3, we derive $\{x_j, x_k\} = 0$ from $\{P_n(x), P_n(y)\} = 0$, $\{x_j, \rho_k\}$ from $\{P_n(x), Q_n(y)\}$ and $\{\rho_j, \rho_k\}$ from $\{Q_n(x), Q_n(y)\} = 0$.

Looked at carefully, if we argued directly for P, Q by induction, we, in essence, use coefficient stripping to relate P_N, Q_N for $\{a_j\}_{j=1}^{N-1} \cup \{b_j\}_{j=1}^N$ to P_{N-1}, Q_{N-1} for $\{a_j\}_{j=2}^{N-1} \cup \{b_j\}_{j=2}^{N-1}$. The analog for OPUC is a new coefficient stripping relation of first and second kind paraorthogonal polynomials, which is new. We present this in Section 4, denoting these first and second kind OPUC by P_N and Q_N .

In Section 5, we use the natural symplectic form for OPUC: in terms of Verblunsky coefficients,

$$\{\alpha_j, \alpha_k\} = 0, \quad \{\alpha_j, \bar{\alpha}_k\} = -i\rho_j^2\delta_{jk}. \tag{1.17}$$

This was introduced by Nenciu–Simon [42]. We will find

$$\{P_n(z), P_n(w)\} = 0 = \{Q_n(z), Q_n(w)\}, \tag{1.18}$$

$$\begin{aligned} \{P_n(z), Q_n(w)\} = & -\frac{i}{2} \left[(P_n(z)Q_n(w) - P_n(w)Q_n(z)) \left(\frac{z+w}{z-w} \right) \right. \\ & \left. - Q_n(z)Q_n(w) + P_n(z)P_n(w) \right] \end{aligned} \tag{1.19}$$

using induction and the coefficient stripping of Section 4. P_n, Q_n will depend on a parameter β . $\{\alpha_j\}_{j=0}^{n-2} \rightarrow \{-\alpha_j\}_{j=0}^{n-2}$ which preserves OPUC Poisson brackets and $\beta \rightarrow -\beta$ interchanges P_n and Q_n , so the right side of (1.20) has to be antisymmetric under interchange of P_n and Q_n , as it is.

In Section 6, we derive the analogs of (1.11). In (1.11), the x 's and ρ 's are global variables fixed by $x_1 < x_2 < \dots$. For OPUC, the eigenvalues are only locally well defined, namely,

$$d\mu = \sum_{j=1}^N \mu_j \delta_{z_j}, \tag{1.20}$$

where

$$z_j = e^{i\theta_j}. \tag{1.21}$$

The analogs of (1.10) are

$$\sum_{j=1}^N \theta_j = c, \quad \sum_{j=1}^N \mu_j = 1 \tag{1.22}$$

and of (1.11),

$$\{\theta_j, \theta_k\} = 0, \quad \{\theta_j, \mu_k\} = \mu_j \delta_{jk} - \mu_j \mu_k. \tag{1.23}$$

The analog of $\{\theta_j, \theta_k\} = 0$ for the periodic case appeared first in [42], and $\{\theta_j, \theta_k\} = 0$ is from [39,40]. The formula for $\{\theta_j, \mu_k\}$ is due to Killip–Nenciu [32].

In Sections 7–11, we discuss three applications of the fundamental relations (1.11) and (1.23): to computations of Jacobians of the maps $(a, b) \rightarrow (x, \rho)$ and $(\alpha) \rightarrow (\theta, \mu)$, to the exact solution of the flows generated by $G(x)$ or $G(\theta)$, and to the periodic case. In Sections 9 and 10, we discuss the differential equations induced on the monic OPRL and OPUC. Sections 12 and 13 compute more Poisson brackets for OPRL and OPUC, respectively.

2. Poisson brackets of first and second kind OPRL

Our goal in this section is to prove (1.13) through (1.16). We note that since decreasing N to $N - 1$ does not change the Poisson brackets for $\{a_k\}_{k=1}^{N-2}$, $\{b_j\}_{j=1}^{N-1}$ nor the $\{P_j\}_{j=1}^{N-1}$, we can suppose $n = N$ in proving (1.13)–(1.16) but then increase N in the induction step.

Proposition 2.1. (1.13) and (1.14) holds for $n = 1$.

Proof. For $N = 1$, $b_1 = c$, $P_0(x) = 1$, $P_1(x) = x - c$, and all Poisson brackets are zero. Thus, we need only show the RHS of (1.14) is zero, that is,

$$\frac{(x - c) - (y - c)}{x - y} - 1 = 0$$

which is obvious. \square

Lemma 2.2.

$$\{a_n^2, P_n(w)\} = -\frac{1}{2} a_n^2 P_{n-1}(w). \tag{2.1}$$

Proof. Since

$$P_{j+1}(x) = (x - b_{j+1})P_j(x) - a_j^2 P_{j-1}(x) \tag{2.2}$$

we have inductively that P_j is a function of $\{b_\ell\}_{\ell=1}^j \cup \{a_\ell\}_{\ell=1}^{j-1}$ as also follows from (2.8). Thus, we have $\{a_n^2, P_{n-1}(w)\} = \{a_n^2, P_{n-2}(w)\} = 0$ so, using (2.2) for $j + 1 = n$,

$$\begin{aligned} \{a_n^2, P_n(w)\} &= \{a_n^2, -b_n\}P_{n-1}(w) \\ &= -\frac{1}{2} a_n^2 P_{n-1}(w). \quad \square \end{aligned}$$

Theorem 2.3. (1.13) and (1.14) hold for all n .

Proof. Proposition 2.1 is the result for $n = 1$. So, by induction, we can suppose (1.13) and (1.14) for $j \leq n$ and need only prove it for $n + 1$. By (2.2) for $j = n$ and the facts that $\{P_{n-1}(x), P_{n-1}(y)\} = \{P_n(x), P_n(y)\} = 0$ and P_n independent of a_n , a_{n+1} and P_{n-1} independent of b_n and b_{n+1} , only the cross terms enter and

$$-\{P_{n+1}(x), P_{n+1}(y)\} = \{(x - b_{n+1})P_n(x), a_n^2 P_{n-1}(y)\} - (x \leftrightarrow y)$$

where $(x \leftrightarrow y)$ is like the first term but x and y are reversed.

This is a sum of three terms: t_1, t_2, t_3 where

$$\begin{aligned}
 2t_1 &= 2(x - b_{n+1})a_n^2\{P_n(x), P_{n-1}(y)\} - (x \leftrightarrow y) \\
 &= (x - y)a_n^2 \left[\frac{(P_n(x) - P_{n-1}(y) - P_n(y)P_{n-1}(x))}{x - y} - P_{n-1}(x)P_{n-1}(y) \right] \tag{2.3}
 \end{aligned}$$

by the induction hypothesis and the symmetry of $\{P_n(y), P_{n-1}(y)\}$ under $x \leftrightarrow y$. Next,

$$\begin{aligned}
 2t_2 &= 2\{xP_n(x), a_n^2\}P_{n-1}(y) - (x \leftrightarrow y) \\
 &= a_n^2(x - y)P_{n-1}(x)P_{n-1}(y) \tag{2.4}
 \end{aligned}$$

by Lemma 2.2. Finally,

$$\begin{aligned}
 2t_3 &= -2\{b_{n+1}, a_n^2\}(P_n(x)P_{n-1}(y) - (x - y)) \\
 &= -a_n^2(P_n(x)P_{n-1}(y) - P_n(y)P_{n-1}(x)). \tag{2.5}
 \end{aligned}$$

There is a fourth term involving $\{b_{n+1}, P_{n-1}(y)\}$, but this is zero.

Clearly, $t_1 + t_2 + t_3 = 0$ since (2.5) cancels the first term in (2.3), and (2.4) the second term. This proves that $\{P_{n+1}(x), P_{n+1}(y)\} = 0$.

Since $\{P_n(x), P_n(y)\} = 0$ and $\{b_{n+1}, P_n(y)\} = 0$ (since $P_n(y)$ only depends on $\{a_j\}_{j=0}^{n-1}$), (2.2) implies

$$\begin{aligned}
 \{P_{n+1}(x), P_n(y)\} &= \{-a_n^2P_{n-1}(x), P_n(y)\} \\
 &= -a_n^2\{P_{n-1}(x), P_n(y)\} - P_{n-1}(x)\{a_n^2, P_n(y)\}. \tag{2.6}
 \end{aligned}$$

The first term is evaluated by induction and the second by Lemma 2.2. The Lemma 2.2 term is $+\frac{1}{2}a_n^2P_{n-1}(x)P_{n-1}(y)$ which cancels one term in $\{P_{n-1}(x), P_n(y)\}$. Thus

$$\begin{aligned}
 2\{P_{n+1}(x), P_n(y)\} &= a_n^2 \left[\frac{P_n(y)P_{n-1}(x) - P_n(x)P_{n-1}(y)}{y - x} \right] \\
 &= \left[-\frac{P_n(y)(P_{n+1}(x) - (x - b_{n+1})P_n(x)) - (x \leftrightarrow y)}{y - x} \right] \\
 &= \frac{(P_{n+1}(y)P_n(y) - P_{n+1}(y)P_n(x))}{x - y} - P_n(x)P_n(y) \tag{2.7}
 \end{aligned}$$

proving the required formula for $n + 1$. In the above, (2.7) comes from (2.2). \square

The monic second kind polynomials, Q_n , can be defined as follows: We note that if $J(b_1, \dots, b_n; a_1, \dots, a_{n-1})$ is the matrix (1.1) with $N = n$, then

$$P_n(x) = \det(x - J(b_1, \dots, b_n; a_1, \dots, a_{n-1})) \tag{2.8}$$

as is well known. Q_n is then defined by removing the top row and left column, that is,

$$Q_n(x) = \det(x - J(b_2, \dots, b_n; a_2, \dots, a_{n-1})). \tag{2.9}$$

Notice that $J(b_1, \dots, b_n; a_1, \dots, a_{n-1})$ and $J(b_n, b_{n-1}, \dots, b_1; a_{n-1}, \dots, a_1)$ are unitarily equivalent under the unitary $U\delta_j = \delta_{n+1-j}$, $j = 1, \dots, n$. This shows making the a, b dependence explicit (we include redundant variables; Q_n is independent of b_1 and a_1):

$$\begin{aligned}
 P_n(x; b_1, \dots, b_n; a_1, \dots, a_{n-1}) &= P_n(x; b_n, \dots, b_1; a_{n-1}, \dots, a_1), \\
 Q_n(x; b_1, \dots, b_n; a_1, \dots, a_{n-1}) &= P_{n-1}(x; b_n, \dots, b_1; a_{n-1}, \dots, a_1).
 \end{aligned}$$

Taking into account that reversing order changes the signs in (1.14) and (1.16), we see that $\{ \cdot, \cdot \}$ changes sign, that is, (1.14) implies (1.16), and we have

Theorem 2.4. (1.15) and (1.16) hold for all n .

Instead of using induction from P_n, P_{n-1} to P_{n+1}, P_n and the above reversal argument, we can directly use induction from P_n, Q_n to P_{n+1}, Q_{n+1} using

$$Q_{n+1}(x; b_1, \dots, b_{n+1}; a_1, \dots, a_n) = P_n(x; b_2, \dots, b_{n+1}; a_2, \dots, a_n) \tag{2.10}$$

and

$$\begin{aligned} &P_{n+1}(x; b_1, \dots, b_{n+1}; a_1, \dots, a_n) \\ &= (x - b_1)P_n(x; b_2, \dots, b_{n+1}; a_2, \dots, a_n) - a_1^2 Q_n(x; b_2, \dots, b_{n+1}; a_2, \dots, a_n) \end{aligned} \tag{2.11}$$

that is, coefficient stripping.

3. Fundamental Poisson brackets for OPRL

Our goal in this section is to prove (1.12) and, most importantly, (1.11). In the calculations below, we will compute Poisson brackets of functions of a, b and free parameter(s), x (e.g., $P_n(x)$) and then set x to a value that is a function, h , of a, b . The order of operations is important, that is, $\{ \cdot, \cdot \}$ first, only then evaluation. We will denote this by $\{f, g\}|_{x=h}$.

Proposition 3.1.

$$\{P_n(x), P_n(y)\}|_{x=x_j, y=x_k} = \{x_j, x_k\} \prod_{\ell \neq j} (x_j - x_\ell) \prod_{\ell \neq k} (x_k - x_\ell). \tag{3.1}$$

In particular,

$$\{P_n(x), P_n(y)\} = 0 \Leftrightarrow \{x_j, x_k\} = 0 \quad \text{for all } j, k. \tag{3.2}$$

Proof. Since P is monic and its zeros are simple and (2.8) holds,

$$P_n(x) = \prod_j (x - x_j). \tag{3.3}$$

Thus, by Leibnitz’s rule,

$$\{P_n(x), P_n(y)\} = \sum_{p,q} \{x_p, x_q\} \prod_{\ell \neq p} (x - x_\ell) \prod_{\ell \neq q} (y - x_\ell). \tag{3.4}$$

Setting $x = x_j, y = x_k$, all terms with $p \neq j$ or $q \neq k$ vanish and we obtain (3.1). (3.1) trivially implies \Rightarrow in (3.2) and (3.4) implies \Leftarrow in (3.2). \square

Proposition 3.2. We have that

$$Q_n(x) = \sum_{j=1}^n \rho_j \prod_{k \neq j} (x - x_k). \tag{3.5}$$

Proof. The m -function is defined by

$$m_n(z) = \sum_{j=1}^n \frac{\rho_j}{x_j - z} \tag{3.6}$$

which is just

$$m_n(z) = \langle \delta_1, (J(b_1, \dots, b_n; a_1, \dots, a_{n-1}) - z)^{-1} \delta_1 \rangle \tag{3.7}$$

which, by Cramer’s rule and (2.8)/(2.9), implies

$$m_n(z) = -\frac{Q_n(z)}{P_n(z)}. \tag{3.8}$$

Thus, by (3.3),

$$\begin{aligned} Q_n(z) &= -m_n(z)P_n(z) \\ &= \sum_{j=1}^n \frac{\rho_j (\prod_{k \neq j} (z - x_k))}{(z - x_j)} \end{aligned}$$

which is (3.5). \square

Proposition 3.3.

$$\{P_n(x), Q_n(y)\}|_{x=x_j, y=x_k} = -\{x_j, \rho_k\} \prod_{\ell \neq j} (x_j - x_\ell) \prod_{\ell \neq k} (x_k - x_\ell). \tag{3.9}$$

This implies that

$$(1.16) \Leftrightarrow \{x_j, \rho_k\} = \frac{1}{2} [\delta_{jk} \rho_j - \rho_j \rho_k]. \tag{3.10}$$

Proof. Since we have proven that $\{x_j, x_k\} = 0$, (3.3) and (3.5) imply that

$$\{P_n(x), Q_n(y)\} = -\sum_{p,q} \{x_p, \rho_q\} \prod_{\ell \neq p} (x - x_\ell) \prod_{\ell \neq q} (y - x_\ell) \tag{3.11}$$

which immediately implies (3.9).

In (1.16), we have, by (3.5), that

$$Q_n(x)Q_n(y)|_{x=x_j, y=x_k} = \{\rho_j \rho_k\} \prod_{\ell \neq j} (x_j - x_\ell) \prod_{\ell \neq k} (x_k - x_\ell). \tag{3.12}$$

The Bezoutian term in (1.16) is 0 if $j \neq k$, but is slightly subtle if $j = k$ because then the formal expression is 0/0. We thus take limits since, of course, $\{P_n(x), Q_n(y)\}$, as a polynomial in x and y , is continuous. Since $P_n(x_k) = 0$, for $x \neq x_k$,

$$\frac{P_n(x)Q_n(x_k) - P_n(x_k)Q_n(x)}{x - x_k} = \left[\prod_{\ell \neq k} (x - x_\ell) \right] \rho_k \left[\prod_{\ell \neq k} (x - x_\ell) \right]. \tag{3.13}$$

Thus

$$\lim_{\substack{x \rightarrow x_j \\ x \neq x_k}} \text{LHS of (3.13)} = \rho_k \left[\prod_{\ell \neq k} (x_k - x_\ell) \right]^2 \delta_{jk}. \tag{3.14}$$

Thus (1.16) implies the right side of (3.10). By (3.11), the right side of (3.10) implies (1.16).

Remark. The symmetry of $\{x_j, \rho_k\}$ under $j \leftrightarrow k$ is equivalent to the symmetry of the right side of (1.16) under $x \leftrightarrow y$.

Proposition 3.4. *Let $j \neq k$. Then*

$$\{\rho_j, \rho_k\} = \sum_{m \neq j} \frac{\rho_j \{x_m, \rho_k\} + \rho_m \{x_j, \rho_k\}}{x_j - x_m} - \sum_{m \neq k} \frac{\rho_k \{x_m, \rho_j\} + \rho_m \{x_k, \rho_j\}}{x_k - x_m} \tag{3.15}$$

$$= \frac{\rho_j \rho_k}{x_j - x_k} - \sum_{m \neq j} \frac{\rho_j \rho_k \rho_m}{x_j - x_m} + \sum_{m \neq k} \frac{\rho_j \rho_k \rho_m}{x_k - x_m}. \tag{3.16}$$

Proof. (3.15) comes from $\{Q_n(x), Q_n(y)\} = 0$. Explicitly,

$$\frac{\{Q_n(x), Q_n(y)\}_{x=x_j, y=x_k}}{\prod_{\ell \neq j} (x_j - x_\ell) \prod_{\ell \neq k} (x_k - x_\ell)} = \{\rho_j, \rho_k\} - \text{RHS of (3.15)}$$

by a straightforward but tedious calculation. (3.16) then follows by using (1.11) for $\{x_p, \rho_q\}$. \square

Remark. There is a sense in which (1.11) implies (1.12) since (1.11) implies $\{P_n(x), P_n(y)\} = 0$ which means $\{Q_n(x), Q_n(y)\} = 0$ (since Q_n is a P_{n-1} for suitable Jacobi parameters), and this yields (3.15).

Finally, we want to translate the basic Poisson brackets relations (1.14)/(1.15) and (1.11) into assertions about the two-form which defines the symplectic structure underlying the Poisson brackets. Recall a symplectic structure is defined by a two-form, ω , that is, an antisymmetric functional on tangent vectors with the requirement that ω is closed (which is equivalent to the Jacobi identity for the Poisson bracket; see, e.g., [9]) and which is nondegenerate; that is, for all tangent vectors $v \neq 0$ at $p \in M$, there is \tilde{v} at p so $\omega(v, \tilde{v}) \neq 0$. Nondegeneracy implies $d = \dim(M)$ is even.

Given such a two-form and function, f , the Hamiltonian vector field H_f is defined by

$$\omega(H_f, v) = \langle df, v \rangle \tag{3.17}$$

for all tangent vectors v . Here df is the one-form $\sum_{i=1}^d \frac{\partial f}{\partial x_i} dx_i$ as usual, and $\langle \cdot, \cdot \rangle$ is the linear algebra pairing of vectors and their duals.

The Poisson bracket associated to ω is then

$$\{f, g\} = H_f(g). \tag{3.18}$$

Proposition 3.5. *Let M be a symplectic manifold of dimension $d = 2\ell$ and suppose there is a local coordinate system $\{x_j\}_{j=1}^\ell, \{y_j\}_{j=1}^\ell$, so that*

$$\omega = \sum_{i,j=1}^\ell W_{ij} dx_i \wedge dy_j + \sum_{i,j=1}^\ell U_{ij} dx_i \wedge dx_j \tag{3.19}$$

(i.e., no $dy_i \wedge dy_j$ terms) with U antisymmetric. Then

(i) *Nondegeneracy of ω is equivalent to invertibility of the $\ell \times \ell$ matrix $(W_{ij})_{1 \leq i, j \leq \ell}$. We will denote its inverse by $(W^{-1})_{ij}$.*

(ii)

$$\omega^\ell = \ell! \det(W) dx_1 \wedge dy_1 \wedge dx_2 \wedge \dots \wedge dy_\ell. \tag{3.20}$$

(iii)

$$\{x_i, x_j\} = 0. \tag{3.21}$$

(iv)

$$\{x_i, y_j\} = (W^{-1})_{ji}. \tag{3.22}$$

Moreover, $\{y_i, y_j\} = 0$ for all i, j if and only if $U \equiv 0$.

Conversely, if a symplectic manifold has a Poisson bracket obeying (3.21) and (3.22), then ω has the form (3.19).

Remarks. (1) In (3.20), ω^ℓ means the ℓ -fold wedge product $\omega \wedge \dots \wedge \omega$.

(2) W and U can be functions of x, y .

(3) These calculations do not use $d\omega = 0$ (or, equivalently, the Jacobi identity).

Proof. Write

$$v = \sum_{k=1}^\ell \left(r_k \frac{\partial}{\partial x_k} + t_k \frac{\partial}{\partial y_k} \right).$$

If $\vec{t} \neq 0$, pick $\tilde{s} = W^{-1}t$ and $\tilde{v} = \sum_{k=1}^\ell \tilde{s}_k \frac{\partial}{\partial x_k}$. If $\vec{t} = 0, \vec{r} \neq 0$, and with $\tilde{s} = W^{-1}r$, we can take $\tilde{v} = \sum_{k=1}^\ell \tilde{s}_k \frac{\partial}{\partial y_k}$. Thus, invertibility implies nondegeneracy. If W is not invertible, then $\det(W) = 0$ and the calculation of (ii) shows ω is degenerate.

(ii) is an easy calculation: using the distributive law to expand ω^ℓ , any term with a $dx_i \wedge dx_j$ term has at least $\ell + 1$ dx 's, and so is zero by antisymmetry. The only products of $\ell dx_i \wedge dy_j$ that are nonzero are of the form $[dx_{\pi(1)} \wedge dy_{\sigma(1)}] \wedge \dots \wedge [dx_{\pi(\ell)} \wedge dy_{\sigma(\ell)}]$ with π, σ permutations, and this is $(-1)^{\sigma\pi^{-1}} dx_1 \wedge dy_1 \wedge \dots \wedge dy_\ell$. The sum over σ for fixed π yields $\det(W)$ and then the sum over π gives $\ell!$.

(iii), (iv). We begin by noting that if $\{\zeta_j\}_{j=1}^{2\ell}$ is any coordinate system, if Ω is a $2\ell \times 2\ell$ antisymmetric real matrix, and

$$\omega = \sum_{j,k=1}^{2\ell} \Omega_{jk} d\zeta_j \wedge d\zeta_k \tag{3.23}$$

is a symplectic form, then for

$$H_{\zeta_j} = \sum_{k=1}^{\ell} \alpha_k \frac{\partial}{\partial \zeta_k} \tag{3.24}$$

we have

$$\omega \left(H_{\zeta_j}, \frac{\partial}{\partial \zeta_k} \right) = \delta_{jk} \Rightarrow \sum_{m=1}^{\ell} \Omega_{mk} \alpha_m = \delta_{jk}$$

or (since $\Omega_{mk} = -\Omega_{km}$)

$$\alpha_k = [(-\Omega)^{-1} \delta_j]_k = -(\Omega^{-1})_{kj} \tag{3.25}$$

so that

$$\{\zeta_j, \zeta_k\} = H_{\zeta_j}(\zeta_k) = -(\Omega^{-1})_{kj}. \tag{3.26}$$

For our case if $\zeta = (x_1, \dots, x_{\ell}, y_1, \dots, y_{\ell})$, Ω has the $\ell \times \ell$ block form

$$\Omega = \left(\begin{array}{c|c} U & W \\ \hline -W^t & 0 \end{array} \right) \tag{3.27}$$

for which

$$\Omega^{-1} = \left(\begin{array}{c|c} 0 & -W_t^{-1} \\ \hline W^{-1} & W^{-1} U W_t^{-1} \end{array} \right) \tag{3.28}$$

so (3.21) and (3.22) follow from (3.26).

For $\{y_i, y_j\} = 0$ if and only if $U \equiv 0$, we note $\{y_i, y_j\} = 0$ if and only if $W^{-1} U W^{-1} = 0$ by (3.18).

For the converse, we note that (3.21) and (3.22) plus antisymmetry imply Ω^{-1} has the form (3.28) for some U , and then (3.23) implies (3.19). \square

From this proposition, we can find the basic symplectic two-form in both (a, b) and (μ, x) coordinates:

Theorem 3.6. *The symplectic form defined by (1.3) / (1.4) has the form:*

$$4 \sum_{1 \leq \ell \leq k \leq N-1} a_{\ell}^{-1} da_{\ell} \wedge db_k. \tag{3.29}$$

Remark. (1.3)/(1.5) only define a two-form. That it is symplectic (i.e., $d\omega = 0$) follows from (3.29).

Proof. Since (1.3)/(1.4) say

$$\{a_k, b_k\} = -\{a_k, b_{k+1}\} = \frac{1}{4} a_k,$$

Proposition 3.5 (which, as we noted, does not use that the Poisson bracket obeys the Jacobi identity) says that (recall that $b_1, \dots, b_{N-1}; a_1, \dots, a_{N-1}$ are a set of coordinates)

$$\omega = \sum_{i,j=1}^{N-1} W_{ij} da_i \wedge db_j \tag{3.30}$$

where $(W^{-1})^t$ has the form

$$\begin{pmatrix} \frac{1}{4} a_1 & -\frac{1}{4} a_1 & 0 & \dots \\ 0 & \frac{1}{4} a_2 & -\frac{1}{4} a_2 & \dots \\ \dots & \dots & \dots & \dots \end{pmatrix} = D(1 - N), \tag{3.31}$$

where D is the diagonal matrix $\frac{1}{4} a_k \delta_{kl}$ and $N = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ \dots & \dots & \dots \end{pmatrix}$ is the standard rank $N - 2$ nilpotent on \mathbb{R}^{N-1} . Thus

$$W = [(D(1 - N))^{-1}]^t = D^{-1}(1 + N^t + (N^t)^2 + \dots + (N^t)^{n-1}) \tag{3.32}$$

$$= \begin{pmatrix} 4a_1^{-1} & 0 & 0 & \dots \\ 4a_1^{-1} & 4a_2^{-1} & 0 & \dots \\ 4a_1^{-1} & 4a_2^{-1} & 4a_3^{-1} & \dots \\ \dots & \dots & \dots & \dots \end{pmatrix} \tag{3.33}$$

so (3.30) is (3.29). \square

Theorem 3.7. *The symplectic form defined by (1.3) / (1.4) has the form:*

$$2 \sum_{j=1}^N dx_j \wedge \rho_j^{-1} d\rho_j + \sum_{i,j} U_{ij} dx_i \wedge dx_j \tag{3.34}$$

for some U .

Remarks. (1) By $\sum_{j=1}^N x_j = c$ and $\sum_{j=1}^N \rho_j = 1$, the x 's and ρ 's are not independent, but they still define functions.

(2) One could compute U from (1.12).

Proof. Define for $k = 1, \dots, N - 1$,

$$y_k = \log \left[\frac{\rho_k}{\rho_N} \right]. \tag{3.35}$$

Then for $j = 1, \dots, N - 1$,

$$\{x_j, y_k\} = \frac{1}{2} \delta_{jk} \tag{3.36}$$

since, by (1.11),

$$\begin{aligned} \{x_j, y_k\} &= \{x_j, \log \rho_k\} - \{x_j, \log \rho_n\} \\ &= \left(\frac{1}{2} \delta_{jk} - \rho_j\right) - (-\rho_j). \end{aligned}$$

$\{x_j\}_{j=1}^{N-1}, \{y_j\}_{j=1}^{N-1}$ are local (indeed, global) coordinates since $\sum_{j=1}^n \rho_j = 1$ lets us invert (3.35),

$$\rho_j = \frac{e^{y_j}}{[1 + \sum_{\ell=1}^{N-1} e^{y_\ell}]} \tag{3.37}$$

$$\rho_N = \frac{1}{[1 + \sum_{\ell=1}^{N-1} e^{y_\ell}]} \tag{3.38}$$

so $\{x_j, x_k\} = 0$ and (3.26) plus Proposition 3.5 imply

$$\omega = 2 \sum_{j=1}^{N-1} dx_j \wedge dy_j + \sum_{i,j} U_{ij} dx_i \wedge dx_j \tag{3.39}$$

(3.34) follows if we note that

$$dy_j = \rho_j^{-1} d\rho_j - \rho_N^{-1} d\rho_N$$

and

$$\sum_{j=1}^{N-1} dx_j \wedge d\rho_N = -dx_N \wedge d\rho_N$$

since $\sum_{j=1}^N dx_j = 0$. \square

4. Coefficient stripping for paraorthogonal polynomials

Following [46,47], we use Φ_N and Ψ_n for the first and second kind orthogonal polynomials. Paraorthogonal polynomials are defined [25] by

$$P_N(z; \{\alpha_j\}_{j=0}^{N-2}, \beta) = z\Phi_{N-1}(z) - \bar{\beta}\Phi_{N-1}^*(z), \tag{4.1}$$

where $\beta \in \partial\mathbb{D}$. Second kind paraorthogonal polynomials are defined by

$$Q_N(z; \{\alpha_j\}_{j=0}^{N-2}, \beta) = z\Psi_{N-1}(z) + \bar{\beta}\Psi_{N-1}^*(z), \tag{4.2}$$

where Ψ are the second kind polynomials. These polynomials have been extensively studied recently [4,5,8,20,50,56] and, in particular, the second kind polynomials are introduced in [50, 56]. The symbols P and Q are new but quite natural. These are relevant for the following reason:

Proposition 4.1. *Let*

$$d\mu = \sum_{j=1}^N \mu_j \delta_{z_j} \tag{4.3}$$

be a pure point probability measure on $\partial\mathbb{D}$ with each $\mu_j > 0$. Let $\Phi_0, \Phi_1, \dots, \Phi_N$ be the monic OPUC where, in particular,

$$\Phi_N(z) = \prod_{j=1}^N (z - z_j). \tag{4.4}$$

Then the recursion relations define parameters $\{\alpha_j\}_{j=0}^{N-2}$ and $\alpha_{N-1} = \beta \in \partial\mathbb{D}$ where

$$\beta = (-1)^{N+1} \prod_{j=1}^N \bar{z}_j. \tag{4.5}$$

Here

$$\Phi_N(z, d\mu) = P_N(z; \{\alpha_j\}_{j=0}^{N-2}, \beta). \tag{4.6}$$

Moreover,

$$\begin{aligned} F(z, d\mu) &\equiv \int \frac{e^{i\theta} + z}{e^{i\theta} - z} d\mu(z) \\ &= \sum_{j=1}^N \mu_j \frac{z_j + z}{z_j - z} \end{aligned} \tag{4.7}$$

$$= - \frac{Q_N(z; \{\alpha_j\}_{j=0}^{N-2}, \beta)}{P_N(z; \{\alpha_j\}_{j=0}^{N-2}, \beta)}. \tag{4.8}$$

In particular,

$$Q_N(z) = \sum_{j=1}^N \mu_j (z + z_j) \prod_{\ell \neq j} (z - z_\ell). \tag{4.9}$$

Remarks. (1) Parts of this proof are close to results of Jones et al. [25] as presented in Theorem 2.2.12 of [46].

(2) (4.9) is, of course, the analog of (3.5).

(3) It is known that $\text{Im } F(e^{i\theta})$ is strictly monotone and pure imaginary between poles, so there is a single zero between poles. Thus, (4.8) shows the zeros of P_n and Q_n interlace, a result of [50, 56] obtained by other means.

Proof. If $d\mu$ has the form (4.3), then $L^2(\partial\mathbb{D}, d\mu)$ is N -dimensional, so $\{\Phi_j\}_{j=0}^{N-1}$ span the whole space so $\Phi_N(e^{i\theta}) = 0$ for $d\mu$ a.e. θ . It follows that Φ_N must obey (we use P_N for later purposes, even though we have not yet proven that Φ_N is a P_N)

$$P_N(z) = \prod_{j=1}^N (z - z_j). \tag{4.10}$$

In particular, $\alpha_{N-1} = \overline{-\Phi_N(0)}$ is given by (4.5) and so is a $\beta \in \partial\mathbb{D}$. Szegő recursion thus says (4.1) holds.

Using $*$ for degree N -polynomials, (4.1)/(4.2) imply

$$P_N^* = -\beta P_N, \quad Q_N^* = \beta Q_N \tag{4.11}$$

so

$$- \frac{Q_N}{P_N} = \frac{Q_N^*}{P_N^*} \tag{4.12}$$

and (4.8) is just (3.2.19) of [46] (this is proven for $(\alpha_0, \dots, \alpha_{N-1}, 0, 0 \dots)$ with $|\alpha_{N-1}| < 1$, but by taking $\alpha_{N-1} \rightarrow \partial\mathbb{D}$, one gets $F = Q_N^*/P_N^*$). (4.7), (4.8) and (4.10) imply (4.9). \square

Since we are about to consider changes in N , we shift notation from N to n . Since P_n, Q_n are defined via a boundary condition on α_{n-1} , if we want to prove something inductively, we need to “strip from the front,” that is, remove α_0 . In [46,47] we called this coefficient stripping. For OPUC, this involves the formula for $\Phi_n(z; \{\alpha_j\}_{j=0}^{n-1})$ and $\Phi_n^*(z; \{\alpha_j\}_{j=0}^{n-1})$ in terms of α_0 and $\Phi_{n-1}(z; \{\alpha_{j+1}\}_{j=0}^{n-2})$ and $\Phi_{n-1}^*(z; \{\alpha_{j+1}\}_{j=0}^{n-2})$. Quite remarkably (and conveniently for use in Sections 5 and 6), coefficient stripping for P_n, Q_n involves another P, Q pair even though P, Q are (Φ, Ψ) mixed! It turns out simpler to state things in terms of

$$C_n(z; \{\alpha_j\}_{j=0}^{n-2}, \beta) = \frac{(P_n + Q_n)}{2}, \tag{4.13}$$

$$S_n(z; \{\alpha_j\}_{j=0}^{n-2}, \beta) = \frac{(P_n - Q_n)}{2} \tag{4.14}$$

which we name analogously to cosh and sinh.

Theorem 4.2. *Let $\beta \in \partial\mathbb{D}$ be fixed and $\{\alpha_j\}_{j=0}^{n-2} \in \mathbb{D}^{n-1}$. Let C_n denote $C_n(z; \{\alpha_j\}_{j=0}^{n-2}, \beta)$, and similarly for S_n . Define C_{n-1} by*

$$C_{n-1} = C_{n-1}(z; \{\alpha_{j+1}\}_{j=0}^{n-3}, \beta) \tag{4.15}$$

(with α_0 removed), and similarly for S_{n-1} . Then

$$C_n(z) = z(C_{n-1}(z) - \alpha_0 z S_{n-1}(z)), \tag{4.16}$$

$$S_n(z) = -\bar{\alpha}_0 C_{n-1}(z) + S_{n-1}(z). \tag{4.17}$$

Proof. As usual, let

$$\Lambda(\alpha, z) = \begin{pmatrix} z & -\bar{\alpha} \\ -\alpha z & 1 \end{pmatrix} \tag{4.18}$$

so with $\delta_+ = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \in \mathbb{C}^2, \delta_- = \begin{pmatrix} 0 \\ 1 \end{pmatrix} \in \mathbb{C}^2$, Szegő recursion says

$$\begin{pmatrix} \Phi_{n-1} \\ \Phi_{n-1}^* \end{pmatrix} = \Lambda(\alpha_{n-2}, z) \dots \Lambda(\alpha_0, z)(\delta_+ + \delta_-) \tag{4.19}$$

$$\begin{pmatrix} \Psi_{n-1} \\ -\Psi_{n-1}^* \end{pmatrix} = \Lambda(\alpha_{n-2}, z) \dots \Lambda(\alpha_0, z)(\delta_+ - \delta_-). \tag{4.20}$$

Let $B(\beta) = \begin{pmatrix} z & 0 \\ 0 & -\beta \end{pmatrix}$ and conclude

$$P_n = \langle (\delta_+ + \delta_-), (B(\beta)\Lambda_{n-2}(\alpha_{n-2}, z) \dots \Lambda(\alpha_0, z))(\delta_+ + \delta_-) \rangle. \tag{4.21}$$

Q_n is similar with the rightmost $(\delta_+ + \delta_-)$ replaced by $(\delta_+ - \delta_-)$. Thus, C_n, S_n correspond to using δ_+ and δ_- , and we have that

$$\begin{pmatrix} C_n \\ S_n \end{pmatrix} = \Lambda(\alpha_0, z)^t \Lambda(\alpha_1, z)^t \dots \Lambda(\alpha_{n-1}, z)^t B(\beta)^t \begin{pmatrix} 1 \\ 1 \end{pmatrix} \tag{4.22}$$

which implies

$$\begin{pmatrix} C_n \\ S_n \end{pmatrix} = \Lambda(\alpha_0, z)^t \begin{pmatrix} C_{n-1} \\ S_{n-1} \end{pmatrix} \tag{4.23}$$

which is (4.16)–(4.17). \square

Remark. One might think the transposes in (4.22) should be adjoints, but the shift of δ_{\pm} from the right to the left side of the inner product introduces a complex conjugate.

While we would not need it, we note the induced formula for P_n, Q_n :

Theorem 4.3. *If $P_n(z) = P_n(z; \{\alpha_j\}_{j=0}^{n-2}, \beta)$ and $P_{n-1}(z) = P_n(z; \{\alpha_{j+1}\}_{j=0}^{n-3}, \beta)$, and similarly for Q_n, Q_{n-1} , then*

$$P_n = \frac{1}{2}(z - \bar{\alpha}_0 + 1 - \alpha_0 z)P_{n-1} + \frac{1}{2}(z - \bar{\alpha}_0 - 1 + \alpha_0 z)Q_{n-1}, \tag{4.24}$$

$$Q_n = \frac{1}{2}(z + \bar{\alpha}_0 + 1 + \alpha_0 z)Q_{n-1} + \frac{1}{2}(z + \bar{\alpha}_0 - 1 - \alpha_0 z)P_{n-1}. \tag{4.25}$$

Proof. This is immediate from (4.23), together with

$$\begin{pmatrix} P_n \\ Q_n \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} \begin{pmatrix} C_n \\ S_n \end{pmatrix}$$

$$\begin{pmatrix} C_{n-1} \\ S_{n-1} \end{pmatrix} = \frac{1}{2} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} \begin{pmatrix} P_{n-1} \\ Q_{n-1} \end{pmatrix}$$

and a calculation of

$$\frac{1}{2} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix} \Lambda^t(\alpha_0, z) \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}. \quad \square$$

The algebraic simplicity of the recursion of (C_n, S_n) relative to (P_n, Q_n) is reminiscent of Schur vs. Carathéodory functions (see Section 3.4 of [46]). This is not a coincidence. Note that by (4.8)

$$\begin{aligned} \frac{C_n}{S_n} &= \frac{P_n + Q_n}{P_n - Q_n} = \frac{1 - [(-Q_n)/P_n]}{1 + [(-Q_n)/P_n]} \\ &= \frac{1 - F}{1 + F} \end{aligned} \tag{4.26}$$

so

$$f(z) = -\frac{z^{-1}C_n}{S_n}. \tag{4.27}$$

Note that S_n is a polynomial of degree $n - 1$ and C_n is a polynomial of degree n vanishing at zero, so (4.27) is a ratio of polynomials of degree $n - 1$. $-S_n$ and $z^{-1}C_n$ are thus essentially paraversions of the Wall polynomials. Indeed, by the Pinter–Nevai formula [46, Theorem 3.2.10],

$$z^{-1}C_n(z) = zB_{n-2}^*(z) + \bar{\beta}A_{n-2}(z),$$

$$S_n(z) = -zA_{n-2}^*(z) - \bar{\beta}B_{n-2}(z).$$

5. Poisson brackets of first and second kind OPUC

Our goal in this section is to prove (1.18) and (1.19). Once we translate to C, S , the induction will be even simpler than in the OPRL because C_{n-1}, S_{n-1} are only dependent on $\{\alpha_{j+1}\}_{j=0}^{n-3}$ and so have zero Poisson bracket with α_0 , so there are no analogs of terms like (2.1). Since we are varying the index, we use n in place of N . We will begin by showing (1.18)/(1.19) is equivalent to

$$\{C_n(z), C_n(w)\} = 0 = \{S_n(z), S_n(w)\}, \tag{5.1}$$

$$\{C_n(z), S_n(w)\} = -i \left(\frac{C_n(z)wS_n(w) - C_n(w)zS_n(z)}{z - w} \right). \tag{5.2}$$

Proposition 5.1. (1.18) and the symmetry of $\{P_n(z), Q_n(w)\}$ under $z \leftrightarrow w$ implies (5.1) and

$$\{C_n(z), S_n(w)\} = -\frac{1}{2} \{P_n(z), Q_n(w)\}. \tag{5.3}$$

Similarly, (5.1) and the symmetry of $\{C_n(z), S_n(w)\}$ under $z \leftrightarrow w$ implies (1.18) and (5.3).

Proof. Given the antisymmetry of the Poisson bracket, the symmetry of $\{P_n(z), Q_n(w)\}$ can be written

$$\{P_n(z), Q_n(w)\} + \{Q_n(z), P_n(w)\} = 0$$

from which (1.18) implies (5.1), and conversely, (5.1) and the symmetry of $\{C_n(z), S_n(w)\}$ implies (1.18).

The antisymmetry plus (1.18) also shows

$$\{C_n(z), S_n(w)\} = \frac{1}{4} 2\{Q_n(z), P_n(w)\}$$

which is (5.3), and the converse is similar. \square

Next, we do a calculation showing the equality of the left side of (5.2) and of (1.19) using $P_n = C_n + S_n$ and $Q_n = C_n - S_n$:

$$\begin{aligned} &\text{RHS of (1.19)} \\ &= -\frac{i}{2} \left[\left(\frac{z+w}{z-w} \right) [-2C_n(z)S_n(w) + 2C_n(w)S_n(z)] + 2C_n(z)S_n(w) + 2C_n(w)S_n(z) \right] \\ &= i \left[\left\{ \left(\frac{z+w}{z-w} \right) - 1 \right\} C_n(z)S_n(w) - \left\{ \left(\frac{z+w}{z-w} \right) + 1 \right\} C_n(w)S_n(z) \right] \\ &= i \left[\frac{2C_n(z)wS_n(w) - 2C_n(w)zS_n(z)}{z - w} \right] \\ &= -2 [\text{RHS of (5.2)}] \end{aligned} \tag{5.4}$$

consistent with (5.1).

Thus, (1.18)/(1.19) is equivalent to (5.1)/(5.2), which we proceed to prove inductively, starting with $n = 1$. We let $G_n(z, w)$ denote the right side of (5.2).

Proposition 5.2. We have that

$$C_1(z) = z, \quad S_1(z) = -\bar{\beta}. \tag{5.5}$$

Thus, $G_1(z, w) = 0$ and (5.1), (5.2) hold for $n = 1$.

Proof. Since

$$P_1(z) = z - \bar{\beta}, \quad Q_1(z) = z + \bar{\beta}. \tag{5.6}$$

(5.5) is immediate, and thus,

$$G_1(z, w) = i\bar{\beta} \left(\frac{zw - wz}{z - w} \right) = 0.$$

Since C_1, S_1 are independent of $\{\alpha_j\}$, all Poisson brackets are zero and (5.1), (5.2) hold for $n = 1$. \square

Remark. It is an interesting exercise to prove

$$(P_1(z)Q_1(w) - P_1(w)Q_1(z)) \frac{z + w}{z - w} - Q_1(z)Q_1(w) + P_1(z)P_1(w)$$

is 0 from (5.6), and explain why $-QQ$ appears with a minus sign.

Lemma 5.3. G_n and G_{n-1} are related by

$$G_n(z, w) = z\rho_0^2 G_{n-1}(z, w) - i\rho_0^2 z S_{n-1}(z) C_{n-1}(w). \tag{5.7}$$

Proof. By (4.16), (4.17), and the fact that

$$\mathcal{B}(f, g)(z, w) = \frac{f(z)g(w) - f(w)g(z)}{z - w} \tag{5.8}$$

is zero if $f = g$:

$$\begin{aligned} G_n(z, w) &= -i\mathcal{B}(C_n, X S_n)(z, w) \\ &= -i\mathcal{B}(X C_{n-1} - \alpha_0 X S_{n-1}, -\bar{\alpha}_0 X C_{n-1} + X S_{n-1}) \\ &= -i[\mathcal{B}(X C_{n-1}, X S_{n-1}) + |\alpha_0|^2 \mathcal{B}(X S_{n-1}, X C_{n-1})] \\ &= -i\rho_0^2 \mathcal{B}(X C_{n-1}, X S_{n-1}), \end{aligned} \tag{5.9}$$

where we use Xf for the function

$$(Xf)(z) = zf(z)$$

and (5.9) using antisymmetry of \mathcal{B} in f and g and $1 - |\alpha_0|^2 = \rho_0^2$.

Next, note

$$\begin{aligned} \mathcal{B}(Xf, g)(z, w) &= \frac{zf(z)g(w) - wf(w)g(z)}{z - w} \\ &= z \left(\frac{f(z)g(w) - f(w)g(z)}{z - w} \right) + \frac{(z - w)}{z - w} f(w)g(z) \\ &= z\mathcal{B}(f, g)(z, w) + g(z)f(w) \end{aligned}$$

to conclude

$$-i\mathcal{B}(X C_{n-1}, X S_{n-1}) = zG_{n-1}(z, w) - izS_{n-1}(z)C_{n-1}(w) \tag{5.10}$$

(5.9) and (5.10) imply (5.7). \square

Theorem 5.4. (5.1), (5.2) hold for all n . (1.18), (1.19) hold for all n .

Proof. As noted after Proposition 5.1 and (5.4), (1.18)/(1.19) is equivalent to (5.1)/(5.2). So, by Proposition 5.2 and induction, we need only prove

$$(5.1)–(5.2) \text{ for } n - 1 \Rightarrow (5.1)–(5.2) \text{ for } n. \tag{5.11}$$

By (4.16) and (4.17), $\{\alpha_0, \alpha_0\} = \{\bar{\alpha}_0, \bar{\alpha}_0\} = 0$ the symmetry of $G_{n-1}(z, w)$ in $z \leftrightarrow w$, and the independence of S_{n-1}, C_{n-1} on $\alpha_0, \bar{\alpha}_0$, we have

$$(5.1)–(5.2) \text{ for } n - 1 \Rightarrow (5.1) \text{ for } n.$$

(Notice how much simpler this is than the analog for OPRL!)

On the other hand, by (4.16) and (4.17),

$$\begin{aligned} \{C_n(z), S_n(w)\} &= z\{C_{n-1}(z) - \alpha_0 S_{n-1}(z), -\bar{\alpha}_0 C_{n-1}(w) + S_{n-1}(w)\} \\ &= z\rho_0^2 G_{n-1}(z, w) + \{\alpha_0, \bar{\alpha}_0\} z S_{n-1}(z) C_{n-1}(w) \\ &= z\rho_0^2 G_{n-1}(z, w) - i\rho_0^2 z S_{n-1}(z) C_{n-1}(w) \\ &= G_n(z, w). \end{aligned} \tag{5.12}$$

$$(5.13)$$

(5.12) uses (1.17) and (5.13) follows from (5.7). We have thus proven (5.2) for n .

6. Fundamental Poisson brackets for OPUC

In this section, we will establish (1.23). Let $\{z_j\}_{j=1}^n$ be the zeros of $P_n(z)$, θ_j given by (1.21) and μ_j by (1.20). Unlike the OPRL case where $x_1 < \dots < x_n$ allowed global variables, these are only defined locally, both because of the z 's not having a unique initial z_1 and because of 2π ambiguities in θ_j .

Throughout this section, we fix $\beta \in \partial\mathbb{D}$ and consider the $2n - 2$ -dimensional manifold \mathbb{D}^{n-1} of Verblunsky coefficients $\{\alpha_j\}_{j=0}^{n-2}$ and $\alpha_{n-1} = \beta$. The associated finite CMV matrix C has

$$\det(C) = (-1)^{n-1} \beta. \tag{6.1}$$

The associated spectral measure given by (1.21) thus has

$$\prod_{j=1}^n z_j = (-1)^{n-1} \beta \tag{6.2}$$

or

$$\sum_{j=1}^n \theta_j = \arg((-1)^{n-1} \beta) \tag{6.3}$$

and, of course,

$$\sum_{j=1}^n \mu_j = 1 \tag{6.4}$$

so as local coordinates $\{\theta_j\}_{j=1}^{n-1} \cup \{\mu_j\}_{j=1}^{n-1}$ coordinatizes \mathbb{D}^{n-1} . β never appears explicitly below—this is not surprising since a rotation can move β to 1.

As usual, one can replace $\text{Re } \alpha, \text{Im } \alpha$ by “independent” coordinates $\alpha, \bar{\alpha}$ with

$$d\alpha = d(\text{Re } \alpha) + i d(\text{Im } \alpha) \quad d\bar{\alpha} = d(\text{Re } \alpha) - i d(\text{Im } \alpha) \tag{6.5}$$

$$\frac{\partial}{\partial \alpha} = \frac{1}{2} \left[\frac{\partial}{\partial \text{Re } \alpha} + \frac{1}{i} \frac{\partial}{\partial \text{Im } \alpha} \right] \quad \frac{\partial}{\partial \bar{\alpha}} = \frac{1}{2} \left[\frac{\partial}{\partial \text{Re } \alpha} - \frac{1}{i} \frac{\partial}{\partial \text{Im } \alpha} \right]. \tag{6.6}$$

Thus, the Poisson bracket defined by (1.17) can be defined in general by

$$\{f, g\} = \sum_{j=1}^n i \rho_j^2 \left(\frac{\partial f}{\partial \bar{\alpha}_j} \frac{\partial g}{\partial \alpha_j} - \frac{\partial f}{\partial \alpha_j} \frac{\partial g}{\partial \bar{\alpha}_j} \right). \tag{6.7}$$

We will denote

$$d^2 \alpha_j = d(\text{Re } \alpha_j) \wedge d(\text{Im } \alpha_j) \tag{6.8}$$

$$= \frac{i}{2} d\alpha_j \wedge d\bar{\alpha}_j. \tag{6.9}$$

Proposition 6.1. For OPUC,

$$\{P_n(z), P_n(w)\}|_{z=z_k, w=z_j} = -z_k z_j \{\theta_k, \theta_j\} \prod_{\ell \neq k} (z_k - z_\ell) \prod_{\ell \neq j} (z_j - z_\ell). \tag{6.10}$$

In particular,

$$\{\theta_k, \theta_j\} = 0. \tag{6.11}$$

Proof. Since $P_n(z) = \prod_{\ell=1}^n (z - z_\ell)$, we have

$$\text{LHS of (6.10)} = \{z_k, z_j\} \prod_{\ell \neq k} (z_k - z_\ell) \prod_{\ell \neq j} (z_j - z_\ell).$$

(6.10) follows if we note

$$\{e^{i\theta_k}, e^{i\theta_j}\} = (i)^2 e^{i\theta_k} e^{i\theta_j} \{\theta_k, \theta_j\}.$$

(1.18) then implies (6.11). \square

Proposition 6.2. For OPUC,

$$\{P_n(z), Q_n(w)\}|_{z=z_j, w=z_k} = -i \{\theta_j, \mu_k\} 2z_j z_k \prod_{\ell \neq k} (z_j - z_\ell) \prod_{\ell \neq j} (z_k - z_\ell) \tag{6.12}$$

while

$$\begin{aligned} & \lim_{\substack{z \rightarrow z_j \\ z \neq z_j}} \text{RHS of (1.19)}|_{w=z_k} \\ &= -\frac{i}{2} \{2z_k \mu_k \delta_{jk} (z_j + z_k) - 2z_j 2z_k \mu_j \mu_k\} \prod_{\ell \neq j} (z_j - z_\ell) \prod_{\ell \neq k} (z_k - z_\ell). \end{aligned} \tag{6.13}$$

In particular,

$$\{\theta_j, \mu_k\} = \mu_j \delta_{jk} - \mu_j \mu_k. \tag{6.14}$$

Proof. Given (4.9), setting $z = z_j$, $w = z_k$ picks out the $\{z_j, \mu_k\}$ term, that is,

$$\text{LHS of (6.12)} = \{-z_j, \mu_k\} \left[\prod_{\ell \neq j} (z_j - z_\ell) \right] \left[2z_k \prod_{\ell \neq k} (z_k - z_\ell) \right],$$

where $2z_k$ is $(w + z_k)|_{w=z_k}$. Since

$$\{-z_j, \mu_k\} = -i z_j \{\theta_j, \mu_k\}$$

we obtain (6.12).

Setting $w = z_k$ in the right side of (1.19) gives (using $P(z_k) = 0$, $Q(z_k) = \mu_k(2z_k) \prod_{\ell \neq k} (z_k - z_\ell)$)

$$\text{RHS of (1.19)}|_{w=z_k} = -\frac{i}{2} \left\{ \frac{P(z)}{z - z_k} \mu_k(z + z_k) - Q(z) \mu_k \right\} (2z_k) \prod_{\ell \neq k} (z_k - z_\ell).$$

Since $P(z)/(z - z_k) = \prod_{\ell \neq k} (z - z_\ell)$, the first term is zero as $z \rightarrow z_j$ if $j \neq k$, and otherwise is $\prod_{\ell \neq j} (z_j - z_\ell)$. This yields (6.13). (6.13) plus (6.12) yields (6.14). \square

From $\{Q(z), Q(w)\} = 0$, one can derive the formula for $\{\mu_j, \mu_k\}$. We do not provide details.

Finally, for this section, we compute the symplectic form, ω , in $d\alpha$, and $d\theta_j, d\mu_j$ coordinates.

The $d\alpha$ coordinate is especially simple since in those coordinates, the Poisson bracket is a product structure. By (1.17) and $\text{Re } \alpha_j = \frac{1}{2}(\alpha_j + \bar{\alpha}_j)$, $\text{Im } \alpha_j = \frac{1}{2i}(\alpha_j - \bar{\alpha}_j)$, we get

$$\{\text{Re } \alpha_j, \text{Im } \alpha_k\} = \frac{\rho_j^2}{2}$$

so by the $U = 0$ case of Proposition 3.5 (and (6.9)):

Proposition 6.3. *The symplectic form for OPUC with $\alpha_{n-1} = \beta \in \partial\mathbb{D}$ fixed is*

$$\omega = \sum_{j=0}^{n-2} \frac{2}{\rho_j^2} d(\text{Re } \alpha_j) \wedge d(\text{Im } \alpha_j) \tag{6.15}$$

$$= \sum_{j=0}^{n-2} \frac{i}{\rho_j^2} (d\alpha_j \wedge d\bar{\alpha}_j). \tag{6.16}$$

The calculation for θ, μ coordinates is essentially the same as for OPRL with x, ρ coordinates since $\sum_{j=1}^n \theta_j$ is constant. Since the $\frac{1}{2}$ in (1.11) is missing from (1.23), the 2 in (3.34) is absent and we get:

Proposition 6.4. *The symplectic form for OPUC in terms of $\{\mu_j, \theta_j\}$ is*

$$\omega = \sum_{j=1}^N d\theta_j \wedge \mu_j^{-1} d\mu_j + \sum_{j,k} U_{jk} d\theta_j \wedge d\theta_k \tag{6.17}$$

for some U .

7. Application 1: Jacobians of coordinate changes

This is the first of three sections in which we present applications of the fundamental Poisson brackets (1.11) for OPRL and (1.23) for OPUC. Those in this section and the next are not new, but we include them here because the literature on these issues is not always so clear nor is it always emphasized that all that is needed are the fundamental Poisson brackets.

Given two local coordinates x_1, \dots, x_m and y_1, \dots, y_m , their Jacobian $|\frac{\partial x}{\partial y}|$ is defined by

$$\left| \frac{\partial(x)}{\partial(y)} \right| \equiv \left| \det \left(\frac{\partial x_j}{\partial y_k} \right) \right| \tag{7.1}$$

named because

$$d^n x = \left| \frac{\partial(x)}{\partial(y)} \right| d^n y \tag{7.2}$$

relates the local volume elements. Of course, differential forms are ideal for this, $dx_j = \sum \frac{\partial x_j}{\partial y_k} dy_k$ means

$$dx_1 \wedge \dots \wedge dx_n = \det \left(\frac{\partial x_j}{\partial y_k} \right) dy_1 \wedge \dots \wedge dy_n. \tag{7.3}$$

Our goal here is to compute the two Jacobians $|\frac{\partial(a,b)}{\partial(x,\rho)}|$ for OPRL and $|\frac{\partial(\text{Re } \alpha, \text{Im } \alpha)}{\partial(\theta, \mu)}|$ for OPUC. It appears that these were first found for OPRL by Dumitriu–Edelman [12], and motivated by that, for OPUC by Killip–Nenciu [31]. Their argument was indirect and Killip–Nenciu asked if there were not a direct calculation. This was provided by Forrester–Rains [16] using forms and the continued fraction expansions of resolvents. Their argument is not unrelated to the one below, but is lacking the connection to Poisson brackets and we feel is more involved. In [32], Killip–Nenciu remark that Deift (unpublished) explained to them how to go from the symplectic form for Jacobi parameters rewritten in terms of (x, ρ) to $|\frac{\partial(a,b)}{\partial(x,\rho)}|$ and they then do the analogous OPUC calculation. As they remark, this is buried at the end of a long paper, and the fact that only the form of Poisson brackets is involved is obscure. We present this idea here to make the OPRL argument explicit and to present the OPUC argument without a need for Lie algebra actions.

Theorem 7.1. *On the $2n - 2$ -dimensional manifold of Jacobi parameters with $\sum_{j=1}^N b_j$ fixed, use $\{a_j, b_j\}_{j=1}^{N-1}$ and $\{x_j, \rho_j\}_{j=1}^{N-1}$ as coordinates. Then*

$$\left| \frac{\partial(a, b)}{\partial(x, \rho)} \right| = \left[\frac{2^{-(N-1)} \prod_{j=1}^{N-1} a_j}{\prod_{j=1}^N \rho_j} \right]. \tag{7.4}$$

If one considers the $2N - 1$ -dimensional manifold without $\sum_{j=1}^N b_j$ fixed (equivalently, without $\sum_{j=1}^N x_j$ fixed), then (7.4) is still true, but where the Jacobian is from $(a_1, \dots, a_{N-1}, b_1, \dots, b_N)$ to $(x_1, \dots, x_N, \rho_1, \dots, \rho_{N-1})$ rather than from $(a_1, \dots, a_{N-1}, b_1, \dots, b_{N-1})$ to $(x_j, \dots, x_{N-1}, \rho_j, \dots, \rho_{N-1})$.

Proof. Let ω be the symmetric form on the $2N - 2$ -dimensional manifold determined by (1.3)–(1.4). By (3.29), the $(N - 1)$ -fold wedge product

$$\omega \wedge \dots \wedge \omega = 4^{N-1}(N - 1)! \prod_{j=1}^{N-1} a_j^{-1} da_1 \wedge db_1 \wedge \dots \wedge da_{N-1} \wedge db_{N-1}. \tag{7.5}$$

By (3.34), noting all terms with at least one $dx_k \wedge dx_j$ must have a repeated x and so vanish,

$$\begin{aligned} \omega \wedge \dots \wedge \omega &= 2^{N-1}(N - 1)! \sum_{j=1}^N \bigwedge_{k \neq j} \rho_k^{-1} (dx_k \wedge d\rho_k) \\ &= 2^{N-1}(N - 1)! \left[\sum_{j=1}^N \left(\prod_{k \neq j} \rho_k^{-1} \right) \right] dx_1 \wedge d\rho_1 \wedge \dots \wedge dx_{N-1} \wedge d\rho_{N-1} \end{aligned} \tag{7.6}$$

$$= 2^{N-1}(N - 1)! \left(\prod_{k=1}^N \rho_k^{-1} \right) dx_1 \wedge d\rho_1 \wedge \dots \wedge dx_{N-1} \wedge d\rho_{N-1}. \tag{7.7}$$

Here (7.6) is obtained from

$$dx_N = - \sum_{j \neq N} dx_j, \quad d\rho_N = - \sum_{j \neq N} d\rho_j$$

and (7.7) uses

$$\sum_{j=1}^N \prod_{k \neq j} \rho_k^{-1} = \prod_{k=1}^N \rho_k^{-1} \sum_{j=1}^N \rho_j = \prod_{k=1}^N \rho_k^{-1}. \tag{7.8}$$

(7.4) is immediate from the equalities of the right side of (7.5) and (7.7).

To get the results for the $2N - 1$ -dimensional manifold, we note that for each fixed $\sum_{j=1}^N x_j$, we have

$$\text{RHS of (7.7)} = \text{RHS of (7.5)}. \tag{7.9}$$

Since $\sum_{j=1}^N x_j = \sum_{j=1}^N b_j$, we can wedge the right side of (7.5) with $db_1 + \dots + db_N$ and of (7.7) by $dx_1 + \dots + dx_N$ and so obtain the analog of (7.9) but with the extra variable. \square

Theorem 7.2. On the $2N - 2$ -dimensional manifold where $\alpha_{N-1} = \beta \in \partial\mathbb{D}$ is fixed and

$$\alpha_j = u_j + i v_j$$

$(u, v \in \mathbb{R})$, use $\{u_j, v_j\}_{j=0}^{N-2}$ and $\{\theta_j, \mu_j\}_{j=1}^{N-1}$ as coordinates. Then

$$\left| \frac{\partial(u, v)}{\partial(\theta, \mu)} \right| = \frac{2^{-(N-1)} \prod_{j=1}^N \rho_j^2}{\prod_{j=1}^N \mu_j}. \tag{7.10}$$

If we consider the $2N - 2$ -dimensional manifold without fixing β and use coordinates $\{u_j, v_j\}_{j=0}^{N-2} \cup \{\psi\}$ where $\beta = e^{i\psi}$ and $\{\theta_j\}_{j=1}^N \cup \{\mu_j\}_{j=1}^{N-1}$, then

$$\left| \frac{\partial(u, v, \psi)}{\partial(\theta, \mu)} \right| = \text{RHS of (7.10)}. \tag{7.11}$$

Proof. By (6.15), the $N - 1$ -fold product

$$\omega \wedge \dots \wedge \omega = (N - 1)! 2^{N-1} \left(\prod_{j=1}^{N-2} \rho_j^{-2} \right) du_0 \wedge dv_0 \wedge \dots \wedge du_{N-2} \wedge dv_{N-2}, \tag{7.12}$$

where, by (6.17) and the same calculation that led to (7.7) (i.e., (7.8) with ρ replaced by μ),

$$\omega \wedge \dots \wedge \omega = (N - 1)! \prod_{j=1}^N \mu_j^{-1} d\theta_1 \wedge d\mu_1 \wedge \dots \wedge d\theta_{N-1} \wedge d\mu_{N-1}. \tag{7.13}$$

(7.10) is immediate and (7.11) then follows from

$$d\psi = \sum_{j=1}^N d\theta_j$$

on account of (6.4). \square

Note that in comparing (7.4) and (7.10) with Forrester–Rains [16] and Dumitriu–Edelman [12], one needs to bear in mind that their q_j are related to ρ_j (and μ_j) by

$$\rho_j = q_j^2 \tag{7.14}$$

and thus

$$\frac{d\rho_j}{\rho_j} = \frac{2dq_j}{q_j} \tag{7.15}$$

and therefore

$$\frac{d\rho_1 \dots d\rho_{N-1}}{\rho_1 \dots \rho_N} = 2^{N-1} \frac{1}{q_N} \frac{dq_1 \dots dq_{N-1}}{q_1 \dots q_N} \tag{7.16}$$

so they have no $2^{-(N-1)}$ in (7.4) but have a $2^{(N-1)}$ in (7.10). There is an extra factor of $1/q_N$ (which is in [16] but missing in [12] due to the fact that their $d^{N-1}q$ of [12] is the measure on a sphere, not Euclidean measure).

8. Application 2: Generalized Toda and Schur flows

In this section, we want to show how (1.11) allows the explicit solution of flows like (1.6) and (1.7) in the ρ (resp. μ) variables. As mentioned earlier, these results are not new. Theorem 8.1 appears in [10] and many times earlier. Theorem 8.3 appears already in [32] as their Corollary 6.5.

Theorem 8.1. Let f be a real-valued C^1 function on \mathbb{R} and let

$$H = \sum_{j=1}^N f(x_j) \tag{8.1}$$

and let a, b solve (with $\dot{} = d/dt$)

$$\dot{a}_k = \{H, a_k\}, \quad \dot{b}_k = \{H, b_k\} \tag{8.2}$$

for some initial conditions for the Jacobi parameters. Then $\{a_k(t), b_k(t)\}_{k=1}^N$ are the Jacobi parameters associated to

$$x_j(t) = x_j(0), \quad \rho_j(t) = \frac{e^{1/2tf'(x_k)} \rho_j(0)}{\sum_{k=1}^N e^{1/2tf'(x_k)} \rho_k(0)}. \tag{8.3}$$

Remarks. (1) Since all that matters is f at the x_j , we can restrict to polynomials where if $f(x) = x^m$, then $H = \text{Tr}(J^m)$.

(2) $\{x_j, x_k\} = 0$ implies all $\text{Tr}(J^m)$ are constants of the motion.

Example 8.2. The Toda flow (1.6)/(1.7) corresponds to $H = 2 \text{Tr}(J^2)$, so $f(x) = 2x^2$ and $f'(x) = 4x$, and (8.3) becomes

$$\rho_j(t) = \frac{e^{2tx_j} \rho_j(0)}{\sum_{k=1}^N e^{2tx_k} \rho_k(0)} \tag{8.4}$$

which is well known to solve Toda [37,9].

Proof. The flow is generated by the vector field $\{H, \cdot\}$ so

$$\dot{x}_j = \{H, x_j\}, \quad \dot{\rho}_j = \{H, \rho_j\}. \tag{8.5}$$

Since $\{x_j, x_k\} = 0$, (8.5) implies $\dot{x}_j = 0$, that is, $x_j(t) = x_j(0)$.

By (1.11), if $j \neq N$,

$$\begin{aligned} \left\{ x_k, \frac{\rho_j}{\rho_N} \right\} &= \rho_N^{-1} \{x_k, \rho_j\} - \frac{\rho_j}{\rho_N^2} \{x_j, \rho_N\} \\ &= \frac{1}{2} \rho_N^{-1} (\delta_{kj} \rho_j - \rho_j \rho_k) - \frac{1}{2} \rho_N^{-1} (\delta_{kN} \rho_j - \rho_j \rho_k) \\ &= \frac{1}{2} (\delta_{kj} - \delta_{kN}) \frac{\rho_j}{\rho_N}. \end{aligned} \tag{8.6}$$

Thus

$$\frac{d}{dt} \left(\frac{\rho_j}{\rho_N} \right) = \frac{1}{2} [f'(x_j) - f'(x_N)] \frac{\rho_j}{\rho_N}. \tag{8.7}$$

Defining $y_k(t)$ by (3.35), we get

$$y_k(t) = y_k(0) + \frac{1}{2} t (f'(x_j) - f'(x_N))$$

which by (3.37) yields (8.3). \square

Theorem 8.3. Let f be a real-valued C^1 function on $\partial\mathbb{D}$

$$H = \sum_{j=1}^N f(e^{i\theta_j}) \tag{8.8}$$

and let α solve

$$\dot{\alpha}_j = \{H, \alpha_j\} \tag{8.9}$$

with some initial conditions $|\alpha_j(0)| < 1$ and boundary conditions $\alpha_{-1} = -1, \alpha_{N-1} = \beta$. Then $\{\alpha_j(t)\}_{j=1}^{N-2} \cup \{\alpha_{N-1} = \beta\}$ are the Verblunsky coefficients associated to the N -point measure with parameters

$$\theta_j(t) = \theta_j(0), \quad \mu_j(t) = \frac{e^{tg(e^{i\theta_j})} \mu_j(0)}{\sum_{k=1}^N e^{tg(e^{i\theta_k})} \mu_k(0)}, \tag{8.10}$$

where g is the function given by

$$g(e^{i\theta}) = \frac{d}{d\theta} f(e^{i\theta}). \tag{8.11}$$

Example 8.4. If $f(e^{i\theta}) = 2 \sin \theta$, then $H = \frac{1}{i} \text{Tr}(\mathcal{C} - \mathcal{C}^{-1})$ and (8.9) is the Schur flow [1,2,13, 21,48]

$$\dot{\alpha}_j = \rho_j^2 (\alpha_{j+1} - \alpha_{j-1}) \tag{8.12}$$

g is $2 \cos \theta$ and (8.10) becomes

$$\mu_j(t) = \frac{e^{2t \cos(\theta_j)} \mu_j(0)}{\sum_{k=1}^N e^{2t \cos(\theta_k)} \mu_k(0)}. \tag{8.13}$$

Proof. As above, $\{\theta_j, \theta_k\} = 0$ implies $\theta_j(t) = \theta_j(0)$. The calculation of $\mu_j(t)/\mu_N(t)$ is identical to the one in Theorem 8.1 except there is no factor of $\frac{1}{2}$ and $df(x)/dx$ is replaced by $df(e^{i\theta})/d\theta$. \square

By taking suitable limits, we get that certain measures on \mathbb{R} (with all moments finite) as initial conditions for difference equations on $\{a_j, b_j\}_{j=1}^\infty$ or $\{\alpha_j\}_{j=1}^\infty$ can be solved by

$$\frac{e^{tf'(x)} d\rho_{t=0}(x)}{\int e^{tf'(x)} d\rho_{t=0}(x)}.$$

See, for example, the discussion in [48] and references therein.

9. Differential equations for OPRL

We have just seen that symplectic flows are naturally defined on Jacobi parameters. On the level of measures, they are solved by (8.3). Here, we want to consider the differential equations

on OPs induced by these flows. In terms of (8.3), we will suppose f is a polynomial; explicitly,

$$\frac{1}{2} f'(x) = \sum_{j=1}^k c_j x^j. \tag{9.1}$$

We drop the c_0 term since it drops out of $d\rho_t$. Here is our main result:

Theorem 9.1. *Under the map,*

$$d\rho_t(x) = \frac{e^{1/2f'(x)t} d\rho_0}{\int e^{1/2f'(x)t} d\rho_0} \tag{9.2}$$

with f' given by (9.1), the monic orthogonal polynomials $P_n(x; d\rho_t) = P_n(x; t)$ obey

$$\dot{P}_n(x; t) = - \sum_{\ell=1}^k \sum_{j=1}^k c_j (J(t)^j)_{n+1, n-\ell+1} \left(\prod_{j=n-\ell+1}^n a_j \right) P_{n-\ell}(x; t), \tag{9.3}$$

where $J(t)$ is the Jacobi matrix of $d\rho_t$.

Remark. The vector indices associated to J start at 1, so (with $p_j = P_j/\|P_j\|$)

$$J_{jk} = \langle p_{j-1}, xp_{k-1} \rangle, \tag{9.4}$$

$$J_{jj} = b_j = \langle p_{j-1}, xp_{j-1} \rangle, \tag{9.5}$$

$$J_{jj+1} = J_{j+1j} = a_j = \langle p_j, xp_{j-1} \rangle. \tag{9.6}$$

Example 9.2. Take $\frac{1}{2}f'(x) = x$ (the Toda case). Then $J_{n+1, n}$ is all that is relevant, and it is a_n , so (9.3) becomes

$$\dot{P}_n = -a_n^2 P_{n-1}. \tag{9.7}$$

This Toda case is known; see, for example, Peherstorfer [43] and Barrios–Hernández [3].

Example 9.3. Take $\frac{1}{2}f'(x) = x^2$. We have

$$(J^2)_{n+1, n-1} = a_n a_{n-1}, \quad (J^2)_{n+1, n} = a_n (b_{n+1} + b_n)$$

so

$$\dot{P}_n = -a_n^2 (b_{n+1} + b_n) P_{n-1} - a_n^2 a_{n-1}^2 P_{n-2}. \tag{9.8}$$

Remark. Using $\langle P_n, P_n \rangle = a_n a_{n-1} \dots a_1$ and (9.5), one can easily go from differential equations for P_n back to those for a_n and b_n .

Proof of Theorem 9.1. Since P_n is monic, \dot{P}_n is a polynomial of degree $n - 1$. Thus

$$\begin{aligned} \dot{P}_n &= \sum_{\ell=1}^n \langle \dot{P}_n, p_{n-\ell} \rangle p_{n-\ell} \\ &= - \sum_{\ell=1}^n \langle P_n, \frac{1}{2} f' p_{n-\ell} \rangle p_{n-\ell} \\ &= - \sum_{\ell=1}^n \langle p_n, \frac{1}{2} f' p_{n-\ell} \rangle \frac{\|P_n\|}{\|P_{n-\ell}\|} P_{n-\ell} \\ &= \text{RHS of (9.3)} \end{aligned} \tag{9.9}$$

where (9.9) comes from

$$\begin{aligned} O &= \frac{d}{dt} \int P_n p_{n-\ell} d\mu_t \\ &= \langle \dot{P}_n, p_{n-\ell} \rangle + \langle P_n, \dot{p}_{n-\ell} \rangle + \langle P_n, \frac{1}{2} f' p_{n-\ell} \rangle \\ &= \langle \dot{P}_n, p_{n-\ell} \rangle + \langle P_n, \frac{1}{2} f' p_{n-\ell} \rangle \end{aligned}$$

since $\dot{p}_{n-\ell}$ is a polynomial of degree $n - \ell$, and so orthogonal to P_n . □

One can also ask about derivatives of $p_n = P_n / \|P_n\|$. Since

$$\dot{p}_n = \frac{\dot{P}_n}{\|P_n\|} - \left[\frac{P_n}{\|P_n\|} \right] \frac{\|P_n\| \cdot}{\|P_n\|} \tag{9.10}$$

we clearly need only find $\frac{d}{dt} \log \|P_n\|$. Following [47, Section 9.10],

Proposition 9.4. *Under the map,*

$$d\rho_t(x) = \frac{e^{1/2 f'(x)t} d\rho_0}{\int e^{1/2 f'(x)t} d\rho_0}$$

with f' given by (9.1), the monic orthogonal polynomials $P_n(x; d\rho_t) = P_n(x; t)$ obey

$$\frac{d}{dt} \log \|P_n\| = \frac{1}{2} \sum_{j=1}^k c_j [(J^j)_{n+1 \ n+1} - (J^j)_{1 \ 1}]. \tag{9.11}$$

Proof. Let

$$N_t = \int e^{1/2 f'(x)t} d\rho_t. \tag{9.12}$$

Since \dot{P}_n is orthogonal to P_n ,

$$\frac{d}{dt} [\|P_n\|^2 N_t] = \left(\int P_n^2 (\frac{1}{2} f') d\rho_t \right) N_t \tag{9.13}$$

$$= \left[\sum_{j=1}^k c_j (J^j)_{n+1 \ n+1} \right] \|P_n\|^2 N_t \tag{9.14}$$

since

$$\int P_n^2 (\frac{1}{2} f') d\rho_t = \|P_n\|^2 \langle p_n, (\frac{1}{2} f') p_n \rangle \tag{9.15}$$

and we have (9.4). Similarly,

$$\begin{aligned} \frac{d}{dt} N_t &= N_t \int (\frac{1}{2} f) d\rho_t \\ &= (9.14) \text{ for } n = 0. \end{aligned} \tag{9.16}$$

Since

$$\frac{d}{dt} \log \|P_n\| = \frac{1}{2} \left[\frac{d}{dt} \log[\|P_n\|^2 N_t] - \frac{d}{dt} \log N_t \right]$$

(9.14) and (9.16) imply (9.11). \square

By (9.3), (9.10), and (9.11), we immediately have:

Theorem 9.5. *Under the map,*

$$d\rho_t(x) = \frac{e^{1/2f'(x)t} d\rho_0}{\int e^{1/2f'(x)t} d\rho_0}$$

with f' given by (9.1), the orthonormal polynomials $p_n = P_n/\|P_n\|$ obey

$$\dot{p}_n = -p_n \left(\frac{1}{2} \sum_{j=1}^k c_j [(J^j)_{n+1, n+1} - (J^j)_{11}] \right) - \sum_{\ell=1}^k \left[\sum_{j=1}^k c_j (J^j)_{n+1, n-\ell+1} \right] p_{n-\ell}. \tag{9.17}$$

10. Differential equations for OPUC

For OPUC, one looks at flows generated by Laurent polynomials, $f(z)$, real on $\partial\mathbb{D}$. If

$$g(e^{i\theta}) = \frac{\partial}{\partial\theta} f(e^{i\theta}) = \sum_{j=-\ell}^{\ell} b_j e^{ij\theta}, \tag{10.1}$$

where $b_{-j} = \bar{b}_j$, then

$$d\mu_t(\theta) = \frac{e^{tg(e^{i\theta})} d\mu_0(\theta)}{\int e^{tg(e^{i\theta})} d\mu_0(\theta)}. \tag{10.2}$$

The analog of the P_n 's are the unnormalized CMV and alternate CMV bases

$$Y_n = \rho_0 \dots \rho_{n-1} \chi_n, \tag{10.3}$$

$$X_n = \rho_0 \dots \rho_{n-1} x_n, \tag{10.4}$$

where $\{\chi_n\}_{n=0}^\infty$ and $\{x_n\}_{n=0}^\infty$ are the CMV and alternate CMV bases (see [46, Section 4.2]). Then the exact same argument as led to Theorem 9.1 leads to:

Theorem 10.1. *Let $d\mu_t$ be given by (10.2) with g given by (10.1). Then if $Y_n(t) = Y_n(d\mu_t)$,*

$$\dot{Y}_n = - \sum_{m=0}^{n-1} \sum_{k=-\ell}^{\ell} b_k (C^k)_{mn} \left(\prod_{j=m}^{n-1} \rho_j \right) Y_m \tag{10.5}$$

with the same equation for \dot{X}_n except the $(C^k)_{mn}$ is replaced by $(C^k)_{nm}$.

Remark. These are analogs of Proposition 9.4 and Theorem 9.5.

Example 10.2 (Schur Flow). We have

$$g(z) = z + z^{-1}. \tag{10.6}$$

Ismail [24] (see also [21]) obtained the following differential equation for the monic OPUC:

$$\dot{\Phi}_n = \Phi_{n+1} - (z + \bar{\alpha}_n \alpha_{n-1})\Phi_n - \rho_{n-1}^2 \Phi_{n-1}. \tag{10.7}$$

We want to show this is equivalent to (10.5). From our point of view, (10.7) is unusual since our arguments show that $\dot{\Phi}_n$ is a linear continuation of $\{\Phi_j\}_{j=0}^{n-1}$ while (10.7) has Φ_{n+1} and Φ_n . (Since there is a $z\Phi_n$, in fact, one can hope—and it happens—that RHS of (10.7) is $\sum_{j=0}^{n-1} c_j \Phi_j$.)

Since $\Phi_{n+1} = z\Phi_n - \bar{\alpha}_n \Phi_n^*$,

$$\text{RHS of (10.7)} = -\bar{\alpha}_n(\Phi_n^* + \alpha_{n-1}\Phi_n) - \rho_{n-1}^2 \Phi_{n-1}. \tag{10.8}$$

Since (see (1.5.41) of [46]) $\Phi_n^* + \alpha_{n-1}\Phi_n = \rho_{n-1}^2 \Phi_{n-1}^*$,

$$\text{RHS of (10.7)} = -\bar{\alpha}_n \rho_{n-1}^2 \Phi_{n-1}^* - \rho_{n-1}^2 \Phi_{n-1} \tag{10.9}$$

verifying that RHS of (10.7) is indeed a polynomial of degree at most $n - 1$.

(10.9) mixes Y 's and X 's, so we use (see (1.5.40) of [46]) $\Phi_{n-1} = \rho_{n-2}^2 z \Phi_{n-2} - \bar{\alpha}_{n-2} \Phi_{n-1}^*$ to find

$$\text{RHS of (10.7)} = -\rho_{n-1}^2 (\bar{\alpha}_n - \bar{\alpha}_{n-2}) \Phi_{n-1}^* - \rho_{n-1}^2 \rho_{n-2}^2 (z \Phi_{n-2})$$

which one can see is exactly (10.7) or its X_n analog (depending on whether n in Φ_n is even or odd). \square

11. Application 3: Poisson commutators for periodic Jacobi and CMV matrices

In this section, we want to show how the fundamental relations (1.11)/(1.23) on finite Jacobi and CMV matrices yield a proof of the basic Poisson bracket relations for the periodic case. For OPRL, the original proofs of this relation by Flaschka [15] via direct calculation of Poisson brackets of eigenvalues are simple, but for OPUC, the two existing proofs by Nenciu–Simon [42] and Nenciu [39,40] are computationally involved.

Given $\{a_j, b_j\}_{j=1}^p \in \mathbb{R}^p$ or $\{\alpha_j\}_{j=0}^{p-1} \in \mathbb{C}^p$ (in the OPUC case, we assume p is even for reasons discussed in Chapter 11 of [47]), we define periodic Jacobi and CMV matrices by first extending the parameters periodically to \mathbb{Z} , that is,

$$a_{j+p} = a_j, \quad b_{j+p} = b_j, \quad \alpha_{j+p} = \alpha_j \tag{11.1}$$

and then letting J, \mathcal{C} act on $\ell^2(\mathbb{Z})$ as two-sided matrices with these periodic parameters (two-sided CMV matrices are discussed on pp. 589–590 of [47]).

In each case, J, \mathcal{C} act on $\ell^\infty(\mathbb{Z})$ and commute with $\mathcal{S}^p: \{u_n\} \rightarrow \{u_{n+p}\}$. For each $e^{i\theta} \in \partial\mathbb{D}$, J, \mathcal{C} leave invariant the p -dimensional space

$$\ell_\theta^\infty = \{u \mid \mathcal{S}^p u = e^{i\theta} u\}$$

and so define $J(\theta), \mathcal{C}(\theta)$ as operators on ℓ_θ^∞ . In terms of the natural basis $\{\delta_j^\theta\}_{j=1}^N$,

$$(\delta_j^\theta)_k = \begin{cases} e^{i\theta \ell} & k = p\ell + j, \\ 0 & k \not\equiv j \pmod p, \end{cases} \tag{11.2}$$

$J(\theta)$ has a matrix of the form

$$\begin{pmatrix} b_1 & a_1 & 0 & \dots & e^{i\theta} a_p \\ a_1 & b_2 & a_2 & \dots & 0 \\ \vdots & \vdots & \vdots & & \vdots \\ e^{-i\theta} a_p & 0 & 0 & \dots & b_p \end{pmatrix} \tag{11.3}$$

of a Jacobi matrix with an extra term in the corners. The $\mathcal{C}(\theta)$ are described on pp. 719–720 of [47].

In each case, there is a natural transfer or monodromy matrix, $T_p(z)$ on \mathbb{C}^2 , so if $u \in \mathbb{C}^\infty$ solves $(J - z)u = 0$ (resp. $(\mathcal{C} - z)v = 0$), then

$$T_p(z) \begin{pmatrix} u_1 \\ u_0 \end{pmatrix} = \begin{pmatrix} u_{p+1} \\ u_p \end{pmatrix}. \tag{11.4}$$

Constancy of Wronskians implies

$$\det(T_p(z)) = \begin{cases} 1 & \text{(OPRL),} \\ z^p & \text{(OPUC).} \end{cases} \tag{11.5}$$

One defines the discriminant, $\Delta(z)$, by

$$\Delta(z) = \begin{cases} \text{Tr}(T_p(z)) & \text{(OPRL),} \\ z^{-p/2} \text{Tr}(T_p(z)) & \text{(OPUC)} \end{cases} \tag{11.6}$$

(recall p is even in the OPUC case). z is clearly an eigenvalue of $J(\theta)$ (resp. $\mathcal{C}(\theta)$) if and only if $e^{i\theta}$ is an eigenvalue of $T_p(z)$ (resp. $z^{-p/2}T_p(z)$; see [47, p. 722]). So, by (11.5),

$$z \text{ is an eigenvalue of } J(\theta) \text{ or } \mathcal{C}(\theta) \Leftrightarrow \Delta(z) = 2 \cos(\theta). \tag{11.7}$$

For later purposes, we want to note that in neither case is $\Delta(z)$ automatically monic. Rather,

$$\Delta(z) = \begin{cases} \frac{1}{a_1 \dots a_p} z^p + \text{lower order} & \text{(OPRL),} \\ \frac{1}{\rho_1 \dots \rho_p} (z^{p/2} + z^{-p/2} + \text{orders in between}) & \text{(OPUC)} \end{cases} \tag{11.8}$$

(see [51] and Chapter 11 of [47]).

$\mathbb{R}^p \times (0, \infty)^{p-1}$ (resp. \mathbb{D}^p) are given Poisson brackets by

$$\{b_k, a_k\} = -\frac{1}{4} a_k, \quad k = 1, 2, \dots, p, \tag{11.9}$$

$$\{b_k, a_{k-1}\} = \frac{1}{4} a_{k-1}, \quad k = 2, \dots, p, \tag{11.10}$$

$$\{b_1, a_p\} = \frac{1}{4} a_p \tag{11.11}$$

and

$$\{\alpha_j, \alpha_k\} = \{\bar{\alpha}_j, \bar{\alpha}_k\} = 0, \quad \{\alpha_j, \bar{\alpha}_k\} = -i \rho_j^2 \delta_{jk}, \quad j, k = 0, \dots, p - 1.$$

One big difference between OPRL and OPUC is that in the OPUC case, the center of the Poisson bracket (i.e., the functions that Poisson commute with all functions) is trivial, that is, $\{\cdot, \cdot\}$

defines a symplectic form. But in the OPRL case,

$$\sum_{j=1}^p b_j \quad \text{and} \quad \prod_{j=1}^p a_j \tag{11.12}$$

are easily seen to have zero Poisson brackets with all a 's and b 's, so the two-form defined by $\{\cdot, \cdot\}$ is degenerate, but on the subspace where

$$\sum_{j=1}^p b_j = \beta \quad \text{and} \quad \prod_{j=1}^p a_j = \alpha \tag{11.13}$$

the form is nondegenerate and the Poisson bracket defines a symplectic form.

Our main results in this section are:

Theorem 11.1 (Flaschka [15]). *For OPRL:*

$$\{\Delta(x), \Delta(y)\} = 0. \tag{11.14}$$

If $\lambda_j(\theta)$ are the simple eigenvalues of $J(\theta)$, then

$$\{\lambda_j(\theta), \lambda_k(\theta')\} = 0. \tag{11.15}$$

Remarks. (1) (11.15) is normally only stated for $\theta = \theta'$, but we will see it is immediate from (11.14) and (11.7).

(2) For $\theta \neq 0, \pi$, all eigenvalues are simple and $\lambda_j(\theta)$ are global nonsingular functions. For $\theta = 0, \pi$, there are subvarieties of codimension 3 where some eigenvalues are degenerate. $\lambda_j(\theta)$ are smooth away from this subvariety and (11.15) only holds there.

Theorem 11.2 (Nenciu–Simon [42]). *For OPUC:*

$$\{\Delta(z), \Delta(w)\} = 0. \tag{11.16}$$

If $\lambda_j(\theta)$ are the simple eigenvalues of $\mathcal{C}(\theta)$, then

$$\{\lambda_j(\theta), \lambda_k(\theta')\} = 0 \tag{11.17}$$

and

$$\left\{ \prod_{k=0}^{p-1} \rho_k, \lambda_j(\theta) \right\} = 0. \tag{11.18}$$

Remarks. (1) As in the OPRL case, $\lambda_j(\theta)$ are simple if $\theta \neq 0, \pi$ and off a closed subvariety of codimension 3 if $\theta = 0$ or $\theta = \pi$.

(2) $\prod_{k=0}^{p-1} \rho_k$ is the inverse leading term in $\Delta(z)$ for OPUC. The analog for OPRL is $\prod_{k=1}^p a_k$. Of course, $\{\prod_{k=1}^p a_k, \lambda_j(\theta)\} = 0$, but we do not say it explicitly since for OPRL, $\prod_{k=1}^p a_k$ Poisson commutes with any function!

We note two general related aspects of these theorems. First, there are no variables analogous to the ρ 's and μ 's allowing a simple exact solution. The natural complementary variables are the

Dirichlet data which are defined specially on each isospectral set rather than as a local function. The proper angle variables for this situation are discussed in terms of the theory of Abelian integrals on a suitable hyperelliptic surface [7,11,19,26–28,34,38,54,55].

Second, there is a sense in which the periodic situation is more subtle and interesting than the finite N structure. For finite N , the isospectral manifolds are open simplexes parametrized by $\{\rho_j\}_{j=1}^N$ (or μ_j) with $\sum_{j=1}^N \rho_j = 1$. Topologically this is \mathbb{R}^{N-1} . For the periodic case, the isospectral manifolds are compact because $\prod_{j=1}^p a_j$ or $\prod_{j=1}^p \rho_j$ const. prevents a_j or ρ_j from going to zero (for OPRL one also needs boundedness of $\sum a_j^2$). By the general theory of completely integrable systems, these isospectral sets are tori generically of dimension one-half the dimension of the symplectic manifold ($p - 1$ for OPRL, p for OPUC).

The idea of the proofs of these theorems is to cut off the periodic matrix to a finite matrix and take limits of Poisson brackets. The limits of finite traces (normalized) will be the moments of the density of states, not the periodic eigenvalues—but we will see they are related. It will be useful to go back and forth between the two sets of fundamental symmetric functions:

$$t_k^{(N)}(\lambda_1, \dots, \lambda_N) = \sum_{j=1}^N \lambda_j^k, \tag{11.19}$$

$$s_k^{(N)}(\lambda_1, \dots, \lambda_N) = \sum_{i_1 < \dots < i_k} \lambda_{i_1} \dots \lambda_{i_k} \quad 1 \leq k \leq N, \tag{11.20}$$

where we will need the following well-known combinatorial result:

Proposition 11.3. *For $k \leq N$, we have*

$$s_k^{(N)}(\lambda_1, \dots, \lambda_N) = \frac{1}{k} t_k^{(N)}(\lambda_1, \dots, \lambda_N) + r, \tag{11.21}$$

where r is a polynomial in $\{t_j^{(N)}\}_{j=1}^{k-1}$ and also a polynomial in $\{s_j^{(N)}\}_{j=1}^{k-1}$.

Proof. Here is a simple proof. Let A be the diagonal matrix with eigenvalues $\lambda_1, \dots, \lambda_N$. Let $\wedge^k(\mathbb{C}^N)$ be the k -fold antisymmetric product and $\otimes^k(\mathbb{C}^N)$ the tensor product (see, e.g., Appendix A of [44]). Let $\pi \in \Sigma_k$, the symmetric group in k -objects, act as \vee_π on $\otimes^k(\mathbb{C}^N)$ via $\vee_\pi(x_1 \otimes \dots \otimes x_k) = x_{\pi(1)} \otimes \dots \otimes x_{\pi(k)}$. Then

$$P_k = \frac{1}{k!} \sum_{\pi \in \Sigma_k} (-1)^\pi \vee_\pi$$

is the projection of $\otimes^k \mathbb{C}^N$ to $\wedge^k(\mathbb{C}^N)$. Thus,

$$\begin{aligned} s_k^{(N)}(\lambda_1, \dots, \lambda_N) &= \text{Tr}_{\wedge^k(\mathbb{C}^N)}(\wedge^k(A)) \\ &= \text{Tr}_{\otimes^k(\mathbb{C}^N)}(P_k \otimes^k A) \\ &= \frac{1}{k!} \sum_{\pi \in \Sigma_k} (-1)^\pi \text{Tr}(\vee_\pi \otimes^k A). \end{aligned} \tag{11.22}$$

It is easy to see [44, Appendix A] that if π 's disjoint cycle decomposition has ℓ_1 one-cycles, ℓ_2 two-cycles, etc., then

$$\begin{aligned} \text{Tr}(\vee_{\pi} \otimes^k A) &= \text{Tr}(A)^{\ell_1} \text{Tr}(A^2)^{\ell_2} \dots \text{Tr}(A^k)^{\ell_k} \\ &= \prod_{j=1}^k t_j^{(N)}(\lambda_1, \dots, \lambda_N)^{\ell_j}. \end{aligned} \tag{11.23}$$

Since there are $(k - 1)!$ permutations with one k -cycle, (11.22) and (11.23) imply (11.21) with r a polynomial in t 's. An easy induction then shows any t_k is a polynomial in $\{s_j\}_{j=1}^k$ and so r is also a polynomial in the s 's. \square

The density of states of a periodic Jacobi or CMV matrix can be defined as the unique measure, $d\gamma$, (on \mathbb{R} or $\partial\mathbb{D}$) so that for all $k \in \{0, 1, 2, \dots\}$,

$$\int \lambda^k d\gamma(\lambda) = \int_0^{2\pi} \left[\frac{1}{p} \sum_{j=1}^p \lambda_j(\theta)^k \right] \frac{d\theta}{2\pi}. \tag{11.24}$$

Moreover, it is easy to see [51] that if $J_m(\{a_k\}_{k=1}^p, \{b_k\}_{k=1}^p)$ is the cutoff Jacobi matrix obtained by taking the two-sided infinite Jacobi matrix and projecting onto the span of $\{\delta_j\}_{j=-m}^m$, then for any $k \in \{0, 1, 2, \dots\}$,

$$\frac{1}{2m + 1} \text{Tr}(J_m^k) \rightarrow \int \lambda^k d\gamma(\lambda). \tag{11.25}$$

Similarly in the CMV case [47], if C_m is defined by taking $\alpha_{-m} \rightarrow -1$ and $\alpha_m \rightarrow \beta$ (for any β), then

$$\frac{1}{2m + 1} \text{Tr}(C_m^k) \rightarrow \int \lambda^k d\gamma(\lambda). \tag{11.26}$$

The key to the proof of Theorem 11.1 is the following. Define for $\{a_j, b_j\}_{j=1}^p$ fixed,

$$t_k(\theta) = \text{Tr}(J(\theta)^k), \tag{11.27}$$

$$s_k(\theta) = \sum_{j_1 < \dots < j_k} \lambda_{j_1}(\theta) \dots \lambda_{j_k}(\theta), \quad k = 0, 1, \dots, p. \tag{11.28}$$

Theorem 11.4. Consider OPRL of period p .

- (i) For $k \leq p - 1$, $s_k(\theta)$ is independent of θ .
- (ii) For $k \leq p - 1$, $t_k(\theta)$ is independent of θ .
- (iii)

$$s_p(\theta) = s_p(0) + (-1)^p \left(\prod_{j=1}^p a_j \right) (2 - 2 \cos \theta) \tag{11.29}$$

- (iv)

$$t_p(\theta) = t_p(0) + (-1)^p p \left(\prod_{j=1}^p a_j \right) (2 - 2 \cos \theta). \tag{11.30}$$

In particular, for $k \leq p$,

$$t_k(0) = \left[(-1)^{p+1} (2p) \prod_{j=1}^p a_j \right] \delta_{kp} + \int \lambda^k d\gamma(\lambda). \tag{11.31}$$

Remark. Directly from the form of the matrix (11.3), one can see (ii) easily, and (iv) with a little more work.

Proof. By (11.7) and (11.8), we see that

$$\prod_{j=1}^p (x - \lambda_j(\theta)) = \left(\prod_{j=1}^p a_j \right) (\Delta(x) - 2 \cos \theta) \tag{11.32}$$

which, expanding the products as

$$\prod_{j=1}^p (x - \lambda_j(\theta)) = x^p + \sum_{j=1}^p (-1)^j s_j(\theta) x^{p-j} \tag{11.33}$$

immediately implies (i) and (iii). (ii) and (iv) then follow from (11.21), and (11.31) then follows from (11.24). \square

For OPUC we have:

Theorem 11.5. Consider OPUC of even period p .

- (i) For $k = 1, 2, \dots, \frac{p}{2} - 1, \frac{p}{2} + 1, \dots, p - 1$, $s_k(\theta)$ is independent of θ .
- (ii) For $k = 1, 2, \dots, \frac{p}{2} - 1$, $t_k(\theta)$ is independent of θ .
- (iii)

$$s_{p/2}(\theta) = s_{p/2}(0) + (-1)^{p/2} \left(\prod_{j=0}^{p-1} \rho_j \right) (2 - 2 \cos \theta) \tag{11.34}$$

(iv)

$$t_{p/2}(\theta) = t_{p/2}(0) + (-1)^{p/2} \frac{p}{2} \left(\prod_{j=0}^{p-1} \rho_j \right) (2 - 2 \cos \theta). \tag{11.35}$$

In particular, for $k \leq p/2$,

$$t_k(0) = (-1)^{p/2+1} p \left(\prod_{j=0}^p \rho_j \right) \delta_{kp/2} + \int \lambda^k d\gamma(\lambda). \tag{11.36}$$

Remark. It will suffice to have control of t_j and \bar{t}_j for $0 \leq |j| \leq p/2$ since $s_{p-j} = \bar{s}_j$, as we will see.

Proof. By (11.8) and (11.7) and the fact that $z^{p/2} \Delta(z)$ is a polynomial nonvanishing at $z = 0$, we have that

$$\prod_{j=1}^p (z - \lambda_j(\theta)) = \left(\prod_{j=0}^{p-1} \rho_j \right) (z^{p/2} \Delta(z) - 2 \cos(\theta) z^{p/2}) \tag{11.37}$$

which, via (11.33), implies (i) and (iii). (ii) and (iv) then follow from (11.21), and (11.36) from (11.24). \square

Proof of Theorem 11.1. Let

$$T_k^{(m)}(\{a_j, b_j\}_{j=0}^p) = \text{Tr} \left([J_m(\{a_k, b_k\}_{k=0}^p)]^k \right) \tag{11.38}$$

and

$$\tilde{T}_k^{(m)}(\{a_j, b_j\}_{j=-m}^m) = \text{Tr} \left([J(\{a_k, b_k\}_{j=-m}^m)]^k \right). \tag{11.39}$$

So $T_k^{(m)}$ is just $\tilde{T}_k^{(m)}$ restricted to periodic sequences.

Let

$$t_k(\{a_j, b_j\}_{j=0}^p) = \int \lambda^k d\gamma(\lambda). \tag{11.40}$$

As noted in (11.25), $\frac{1}{2m+1} T_k^{(m)} \rightarrow t_k$. This can be seen by noting diagonal matrix elements of $(J_m)^k$ are uniformly bounded in m for k fixed and equal to those of the infinite matrix so long as their index j_0 obeys $|j_0| \leq m - 2k - 1$.

This same equality and polynomial nature show for $1 \leq j \leq p$,

$$\frac{\partial}{\partial a_{j+\ell p}} \tilde{T}_k^{(m)} \Big|_{\substack{a_j=a_j^{(0)} \\ b_j=b_j^{(0)}}} = \frac{\partial}{\partial a_j} t_k \left(\{a_j^{(0)}, b_j^{(0)}\}_{j=1}^p \right) \tag{11.41}$$

where $\{a_j^{(0)}, b_j^{(0)}\}$ is periodic, and similarly for b derivatives. (11.41) holds so long as $|j + \ell p| \leq m - 2k - 1$. It follows that

$$\frac{p}{m} \{ \tilde{T}_k^{(m)}, \tilde{T}_j^{(m)} \} \Big|_{\substack{a=a^{(0)} \\ b=b^{(0)}}} \rightarrow \{t_k, t_\ell\}. \tag{11.42}$$

Since $\{ \tilde{T}_l^{(m)}, \tilde{T}_j^{(m)} \} = 0$, we conclude

$$\{t_k, t_\ell\} = 0 \tag{11.43}$$

for all k, ℓ . By (11.31)

$$\{t_k(0), t_\ell(0)\} = 0 \tag{11.44}$$

for all k, ℓ . Thus by (11.21), we see for $1 \leq k, \ell \leq p$,

$$\{s_k, s_\ell\} = 0. \tag{11.45}$$

Since $\Delta(x) - 2 = (a_1 \dots a_p)^{-1} \prod_{j=1}^p (x - \lambda_j(0))$ and (11.33) holds and $(a_1 \dots a_p)^{-1}$ is in the Poisson center, we see that

$$\{\Delta(x), \Delta(y)\} = 0. \tag{11.46}$$

Thus

$$\{\Delta(x) - 2 \cos(\theta), \Delta(y) - 2 \cos(\theta')\} = 0. \tag{11.47}$$

Evaluating this at $x = \lambda_j(\theta), y = \lambda_k(\theta')$, we get (11.15) as usual. \square

Proof of Theorem 11.2. By repeating the arguments from the last proof, we see that (11.44) holds and that

$$\{t_k(0), t_\ell(0)\} = \{\overline{t_k(0)}, t_\ell(0)\} = 0. \tag{11.48}$$

From this and Theorem 11.5, we see for $1 \leq k, \ell \leq p/2$,

$$\{s_k(0), s_\ell(0)\} = \{s_k(0), \overline{s_\ell(0)}\} = 0. \tag{11.49}$$

By (11.8), the product of the roots is 1, that is,

$$\prod_{j=1}^p \lambda_j(\theta) = 1.$$

Since $\lambda_j \bar{\lambda}_j = 1$ also, we see

$$\bar{s}_{p-j}(\theta) = \prod_{j=1}^p \lambda_j(\theta) \overline{s_{p-j}(\theta)} = s_j(\theta) \tag{11.50}$$

so for $1 \leq k, \ell \leq p - 1$,

$$\{s_k(0), s_\ell(0)\} = 0. \tag{11.51}$$

This, (11.37), and (11.8) imply that

$$\left\{ \left(\prod_{j=0}^{p-1} \rho_j \right) \Delta(z), \left(\prod_{j=0}^{p-1} \rho_j \right) \Delta(w) \right\} = 0. \tag{11.52}$$

By a simple calculation if $\alpha_j = |\alpha_j| e^{i\theta_j}$, then

$$\left\{ \prod_{j=0}^{p-1} \rho_j^2, g \right\} = - \prod_{j=0}^{p-1} \rho_j^2 \sum_{j=0}^{p-1} \frac{\partial g}{\partial \theta_j} \tag{11.53}$$

so $-\log(\prod_{j=0}^{p-1} \rho_j^2)$ generates the Hamiltonian flow

$$\alpha_j \rightarrow \alpha_j e^{it}. \tag{11.54}$$

This transformation is implementable by a unitary (see, e.g., [49]), and so it leaves the $\lambda_j(\theta)$ fixed. Thus, (11.18) holds, which implies

$$\left\{ \prod_{j=0}^{p-1} \rho_j, \left(\prod_{j=0}^{p-1} \rho_j \right) \Delta(z) \right\} = 0 \tag{11.55}$$

so (11.52) implies (11.16), which in turn implies (11.17). \square

Remark. Unlike OPRL where $\prod_{j=1}^p a_j$ is in the Poisson center, $\prod_{j=0}^{p-1} \rho_j$ generates a nontrivial flow, but one that leaves $\lambda_j(\theta)$ invariant.

12. More Poisson brackets for OPRL

In this section, we will discuss some additional Poisson brackets for OPRL and related functions as a way of illuminating and extending the major results that we proved earlier. We want to begin by showing that one can go backwards from (1.11) to (1.15)/(1.16) (or, more precisely, to the Poisson brackets (1.16) and $\{P_n(x), P_n(y)\} = 0$; that $\{Q_n(x), Q_n(y)\} = 0$ is then a consequence of (2.10)).

Clearly,

$$\{P_n(x), P_n(y)\} = 0 \Leftrightarrow \{x_j, x_k\} = 0. \tag{12.1}$$

To see that

$$\{x_j, \rho_k\} = \frac{1}{2} [\delta_{jk} \rho_j - \rho_j \rho_k] \Leftrightarrow (1.16) \tag{12.2}$$

we compute using (3.5) and $\{x_j, x_k\} = 0$,

$$\begin{aligned} 2\{P_n(x), Q_n(y)\} &= 2 \sum_{j,k} -\{x_j, \rho_k\} \frac{P_n(x)}{x - x_j} \frac{P_n(y)}{y - y_k} \\ &= \sum_{j,k} \rho_j \rho_k \frac{P_n(x)}{x - x_j} \frac{P_n(y)}{y - x_k} - \sum_j \rho_j \frac{P_n(x) P_n(y)}{(x - x_j)(y - x_j)}. \end{aligned} \tag{12.3}$$

The first sum gives $Q_n(x)Q_n(y)$ by (2.10) and the second sum is $(x - y)^{-1}[P_n(x)Q_n(y) - P_n(y)Q_n(x)]$ if we note that

$$\frac{1}{(x - x_j)} \frac{1}{(y - x_j)} = \frac{1}{x - y} \left[\frac{1}{y - x_j} - \frac{1}{x - x_j} \right]. \tag{12.4}$$

This establishes (12.2).

By (2.11) (with n replaced by $n - 1$) used in the first factor $\{P_n(x), P_n(y)\}$, we get a relation among $\{P_{n-1}(x), P_n(y)\}$, $\{b_1, P_n(y)\}$, $\{a_1^2, P_n(y)\}$, and $\{Q_{n-1}(x), P_n(y)\}$. Since the first three are computed earlier, we get a formula for $\{Q_{n-1}(x), P_n(y)\}$, namely,

$$\begin{aligned} \{Q_{n-1}(x), P_n(y)\} &= 2P_{n-1}(x) - 2Q_{n-1}(x) \\ &\quad - b_1 a_1^{-2} \left(\frac{P_n(x) P_{n-1}(y) - P_{n-1}(x) P_n(y)}{x - y} - P_{n-1}(x) P_{n-1}(y) \right). \end{aligned} \tag{12.5}$$

Next, we want to discuss Poisson brackets with

$$m_n(x) = -\frac{Q_n(x)}{P_n(x)} \tag{12.6}$$

both to link to earlier work of Faybusovich–Gekhtman [14] and because one can then take $n \rightarrow \infty$.

Theorem 12.1. *We have that*

$$\{P_n(x), m_n(y)\} = \frac{1}{2} \left[Q_n(x)m_n(y) - \frac{P_n(x)m_n(y) + Q_n(x)}{x - y} \right], \tag{12.7}$$

$$\{Q_n(x), m_n(y)\} = \frac{1}{2} \left[-Q_n(x)m_n(y)^2 + m_n(y) \left[\frac{P_n(x)m_n(y) + Q_n(x)}{x - y} \right] \right]. \tag{12.8}$$

Proof. We have, since $\{P_n(x), P_n(y)\} = 0$, that

$$\{P_n(x), m_n(y)\} = -P_n(y)^{-1} \{P_n(x), Q_n(y)\}. \tag{12.9}$$

The second term on the right of (1.16) leads to

$$-\frac{1}{2} P_n(y)^{-1} Q_n(x)Q_n(y) = \frac{1}{2} Q_n(x)m_n(y).$$

The first term on the right leads to

$$\begin{aligned} &\frac{1}{2} P_n(y)^{-1} [P_n(x)Q_n(y) - Q_n(x)P_n(y)](x - y)^{-1} \\ &= (-\frac{1}{2} P_n(x)m_n(y) - \frac{1}{2} Q_n(x))(x - y)^{-1}, \end{aligned}$$

proving (12.7).

On the other hand, since $\{Q_n(x), Q_n(y)\} = 0$,

$$\begin{aligned} \{Q_n(x), m_n(y)\} &= Q_n(y) \{Q_n(x), -P_n(y)^{-1}\} \\ &= m_n(y) P_n(y)^{-1} \{P_n(y), Q_n(x)\} \\ &= m_n(y) P_n(y)^{-1} \{P_n(x), Q_n(y)\} \end{aligned} \tag{12.10}$$

$$= m_n(y) \{P_n(x), m_n(y)\} \tag{12.11}$$

so (12.7) implies (12.8). (12.10) follows from the symmetry of $\{P_n(x), Q_n(y)\}$ under $x \leftrightarrow y$, and (12.11) from (12.9). \square

Theorem 12.2 (Faybusovich and Gekhtman [14]). *We have that*

$$\{m_n(z), m_n(w)\} = \frac{1}{2} (m_n(z) - m_n(w)) \left[-\frac{m_n(z) - m_n(w)}{z - w} + m_n(z)m_n(w) \right]. \tag{12.12}$$

Proof. Since $\{P_n(z), P_n(w)\} = 0 = \{Q_n(z), Q_n(w)\}$ and $\{P_n(z), Q_n(w)\}$ is symmetric under $z \leftrightarrow w$ and $\{f, g\}$ is antisymmetric under $f \leftrightarrow g$,

$$\begin{aligned} \{m_n(z), m_n(w)\} &= -\frac{Q_n(z)}{P_n(w)Q_n(z)^2} \{P_n(z), Q_n(w)\} - (z \leftrightarrow w) \\ &= \frac{m_n(z)}{2} \left[m_n(z)m_n(w) - \frac{m_n(z) - m_n(w)}{z - w} \right] - (z \leftrightarrow w). \end{aligned} \tag{12.13}$$

which is (12.12).

As $n \rightarrow \infty$, m_n has a limit so long as $\{a_j, b_j\}_{j=0}^\infty$ are bounded (actually so long as the moment problem is determinate), so (12.12) holds if $m_n(z)$ is replaced by $m(z)$ for semi-infinite Jacobi matrices.

The unnormalized transfer matrix has the form

$$\begin{pmatrix} P_n & Q_n \\ P_{n-1} & Q_{n-1} \end{pmatrix}.$$

Of the 16 Poisson brackets for these four functions, we have computed 12. It would be interesting to know $\{Q_n(z), P_{n-1}(w)\}$. Similarly, it would be interesting to know the Poisson brackets for the periodic transfer matrix and use them to prove (11.14). During the refereeing of this paper, Tsiganov [53] posted a preprint that computes Poisson brackets of transfer matrices associated to Toda lattices.

13. More Poisson brackets for OPUC

We begin by showing that one can go backwards from (1.23) to (1.18)/(1.19) (or, more precisely, to (1.19) and $\{P_n(z), P_n(w)\} = 0$; $\{Q_n(z), Q_n(w)\}$ then follows from the symmetry discussed after (1.19)).

Clearly,

$$\{P_n(z), P_n(w)\} = 0 \Leftrightarrow \{\theta_j, \theta_k\} = 0. \tag{13.1}$$

To see that

$$\{\theta_j, \mu_k\} = \mu_j \delta_{jk} - \mu_j \mu_k \Rightarrow (1.16) \tag{13.2}$$

we first need

Lemma 13.1. For any distinct $z, w \in \mathbb{C}$ and $e^{i\theta} \in \mathbb{D}$,

$$\frac{e^{i\theta} + z}{e^{i\theta} - z} \frac{e^{i\theta} + w}{e^{i\theta} - w} = \frac{z + w}{z - w} \left(\frac{e^{i\theta} + z}{e^{i\theta} - z} - \frac{e^{i\theta} + w}{e^{i\theta} - w} \right) + 1. \tag{13.3}$$

Proof. Consider first $e^{i\theta} = 1$. Then

$$\frac{1 + z}{1 - z} \frac{1 + w}{1 - w} - 1 = \frac{2(z + w)}{(1 - z)(1 - w)}$$

and

$$\frac{1 + z}{1 - z} - \frac{1 + w}{1 - w} = \frac{2(z - w)}{(1 - z)(1 - w)}$$

which implies (13.3) for $e^{i\theta} = 1$.

In that formula, replace z by $e^{-i\theta}z$ and w by $e^{-i\theta}w$ and thereby obtain the formula for general $e^{i\theta}$. \square

Now we compute using (4.9) with $z_j = e^{i\theta_j}$,

$$\begin{aligned} & \{P_n(z), Q_n(w)\} \\ &= \sum_{j,k} -\{z_j, \mu_k\} \frac{P_n(z)}{z - z_j} P_n(w) \frac{w + z_k}{w - z_k} \end{aligned} \tag{13.4}$$

$$\begin{aligned} &= -i \sum_{j,k} z_j [\delta_{jk} \mu_j - \mu_j \mu_k] \frac{P_n(z)}{z - z_j} P_n(w) \frac{w + z_k}{w - z_k} \\ &= -\frac{i}{2} \sum_{j,k} [\delta_{jk} \mu_j - \mu_j \mu_k] \left(\frac{e^{i\theta_j} + z}{e^{i\theta_j} - z} \right) \left(\frac{e^{i\theta_k} + w}{e^{i\theta_k} - w} \right) P_n(z) P_n(w) \end{aligned} \tag{13.5}$$

$$= \text{RHS of (1.19)} \tag{13.6}$$

verifying (13.2). In the above, we get (13.5) by using

$$\frac{z_j}{z - z_j} = \frac{1}{2} \left[\frac{z_j + z}{z_j - z} - 1 \right] \tag{13.7}$$

and noting that the -1 term has j dependence only from $[\delta_{jk} \mu_j - \mu_j \mu_k]$ which sums to zero for each fixed k . To get (13.6), we note that (4.9) says

$$Q_n(z) = - \sum_k \mu_k \left(\frac{e^{i\theta_j} + z}{e^{i\theta_j} - z} \right) P_n(z) \tag{13.8}$$

and use Lemma 13.1 on the $\delta_{jk} \mu_j$ term. The first term in (13.3) gives $\frac{z+w}{z-w} [P_n(z) Q_n(w) - Q_n(z) P_n(w)]$ by (13.8). The second term in (13.3) gives $P_n(z) P_n(w)$ if we note that $\sum_{j,k} \mu_j \delta_{jk} = \sum_j \mu_j = 1$. The $\mu_j \mu_k$ term in (13.5) gives the $-Q_n(z) Q_n(w)$ by (13.8).

Next, we consider Poisson brackets with $F_n(z)$ given by (4.8) and $f_n(z) = z^{-1} [\frac{F_n(z)-1}{F_n(z)+1}]$.

Theorem 13.2. *We have*

$$\begin{aligned} \{P_n(z), F_n(w)\} &= -\frac{i}{2} \left[(P_n(z) m_n(w) + Q_n(z)) \left(\frac{z + w}{z - w} \right) \right. \\ &\quad \left. - P_n(z) - Q_n(z) m_n(w) \right], \end{aligned} \tag{13.9}$$

$$\{Q_n(z), F_n(w)\} = -F_n(w) (\text{RHS of (13.9)}), \tag{13.10}$$

$$\{P_n(z), f_n(w)\} = (1 - wf(w)) W_n(z, w), \tag{13.11}$$

$$\{Q_n(z), f_n(w)\} = -(1 + wf(w)) W_n(z, w), \tag{13.12}$$

where

$$W_n(z, w) = -\frac{i}{2} \left[\frac{(P_n(z) + Q_n(z)) - z(P_n(z) - Q_n(z)) f_n(w)}{z - w} \right]. \tag{13.13}$$

Proof. (12.9) is valid with m_n replaced by F_n . Thus

$$\begin{aligned} \{P_n(z), F_n(w)\} &= -\frac{i}{2} \left[-P_n(w)^{-1} \left[\left(\frac{z+w}{z-w} \right) (P_n(z)Q_n(w) - P_n(w)Q_n(z)) \right. \right. \\ &\quad \left. \left. + P_n(z)P_n(w) - Q_n(z)Q_n(w) \right] \right] \\ &= \text{RHS of (13.9)}. \end{aligned} \tag{13.14}$$

(12.11) is also valid with m_n replaced by F_n , so (13.10) follows from (13.9). For the formula involving f_n , we use (4.27). Analogous to (12.11), one finds

$$\{C_n(z), f_n(w)\} = -wf(w)\{S_n(z), f(w)\} \tag{13.15}$$

and, by (5.2),

$$\{S_n(z), f_n(w)\} = -w^{-1}S_n(w)^{-1}\{S_n(z), C_n(w)\} \tag{13.16}$$

$$= -i \left[\frac{C_n(z) + zS_n(z)f_n(w)}{z-w} \right] \tag{13.17}$$

$$= W(w, z) \tag{13.18}$$

by (4.13)/(4.14).

We then get (13.11) and (13.12) using $P_n = C_n + S_n$, $Q_n = C_n - S_n$ and (13.15), (13.18). \square

Theorem 13.3 (Gekhtman–Nenciu [18]). *We have that*

$$\begin{aligned} \{F_n(z), F_n(w)\} &= -\frac{i}{2} (F_n(z) - F_n(w)) \left[\left(\frac{z+w}{z-w} \right) \right. \\ &\quad \left. (F_n(z) - F_n(w)) + 1 - F_n(z)F_n(w) \right], \end{aligned} \tag{13.19}$$

$$\{f_n(z), f_n(w)\} = -i \frac{f(z) - f(w)}{z-w} (zf(z) - wf(w)). \tag{13.20}$$

Remarks. (1) [18] has $-i$ and $-2i$ where we have $-\frac{i}{2}$ and $-i$ because their $\{ \cdot, \cdot \}$ is twice ours since, following [32], they dropped the normalization of [42] which we keep.

(2) While we will separately derive (13.19) and (13.20), it is an illuminating calculation to go from one to the other using $F(z) = (1 + zf(z))/(1 - zf(z))$.

Proof. By the same calculation that led to (12.13),

$$\{F_n(z), F_n(w)\} = -\frac{Q_n(z)}{P_n(w)P_n(z)^2} \{P_n(z), Q_n(w)\} - (z \leftrightarrow w) \tag{13.21}$$

$$\begin{aligned} &= -\frac{i}{2} F_n(z) \left[\left(\frac{z+w}{z-w} \right) (F_n(z) - F_n(w)) \right. \\ &\quad \left. + 1 - F_n(z)F_n(w) \right] - (z \leftrightarrow w) \\ &= \text{RHS of (13.19)}, \end{aligned} \tag{13.22}$$

where (13.22) follows from (1.19).

Similarly, by (4.27) and the same calculation that led to (12.13),

$$\begin{aligned}
 \{f_n(z), f_n(w)\} &= -(zw)^{-1} \frac{C_n(z)}{S_n(z)^2 S_n(w)} \{S_n(z), C_n(w)\} - (z \leftrightarrow w) \\
 &= -f_n(z) \frac{\{C_n(z), S_n(w)\}}{w S_n(z) S_n(w)} - (z \leftrightarrow w) \\
 &= \frac{i f_n(z)}{z - w} \left[\frac{C_n(z)}{S_n(z)} - \frac{z}{w} \frac{C_n(w)}{S_n(w)} \right] - (z \leftrightarrow w) \\
 &= -\frac{i f_n(z)}{z - w} \left[z f_n(z) - \frac{z}{w} w f_n(w) \right] - (z \leftrightarrow w) \\
 &= -\frac{i z f_n(z)}{z - w} (f_n(z) - f_n(w)) - (z \leftrightarrow w) \\
 &= \text{RHS of (13.20)}. \quad \square
 \end{aligned}
 \tag{13.23}$$

Finally, we want to note that one can compute $\{\Phi_n(z), \Psi_n(z)\}$, $\{\Phi_n^*(z), \Phi_n^*(w)\}$, and $\{\Phi_n(z), \Phi_n^*(w)\}$, $\{\Psi_n(z), \Psi_n^*(w)\}$. We did not know $\{\Phi_n(z), \Psi_n^*(z)\}$ or $\{\Psi_n(z), \Phi_n^*(z)\}$. Nenciu [41], motivated by a preliminary version of this paper, has found this last missing bracket. With her formulae, one can get formulae for the brackets of the elements of the transfer matrix.

Theorem 13.4. *We have*

$$\begin{aligned}
 \{\Phi_n(z), \Phi_n(w)\} &= \{\Psi_n(z), \Psi_n(w)\} = \{\Phi_n^*(z), \Phi_n^*(w)\} \\
 &= \{\Psi_n^*(z), \Psi_n^*(w)\} = 0
 \end{aligned}
 \tag{13.24}$$

$$\begin{aligned}
 \{\Phi_n^*(z), \Psi_n^*(w)\} &= -\frac{i}{2} \left[(\Phi_n^*(z)\Psi_n^*(w) - \Psi_n^*(z)\Phi_n^*(w)) \right. \\
 &\quad \left. \left(\frac{z+w}{z-w} \right) - \Phi_n^*(z)\Phi_n^*(w) + \Psi_n^*(z)\Psi_n^*(w) \right]
 \end{aligned}
 \tag{13.25}$$

$$\begin{aligned}
 \{\Phi_n(z), \Psi_n(w)\} &= -\frac{i}{2} \left[(\Phi_n(z)\Psi_n(w) - \Psi_n(z)\Phi_n(w)) \right. \\
 &\quad \left. \left(\frac{z+w}{z-w} \right) + \Phi_n(z)\Phi_n(w) - \Psi_n(z)\Psi_n(w) \right].
 \end{aligned}
 \tag{13.26}$$

Proof. (1.18) and (1.19) depend on β and are quadratic polynomials in $\bar{\beta}$. Equality for all $\bar{\beta} \in \partial\mathbb{D}$ implies equality for all β , and so equality of the coefficients of the terms multiplying $\bar{\beta}\bar{\beta}$, $\bar{\beta}$, and 1. The $\bar{\beta}\bar{\beta}$ yield the three Φ_n^* , Ψ_n^* Poisson brackets and the terms with no β yield the three Φ_n , Ψ_n Poisson brackets. \square

Remarks. (1) The no β terms involve $z\Phi_n(z)$ and $w\Psi_n(w)$, but zw factors out. The change of sign between (13.25) and (13.26) comes from the minus sign in (4.1) vs. the plus sign in (4.2).

(2) It is an interesting exercise to use $\Phi_n^*(z) = z^n \overline{\Phi_n(1/\bar{z})}$ to go from (13.26) to (13.25). Since Poisson brackets of real functions are real, $\{\bar{f}, \bar{g}\} = \{f, g\}$. We get a minus sign

from $\overline{-i/2} = i/2$, explaining the sign changes in the second and third terms. Because $\frac{1/\bar{z}+1/\bar{w}}{1/\bar{z}-1/\bar{w}} = -\frac{z+w}{z-w}$, the first term does not change.

Theorem 13.5. *We have that*

$$\{\Phi_n(z), \Phi_n^*(w)\} = i w \left(\frac{\Phi_n(z)\Phi_n^*(w) - \Phi_n(w)\Phi_n^*(z)}{z - w} \right). \tag{13.27}$$

Remarks. (1) The same formula holds if Φ_n, Φ_n^* is replaced by Ψ_n, Ψ_n^* (since $\alpha_n \rightarrow -\alpha_n$ preserves $\{\cdot, \cdot\}$).

(2) The proof provides another proof of $\{\Phi_n(z), \Phi_n(w)\} = \{\Phi_n^*(z), \Phi_n^*(w)\} = 0$.

Proof. Define

$$\tilde{C}_n(z) = z\Phi_n(z), \quad \tilde{S}_n(z) = \Phi_n^*(z). \tag{13.28}$$

Then Szegő recursion becomes

$$\tilde{C}_{n+1}(z) = z\tilde{C}_n(z) - \bar{\alpha}_n z\tilde{S}_n(z), \tag{13.29}$$

$$\tilde{S}_{n+1}(z) = \tilde{S}_n(z) - \alpha_n \tilde{C}_n(z), \tag{13.30}$$

which has the same structure as (4.16) and (4.17) except that $\alpha_0, \alpha_1, \dots$ is replaced by $\bar{\alpha}_0, \bar{\alpha}_1, \dots, \bar{\alpha}_n$. Moreover, at $n = 0$,

$$\tilde{C}_n(z)w\tilde{S}_n(w) - \tilde{C}_n(w)z\tilde{S}_n(z) = zw - zw = 0.$$

So we can start the induction. Thus, by induction as in Section 5 ($-i$ becomes the $+i$ because of the complex conjugate change which flips signs of $\{\alpha_n, \bar{\alpha}_n\}$ to $\{\bar{\alpha}_n, \alpha_n\}$),

$$\{\tilde{C}_n(z), \tilde{S}_n(w)\} = i \left(\frac{\tilde{C}_n(z)w\tilde{S}_n(w) - \tilde{C}_n(w)z\tilde{S}_n(z)}{z - w} \right)$$

which is equivalent to (13.27). \square

Using the recursion relations for Φ_n in terms of Φ_{n-1} , one obtains:

Theorem 13.6. *We have*

$$\{\Phi_{n-1}(z), \Phi_n(w)\} = -i\bar{\alpha}_{n-1}w \left(\frac{\Phi_{n-1}(z)\Phi_{n-1}^*(w) - \Phi_{n-1}(w)\Phi_{n-1}^*(z)}{z - w} \right), \tag{13.31}$$

$$\{\Phi_n(z), \Phi_{n-1}^*(w)\} = -izw \left(\frac{\Phi_{n-1}(z)\Phi_{n-1}^*(w) - \Phi_{n-1}(w)\Phi_{n-1}^*(z)}{z - w} \right). \tag{13.32}$$

Acknowledgments

It is a pleasure to thank Rafael Hernández Heredero and Irina Nenciu for useful discussions. M. J. Cantero would like to thank Tom Tombrello and Gary Lorden for the hospitality of Caltech.

George Lorentz was a giant of approximation theory. It is a pleasure and an honor to dedicate this paper to his memory.

References

- [1] M.J. Ablowitz, J.F. Ladik, Nonlinear differential-difference equations, *J. Math. Phys.* 15 (1975) 598–603.
- [2] G.S. Ammar, W.B. Gragg, Schur flows for orthogonal Hessenberg matrices, in: *Hamiltonian and Gradient Flows, Algorithms and Control*, in: *Fields Institute Communications*, vol. 3, American Mathematical Society, Providence, RI, 1994, pp. 27–34.
- [3] D. Barrios, R. Hernández, On the relation between the complex Toda and Volterra lattices, Preprint, 2006.
- [4] M.J. Cantero, L. Moral, L. Velázquez, Measures and para-orthogonal polynomials on the unit circle, *East J. Approx.* 8 (2002) 447–464.
- [5] M.J. Cantero, L. Moral, L. Velázquez, Measures on the unit circle and unitary truncations of unitary operators, *J. Approx. Theory* 139 (2006) 430–468.
- [6] T.S. Chihara, *An Introduction to Orthogonal Polynomials*, in: *Mathematics and Its Applications*, vol. 13, Gordon and Breach, New York, London, Paris, 1978.
- [7] E. Date, S. Tanaka, Analogue of inverse scattering theory for the discrete Hill’s equation and exact solutions for the periodic Toda lattice, *Progr. Theoret. Phys.* 56 (1976) 457–465.
- [8] E.B. Davies, B. Simon, Eigenvalue estimates for non-normal matrices and the zeros of random orthogonal polynomials on the unit circle, *J. Approx. Theory* 141 (2006) 189–213.
- [9] P. Deift, *Integrable Hamiltonian systems*, in: *Dynamical Systems and Probabilistic Methods in Partial Differential Equations* (Berkeley, 1994), in: *Lectures in Applied Mathematics*, vol. 31, American Mathematical Society, Providence, RI, 1996, pp. 103–138.
- [10] P. Deift, L.C. Li, C. Tomei, Toda flows with infinitely many variables, *J. Funct. Anal.* 64 (1985) 358–402.
- [11] B.A. Dubrovin, V.B. Matveev, S.P. Novikov, Non-linear equations of Korteweg–de Vries type, finite-zone linear operators, and Abelian varieties, *Russian Math. Surv.* 31 (1) (1976) 59–146.
- [12] I. Dumitriu, A. Edelman, Matrix models for beta ensembles, *J. Math. Phys.* 43 (2002) 5830–5847.
- [13] L. Faybusovich, M. Gekhtman, On Schur flows, *J. Phys. A* 32 (1999) 4671–4680.
- [14] L. Faybusovich, M. Gekhtman, Poisson brackets on rational functions and multi-Hamiltonian structure for integrable lattices, *Phys. Lett. A* 272 (2000) 236–244.
- [15] H. Flaschka, The Toda lattice, I. Existence of integrals, *Phys. Rev. B* 9 (3) (1974) 1924–1925.
- [16] P.J. Forrester, E.M. Rains, Jacobians and rank 1 perturbations relating to unitary Hessenberg matrices, *Int. Math. Res. Not.* (2006) Article ID 48306, 36 pp.
- [17] G. Freud, *Orthogonal Polynomials*, Pergamon Press, Oxford, New York, 1971.
- [18] M. Gekhtman, I. Nenciu, Multi-Hamiltonian structures for the defocusing Ablowitz–Ladik equation, *Comm. Pure Appl. Math.* (in press).
- [19] F. Gesztesy, J.S. Geronimo, H. Holden, Algebro-geometric solutions of a discrete system related to the trigonometric moment problem, *Comm. Math. Phys.* 258 (2005) 149–177.
- [20] L. Golinskii, Quadrature formula and zeros of para-orthogonal polynomials on the unit circle, *Acta Math. Hungar.* 96 (2002) 169–186.
- [21] L. Golinskii, Schur flows and orthogonal polynomials on the unit circle, *Mat. Sb.* 197 (2006) 41–62.
- [22] Yu.A. Grigoryev, A.V. Tsiganov, On the Darboux–Nijenhuis variables for the open Toda lattice, *SIGMA Symmetry Integrability Geom. Methods Appl.* 2 (2006) Paper 097, 15 pp. (electronic).
- [23] U. Helmke, P.A. Fuhrmann, Bezoutians, *Linear Algebra Appl.* 122/123/124 (1989) 1039–1097.
- [24] M.E.H. Ismail, *Classical and Quantum Orthogonal Polynomials in One Variable*, in: *Encyclopedia of Mathematics and its Applications*, vol. 98, Cambridge University Press, Cambridge, 2005.
- [25] W.B. Jones, O. Njåstad, W.J. Thron, Moment theory, orthogonal polynomials, quadrature, and continued fractions associated with the unit circle, *Bull. London Math. Soc.* 21 (1989) 113–152.
- [26] M. Kac, P. van Moerbeke, On an explicitly soluble system of nonlinear differential equations related to certain Toda lattices, *Adv. Math.* 16 (1975) 160–169.
- [27] M. Kac, P. van Moerbeke, On some periodic Toda lattices, *Proc. Nat. Acad. Sci. USA* 72 (1975) 1627–1629.
- [28] M. Kac, P. van Moerbeke, A complete solution of the periodic Toda problem, *Proc. Nat. Acad. Sci. USA* 72 (1975) 2879–2880.
- [29] F. Kako, N. Mugibayashi, Complete integrability of general nonlinear differential-difference equations solvable by the inverse method. I, *Progr. Theoret. Phys.* 60 (1978) 975–984.
- [30] F. Kako, N. Mugibayashi, Complete integrability of general nonlinear differential-difference equations solvable by the inverse method. II, *Progr. Theoret. Phys.* 61 (1979) 776–790.
- [31] R. Killip, I. Nenciu, Matrix models for circular ensembles, *Int. Math. Res. Not.* 2004 (50) (2004) 2665–2701.
- [32] R. Killip, I. Nenciu, CMV: The unitary analogue of Jacobi matrices, *Comm. Pure Appl. Math.* 60 (2007) 1148–1188.

- [33] B. Kostant, The solution to a generalized Toda lattice and representation theory, *Adv. Math.* 34 (1979) 195–338.
- [34] I.M. Krichever, Algebraic curves and non-linear difference equations, *Russian Math. Surv.* 33 (4) (1978) 255–256.
- [35] P.P. Kulish, Quantum difference nonlinear Schrödinger equation, *Lett. Math. Phys.* 5 (1981) 191–197.
- [36] L.-C. Li, Some remarks on CMV matrices and dressing orbits, *Int. Math. Res. Not.* (40) (2005) 2437–2446.
- [37] J. Moser, Finitely many mass points on the line under the influence of an exponential potential—an integrable system, in: *Dynamical Systems, Theory and Applications (Rencontres, Battelle Res. Inst., Seattle, 1974)*, in: *Lecture Notes in Physics*, vol. 38, Springer, Berlin, 1975, pp. 467–497.
- [38] D. Mumford, An algebro-geometric construction of commuting operators and of solutions to the Toda lattice equation, Korteweg–de Vries equation and related non-linear equations, in: *International Symposium on Algebraic Geometry*, Kyoto, 1977, pp. 115–153.
- [39] I. Nenciu, Lax pairs for the Ablowitz–Ladik system via orthogonal polynomials on the unit circle, Ph.D. Dissertation, California Institute of Technology, 2005.
- [40] I. Nenciu, Lax pairs for the Ablowitz–Ladik system via orthogonal polynomials on the unit circle, *Int. Math. Res. Not.* (11) (2005) 647–686.
- [41] I. Nenciu, Poisson brackets for orthogonal polynomials on the unit circle, Preprint.
- [42] I. Nenciu, B. Simon, unpublished; appeared as Section 11.11 of [47].
- [43] F. Peherstorfer, On Toda lattices and orthogonal polynomials, *J. Comput. Appl. Math.* 133 (2001) 519–534.
- [44] B. Simon, Representations of Finite and Compact Groups, in: *Graduate Studies in Mathematics*, vol. 10, American Mathematical Society, Providence, RI, 1996.
- [45] B. Simon, OPUC on one foot, *Bull. Amer. Math. Soc.* 42 (2005) 431–460.
- [46] B. Simon, Orthogonal Polynomials on the Unit Circle, Part 1: Classical Theory, in: *AMS Colloquium Series*, American Mathematical Society, Providence, RI, 2005.
- [47] B. Simon, Orthogonal Polynomials on the Unit Circle, Part 2: Spectral Theory, in: *AMS Colloquium Series*, American Mathematical Society, Providence, RI, 2005.
- [48] B. Simon, Zeros of OPUC and long time asymptotics of Schur and related flows, *Inverse Problems and Imaging* 1 (2007) 155–181.
- [49] B. Simon, CMV matrices: Five years after, *J. Comput. Appl. Math.* 208 (2007) 120–154.
- [50] B. Simon, Rank one perturbations and the zeros of paraorthogonal polynomials on the unit circle, *J. Math. Anal. Appl.* 329 (2007) 376–382.
- [51] B. Simon, Szegő’s Theorem and Its Descendants: Spectral Theory for L^2 Perturbations of Orthogonal Polynomials, to be published by Princeton University Press, Princeton, NJ (in preparation).
- [52] G. Szegő, Orthogonal Polynomials, American Mathematical Society Colloquium Publications, vol. 23, American Mathematical Society, Providence, RI, 1939; 3rd edition, 1967.
- [53] A.V. Tsiganov, A family of the Poisson brackets compatible with the Sklyanin bracket, *J. Phys. A* 40 (2007) 4803–4816.
- [54] P. van Moerbeke, The spectrum of Jacobi matrices, *Invent. Math.* 37 (1976) 45–81.
- [55] P. van Moerbeke, D. Mumford, The spectrum of difference operators and algebraic curves, *Acta Math.* 143 (1979) 93–154.
- [56] M.-W.L. Wong, First and second kind paraorthogonal polynomials and their zeros, *J. Approx. Theory.* 146 (2007) 282–293.