



Natural boundaries and spectral theory

Jonathan Breuer^a, Barry Simon^{b,*}

^a *Einstein Institute of Mathematics, The Hebrew University of Jerusalem, Jerusalem 91904, Israel*

^b *Mathematics 253-37, California Institute of Technology, Pasadena, CA 91125, USA*

Received 9 August 2010; accepted 21 December 2010

Available online 30 December 2010

Communicated by Charles Fefferman

Abstract

We present and exploit an analogy between lack of absolutely continuous spectrum for Schrödinger operators and natural boundaries for power series. Among our new results are generalizations of Hecke's example and natural boundary examples for random power series where independence is not assumed.

© 2011 Elsevier Inc. All rights reserved.

MSC: 30B30; 35P05; 30B20

Keywords: Reflectionless; Right limit; Natural boundary

1. Introduction

In this paper, we'll present and exploit a powerful analogy between spectral theory and the question of when a power series $f(z) = \sum_{n=0}^{\infty} a_n z^n$ defining an analytic function on $\mathbb{D} = \{z \mid |z| < 1\}$ has a natural boundary on $\partial\mathbb{D}$, in that for no $z_0 = e^{i\theta}$ does f have an analytic continuation to $\{z \mid |z - z_0| < \delta\}$ for some $\delta > 0$. In particular, we shall import two notions ("reflectionless" and "right limit") from the spectral theory of Jacobi matrices and obtain a general theorem regarding the consequence of the possibility to analytically continue f across an arc. While spectral theory ideas motivated our approach to natural boundaries, what we develop doesn't require any spectral theory. The reader not knowledgeable in spectral theory and not interested in the background should skip to the paragraph containing (1.4).

* Corresponding author.

E-mail addresses: jbreuer@math.huji.ac.il (J. Breuer), bsimon@caltech.edu (B. Simon).

¹ Supported in part by NSF grant DMS-0652919.

As it turns out, some of the results presented in this paper are not new. In particular, in [1] Agmon treated natural boundaries for general series with radius of convergence 1 using a related approach. Our Theorem 1.3 is a special case of his results. To the best of our knowledge, however, our treatment of *strong* natural boundaries using this approach (see Theorem 1.4 below) is completely new.

Of course, since the relevant spectral theoretic notions were not yet defined at the time of publication of [1], Agmon was unaware of the analogy. Surprisingly, [1] is rarely quoted in the relevant literature and seems to be little known. Thus, a secondary aim of our paper is to draw attention to Agmon’s work.

Moreover, we believe the significance of the analogy presented in this paper is more than merely anecdotal. As an example, our knowledge of the applications of these ideas to random potentials in the spectral theoretic setting, led us to apply these methods to random series. In this context we have obtained results for series where only a subsequence is random and for general bounded ergodic nondeterministic series. Another example is Theorem 1.8 that has Hecke’s famous example (1.14) as a special case.

Thus, Theorems 1.3 and 1.4 provide a general framework in which many existing and new results concerning natural boundaries can be derived.

To set the stage, consider a Jacobi matrix,

$$J = \begin{pmatrix} b_1 & a_1 & 0 & \dots \\ a_1 & b_2 & a_2 & \dots \\ 0 & a_2 & b_3 & \dots \\ \vdots & \vdots & \vdots & \ddots \end{pmatrix} \tag{1.1}$$

where $\{a_n, b_n\}_{n=1}^\infty$ are bounded. By a right limit of J , we mean a two-sided Jacobi matrix, $J^{(r)}$, with parameters $\{a_n^{(r)}, b_n^{(r)}\}_{n=-\infty}^\infty$ given by

$$a_n^{(r)} = \lim_{k \rightarrow \infty} a_{n+n_k}, \quad b_n^{(r)} = \lim_{k \rightarrow \infty} b_{n+n_k} \tag{1.2}$$

where $n_k \rightarrow \infty$ is a subsequence. By compactness, right limits exist. This definition is from Last and Simon [23], who used it in

Theorem 1.1. (See [23].) *If $\Sigma_{ac}(A)$ is the essential support of the a.c. spectrum of an operator, A , then*

$$\Sigma_{ac}(J) \subset \Sigma_{ac}(J^{(r)}) \tag{1.3}$$

for every right limit $J^{(r)}$ of J .

Right limits are also relevant to essential spectrum where they have been exploited by several authors (e.g., [10,11,18,23–25,35]), but their relevance for a.c. spectrum goes back to Last and Simon [23]. Recently, Remling [36] found a much stronger property of right limits when there is a.c. spectrum. It depends on the notion of reflectionless two-sided Jacobi matrix. The precise definition is irrelevant for our discussion here (see, e.g., [5,36,44,45]), but we note that if \tilde{J} is a two-sided Jacobi matrix, reflectionless on some $\epsilon \subset \mathbb{R}$ with the Lebesgue measure, $|\epsilon|$, of ϵ nonzero, then $\epsilon \subset \Sigma_{ac}(\tilde{J})$ and $\{\tilde{a}_n, \tilde{b}_n\}_{n=-\infty}^{-1}$ determine $\{\tilde{a}_n, \tilde{b}_n\}_{n=0}^\infty$. Remling [36] proved

Theorem 1.2. (See [36].) *If ϵ is $\Sigma_{ac}(J)$ for a Jacobi matrix, then any right limit, $J^{(\epsilon)}$, is reflectionless on ϵ .*

With this result, Remling was able to recover and extend virtually every result on the absence of a.c. spectrum for situations where $\{a_n, b_n\}_{n=1}^\infty$ are bounded, and the lack of a.c. spectrum is due to a part of the a 's and b 's that is dominant at infinity (rather than a perturbation that goes to zero at infinity).

Our work here began by our noticing that the major classes of Jacobi matrices with no a.c. spectrum have analogs in the major classes of results on the occurrence of natural boundaries (see Remmert [37, Chapter 11] for a summary of classical results on natural boundaries), as seen in

- Gap theorems [19,17,16,37] \sim sparse potentials [32,36].
- Finite-valued power series [47,37] \sim finite-valued Jacobi matrices [22,36].
- Random power series [46,30,21] \sim Anderson localization [8,31].

Let us describe our major abstract results on natural boundaries motivated by Last and Simon [23] and Remling [36]. Given a power series $f(z) = \sum_{n=0}^\infty a_n z^n$ with

$$\sup_n |a_n| < \infty \tag{1.4}$$

we define a right limit of $\{a_n\}_{n=0}^\infty$ to be a two-sided sequence $\{b_n\}_{n=-\infty}^\infty$ with

$$b_n = \lim_{j \rightarrow \infty} a_{n+n_j} \tag{1.5}$$

for some $n_j \rightarrow \infty$. By compactness and (1.4), right limits exist.

Given a two-sided bounded sequence, $\{b_n\}_{n=-\infty}^\infty$, we consider two functions, $f_+(z)$ on \mathbb{D} and $f_-(z)$ on $\mathbb{C} \cup \{\infty\} \setminus \overline{\mathbb{D}}$, defined by

$$f_+(z) = \sum_{n=0}^\infty b_n z^n, \quad f_-(z) = \sum_{n=-\infty}^{-1} b_n z^n \tag{1.6}$$

where the series are guaranteed to converge on the indicated sets. Let I be an open interval in $\partial\mathbb{D}$. We say $\{b_n\}_{n=-\infty}^\infty$ is *reflectionless* on I if and only if f_+ has an analytic continuation from \mathbb{D} to $\mathbb{C} \cup \{\infty\} \setminus (\partial\mathbb{D} \setminus I)$, so that on $\mathbb{C} \cup \{\infty\} \setminus \overline{\mathbb{D}}$, we have that

$$f_+(z) + f_-(z) = 0. \tag{1.7}$$

Obviously, it suffices that f_+ have a continuation to a neighborhood of I so that (1.7) holds in the intersection of that neighborhood and $\mathbb{C} \setminus \overline{\mathbb{D}}$.

Example. Let $b_n \equiv 1$. Then $f_+(z) = (1 - z)^{-1}$ and $f_-(z) = -(1 - z)^{-1}$. This series is reflectionless on $I = \{e^{i\theta} \mid 0 < \theta < 2\pi\}$. Similarly, it is easy to see that a periodic b_n of period p is reflectionless on any I in $\partial\mathbb{D}$ with all the p -th roots of unity removed.

Our first main theorem is:

Theorem 1.3. *Let $f(z) = \sum_{n=0}^{\infty} a_n z^n$ be a power series with (1.4). Suppose $I \subset \partial\mathbb{D}$ is an open interval so that $f(z)$ has an analytic continuation to a neighborhood of I . Then every right limit of $\{a_n\}_{n=0}^{\infty}$ is reflectionless on I .*

As remarked above, this theorem is a special case of the results in [1]. While our treatment is limited to bounded sequences and produces interesting results only when $a_n \not\rightarrow 0$, Agmon treats general (not necessarily bounded) series. He introduces a way to renormalize the coefficients so that any series with radius of convergence 1 has, after renormalization, nontrivial right limits. He then shows that if the corresponding function has an analytic continuation to an arc on $\partial\mathbb{D}$, then the renormalized right limits are reflectionless across that arc. For bounded series, we will actually prove a stronger result (we state Theorem 1.3 both for conceptual reasons and because we need it in the proof of the stronger result):

Theorem 1.4. *Let $f(z) = \sum_{n=0}^{\infty} a_n z^n$ be a power series with (1.4). Suppose $I \subset \partial\mathbb{D}$ is an open interval so that*

$$\sup_{\substack{0 < r < 1 \\ e^{i\theta} \in I}} \int |f(re^{i\theta})| \frac{d\theta}{2\pi} < \infty. \tag{1.8}$$

Then every right limit of $\{a_n\}$ is reflectionless on I .

These theorems imply natural boundaries. We say f has a *strong natural boundary* on $\partial\mathbb{D}$ if (1.8) fails for every $I \subset \partial\mathbb{D}$. In particular, in that case, f is unbounded in every sector $\{re^{i\theta} \mid 0 < r < 1, e^{i\theta} \in I\}$.

Corollary 1.5. *Let $f(z) = \sum_{n=0}^{\infty} a_n z^n$ be such that for any open interval $I \subset \partial\mathbb{D}$, there is a right limit of $\{a_n\}_{n=0}^{\infty}$ which is not reflectionless on I . Then f has a strong natural boundary on $\partial\mathbb{D}$.*

Of course, if $a_n \rightarrow 0$, it can happen that there are natural boundaries which are not strong natural boundaries. For example, by the Hadamard gap theorem [37],

$$f(z) = \sum_{n=1}^{\infty} \frac{z^{n!}}{(n!)^n} \tag{1.9}$$

has a natural boundary on $\partial\mathbb{D}$ but, for all k , $f^{(k)}(z)$ is bounded on \mathbb{D} .

We note that Duffin and Schaeffer [13] and Boas [4] long ago had results on what we call strong natural boundaries. These results are all for series with gaps or finite-valued series. These authors and also Agmon [1] have results that prove that certain functions rather than merely having classical natural boundaries are unbounded in every sector (we call this L^∞ natural boundaries). We note that our approach for going from classical to strong natural boundaries is very close to the method that Agmon [1] uses to go from classical natural boundaries to unboundedness in every sector. Theorem 1.4 is, to the best of our knowledge, the first general theorem concerning strong natural boundaries.

For us, the point of these theorems is conceptual: they provide a unified framework that makes many theorems transparent. That said, the following results seem to be new:

Theorem 1.6. *Suppose $\{a_n\}_{n=0}^\infty$ obeys (1.4) and there exists $n_j \rightarrow \infty$ so that for all $k < 0$,*

$$\lim_{j \rightarrow \infty} a_{n_j+k} = 0, \tag{1.10}$$

$$\liminf |a_{n_j}| > 0. \tag{1.11}$$

Then $f(z) = \sum_{n=0}^\infty a_n z^n$ has a strong natural boundary on $\partial\mathbb{D}$.

Remarks.

1. This result (as well as its extension—Theorem 4.3), for L^∞ natural boundaries, appears in Agmon [1].
2. The result is true if (1.10) is replaced by the assumption for all $k > 0$ or for all $k < K$, some $K < 0$ or for all $k > K$, some $K > 0$. The proofs are essentially identical. In addition, (1.10) can be replaced by an exponential decay condition on the limit; see Theorem 4.3.
3. This includes the famous examples $\sum_{n=1}^\infty z^{n!}$ of Weierstrass and $\sum_{n=0}^\infty z^{n^2}$ of Kronecker.
4. This allows gaps where the set of zeros has zero density. At first sight, this seems a violation of the result of Pólya [33] and Erdős [15] that the Fabry gap theorem is optimal. We'll explain this apparent discrepancy in Section 4.

Theorem 1.7. *Let $\{a_n(\omega)\}_{\omega \in \Omega}$ be a translation invariant, ergodic, stochastic process, which is nondeterministic, so that*

$$\sup_{n, \omega} |a_n(\omega)| < \infty. \tag{1.12}$$

Then for a.e. ω , $\sum_{n=0}^\infty a_n(\omega)z^n$ has a strong natural boundary.

Remarks.

1. So far as we know, all previous results on random power series rely on independence and only obtain natural boundaries, not strong natural boundaries.
2. Recall that a stochastic process $\{a_n(\omega)\}$ is deterministic if $a_0(\omega)$ is a measurable function of $\{a_n(\omega)\}_{n=-\infty}^{-1}$ for a.e. ω , and nondeterministic if it is not deterministic.
3. By taking an average of a deterministic and a nondeterministic process, it is easy to see that ergodicity is essential for this theorem to be true.

Theorem 1.8. *Let $f: \partial\mathbb{D} \rightarrow \mathbb{C}$ be a bounded and piecewise continuous function with only a finite number of discontinuities, at one of which the one-sided limits exist and are unequal. Then for any irrational number q and every $\theta \in \mathbb{R}$, we have that*

$$\sum_{n=0}^\infty f(e^{2\pi i(qn+\theta)})z^n \tag{1.13}$$

has a strong natural boundary.

Remark. This includes Hecke’s famous example [20],

$$f(z) = \sum_{n=0}^{\infty} \{nq\}z^n \tag{1.14}$$

where $\{x\}$ is the fractional part of $x \in \mathbb{R}$.

The following is elementary and does not require our full machinery. It was suggested to us by the Wonderland theorem of Simon [43] and seems to be new.

Theorem 1.9. *Let $\Omega \subset \mathbb{C}$ be a compact set with more than one point. Let Ω^∞ be a countable product of copies of Ω in the weak topology. Then $\{\{a_n\} \in \Omega^\infty \mid \sum_{n=0}^{\infty} a_n z^n \text{ has a natural boundary on } \partial\mathbb{D}\}$ is a dense G_δ in Ω^∞ .*

In Section 2, we prove Theorem 1.3. The key is a lemma of M. Riesz whose proof we include for completeness. In Section 3, we prove Theorem 1.4 using the theory of H^p spaces on a sector (Duren [14]). In Section 4, we discuss gap theorems, including Theorem 1.6. In Section 5, we discuss Szegő’s theorem on finite-valued power series using Theorem 1.4. In Section 6, we discuss random power series, including Theorem 1.7. In Section 7, we prove Theorem 1.8, following a spectral theory analysis of Damanik and Killip [12]. In Section 8, we prove Theorem 1.9.

We believe our work here opens up numerous new directions in the study of power series and of spectral theory. In particular, there is a dynamical view of reflectionless in spectral theory (see [5]) and there is the distinction in spectral theory between pure point and singular continuous spectra. What are the analogs for power series?

2. Classical natural boundaries

In this section, we prove Theorem 1.3. The key will be a lemma of M. Riesz [38] used by many other authors in the study of natural boundaries. The use of right limits and reflectionless power series can be viewed as a tool for squeezing maximum benefit from Riesz’s lemma.

Given a power series $f(z) = \sum_{n=0}^{\infty} a_n z^n$, where the a_n obey (1.4), we define

$$f_+^{(N)}(z) = \sum_{n=0}^{\infty} a_{n+N} z^n, \quad f_-^{(N)}(z) = \sum_{n=-N}^{-1} a_{n+N} z^n \tag{2.1}$$

so for $z \in \mathbb{D} \setminus \{0\}$,

$$f_+^{(N)}(z) + f_-^{(N)}(z) = z^{-N} f(z). \tag{2.2}$$

Clearly, f_- is defined and analytic on $\mathbb{C} \setminus \{0\}$, f_+ is defined initially on \mathbb{D} , but by (2.2), has analytic continuation to any region that f does.

Theorem 2.1 (*M. Riesz’s lemma*). *Suppose $\{a_n\}_{n=0}^{\infty}$ obeys (1.4) and that f has an analytic continuation to a neighborhood of $\mathbb{D} \cup S$ where*

$$S = \{r e^{i\theta} \mid 0 < r \leq R, \alpha \leq \theta \leq \beta\} \tag{2.3}$$

for some $R > 1$ and $\alpha < \beta$. Then

$$\sup_{\substack{z \in S \\ N=0,1,2,\dots}} |f_+^{(N)}(z)| < \infty. \tag{2.4}$$

Remark. Riesz’s lemma is usually in terms of $z^{-N}(f(z) - \sum_{n=0}^{N-1} a_n z^n)$, but this is exactly $f_+^{(N)}(z)$.

Proof of Theorem 2.1. While the result is classical and appears in many places (e.g., [37]), we sketch the proof for completeness.

By comparing with a geometric series, for $z \in \mathbb{D}$ and all N ,

$$|f_+^{(N)}(z)| \leq (1 - |z|)^{-1} \sup_n |a_n| \tag{2.5}$$

and similarly, for $z \in \mathbb{C} \setminus \bar{\mathbb{D}}$,

$$|f_-^{(N)}(z)| \leq (1 - |z|^{-1})^{-1} \sup_n |a_n|. \tag{2.6}$$

Let \tilde{S} have the form of S with α, β replaced by $\tilde{\alpha}, \tilde{\beta}$ and $\tilde{\alpha} < \alpha < \beta < \tilde{\beta}$ so that \tilde{S} lies in the neighborhood of analyticity of f . Let $z_1 = e^{i\tilde{\alpha}}, z_2 = e^{i\tilde{\beta}}$, and define

$$g_{\pm}^{(N)}(z) = (z - z_1)(z - z_2) f_{\pm}^{(N)}(z). \tag{2.7}$$

Clearly, (2.4) is implied by

$$\sup_{\substack{z \in \tilde{S} \\ N=0,1,2,\dots}} |g_+^{(N)}(z)| < \infty. \tag{2.8}$$

By the maximum principle, we need only check this on $\partial\tilde{S} \setminus \{z_1, z_2\}$.

Because of (2.5) and the zeros of $g_+^{(N)}$, we have

$$\sup_{\substack{z \in \partial\tilde{S} \cap \mathbb{D} \\ N=0,1,2,\dots}} |g_+^{(N)}(z)| < \infty. \tag{2.9}$$

Similarly, by (2.6),

$$\sup_{\substack{z \in \partial\tilde{S} \cap \mathbb{C} \setminus \bar{\mathbb{D}} \\ N=0,1,2,\dots}} |g_-^{(N)}(z)| < \infty. \tag{2.10}$$

Since $|z^{-N}| < 1$ on $\mathbb{C} \setminus \bar{\mathbb{D}}$, (2.2) and (2.10) imply

$$\sup_{\substack{z \in \partial \tilde{\mathbb{D}} \cap \mathbb{C} \setminus \mathbb{D} \\ N=0,1,2,\dots}} |g_+^{(N)}(z)| < \infty \tag{2.11}$$

proving (2.8). \square

Proof of Theorem 1.3. Suppose $a_{N_j+n} \rightarrow b_n$ for $n \in \mathbb{Z}$ and let f_{\pm} be the functions in (1.6). Then, by estimating Taylor series, we have

$$f_+^{(N_j)}(z) \rightarrow f_+(z), \quad f_-^{(N_j)}(z) \rightarrow f_-(z) \tag{2.12}$$

uniformly on compact subsets of \mathbb{D} and $\mathbb{C} \cup \{\infty\} \setminus \bar{\mathbb{D}}$, respectively.

If f has an analytic continuation to a neighborhood of $I = \{e^{i\theta} \mid \alpha_0 < \theta < \beta_0\}$, we can apply Riesz’s lemma for any S of the form (2.3) with $\alpha_0 < \alpha < \beta < \beta_0$ and some suitable $R > 1$ (depending on α, β). Thus, by the Vitali convergence theorem, $f_+^{(N_j)}$ converges uniformly on S , so $f_+(z)$ has an analytic continuation to S . Moreover, by the analytic continuation of (2.2) to S and $z^{-N_j} \rightarrow 0$ on $\mathbb{C} \setminus \bar{\mathbb{D}}$, we see on $S \setminus \bar{\mathbb{D}}$,

$$f_+(z) + f_-(z) = 0. \tag{2.13}$$

Thus, $\{b_n\}_{n=-\infty}^{\infty}$ is reflectionless across I . \square

The proof shows that if $\Omega_I = \mathbb{C} \cup \{\infty\} \setminus (\partial \mathbb{D} \setminus I)$ and if \mathcal{R} is the family of all right limits and $f_b(z)$ the function on Ω_I equal to $\sum_{n=0}^{\infty} b_n z^n$ on \mathbb{D} , then

Theorem 2.2. Fix I an interval in $\partial \mathbb{D}$ and $A \in (0, \infty)$. Let \mathcal{R} be the set of two-sided sequences reflectionless across I with

$$\sup_{-\infty < n < \infty} |b_n| \leq A. \tag{2.14}$$

Then, one has that, for any compact $K \subset \Omega_I$,

$$\sup_{b \in \mathcal{R}} \sup_{z \in K} |f_b(z)| < \infty. \tag{2.15}$$

Moreover, \mathcal{R} is compact in the topology of uniform convergence on compacts of Ω_I and on

$$\{\{b_n\}_{n=-\infty}^{-1} \mid \{b_n\}_{n=-\infty}^{\infty} \in \mathcal{R}\}, \tag{2.16}$$

b_0, b_1, \dots are continuous functions of $\{b_n\}_{n=-\infty}^{-1}$. In fact, for any $k < \ell$, there is a homeomorphism of $\{b_n\}_{n=-\infty}^k$ to $\{b_n\}_{n=\ell}^{\infty}$ by associating the two ends of a two-sided sequence.

Proof. If $K \subset \mathbb{D}$ or $K \subset \mathbb{C} \setminus \bar{\mathbb{D}}$, one can use (2.5) or (2.6) for f_b and $\sup_n |b_n|$. For K straddling I , use the argument in the proof of Riesz’s lemma and the fact that f_b is $\sum_{n=0}^{\infty} b_n z^n$ on \mathbb{D} and $-\sum_{n=-\infty}^{-1} b_n z^n$ on $\mathbb{C} \setminus \bar{\mathbb{D}}$. This proves (2.15).

Compactness then follows from Montel’s theorem and the fact that, by these bounds, the set of reflectionless functions is closed. The continuity is immediate if one notes that $\{b_n\}_{n=-\infty}^{-1}$

determines f near ∞ and f near 0 determines $\{b_n\}_{n=0}^\infty$. The “in fact” statement uses that if $\{b_n\}_{n=0}^\infty$ is reflectionless, so are its translates. \square

Since Theorem 1.3 says that if some right limit is not reflectionless, then f has a natural boundary, and since reflectionless $\{b_n\}_{n=-\infty}^\infty$ have one half determining the other half, we have the following, which implies almost all the natural boundary results of this paper:

Theorem 2.3. *Let $\{a_n\}_{n=0}^\infty$ be a bounded sequence with two right limits, $\{b_n\}_{n=-\infty}^\infty$ and $\{c_n\}_{n=-\infty}^\infty$, that obey*

$$b_0 \neq c_0 \tag{2.17}$$

and either for some $k > 0$ and all $j \geq k$ or for some $k < 0$ and all $j \leq k$,

$$b_j = c_j. \tag{2.18}$$

Then $\sum_{n=0}^\infty a_n z^n$ has a natural boundary.

Remarks.

1. For example, if $k > 0$, we look at the right limits $\{b_{n-k}\}_{n=-\infty}^\infty$ and $\{c_{n-k}\}_{n=-\infty}^\infty$ which have the same f_+ ’s but unequal f_- ’s.
2. In the next section, we extend this to conclude strong natural boundaries.

3. Strong natural boundaries

In this section, we prove Theorem 1.4. We suppose that f obeys (1.8) where $I = (\alpha, \beta)$. Since (1.8) implies $\int_0^1 (\int_\alpha^\beta |f(re^{i\theta})| \frac{d\theta}{2\pi}) dr < \infty$, for a.e. θ_0 in (α, β) , we have $\int_0^1 |f(re^{i\theta_0})| dr < \infty$. So, for a sequence $\varepsilon_n \downarrow 0$, we have $f \in E(S_n)$, where $S_n = \{re^{i\theta} \mid 0 < r < 1, \alpha + \varepsilon_n < \theta < \beta - \varepsilon_n\}$ and $E(S_n)$ is the space introduced in [14, Section 10.1]. It follows that:

- (a) $\lim_{r \uparrow 1} f(re^{i\theta}) = f(e^{i\theta})$ exists for a.e. $\theta \in (\alpha, \beta)$ [14, Theorem 10.3].
- (b) For all ε , $\int_{\alpha+\varepsilon}^{\beta-\varepsilon} |f(e^{i\theta})| \frac{d\theta}{2\pi} < \infty$ [14, Theorem 10.3].
- (c) For every $\varepsilon > 0$,

$$\lim_{r \uparrow 1} \int_{\alpha+\varepsilon}^{\beta-\varepsilon} |f(re^{i\theta}) - f(e^{i\theta})| \frac{d\theta}{2\pi} = 0. \tag{3.1}$$

- (d) If we define

$$F(z) = \int_{\alpha+\varepsilon}^{\beta-\varepsilon} f(e^{i\theta})(e^{i\theta} - z)^{-1} \frac{d\theta}{2\pi i} \tag{3.2}$$

then F is analytic in $\mathbb{C} \setminus \{e^{i\theta} \mid \alpha + \varepsilon < \theta < \beta - \varepsilon\}$ and

$$\lim_{r \uparrow 1} F(re^{i\theta}) - \lim_{r \downarrow 1} F(re^{i\theta}) = f(e^{i\theta}) \tag{3.3}$$

(follows from [14, Theorem 10.4]).

We also need the following Painlevé-type theorem, which follows easily from Morera’s theorem:

(e) If f_+ is analytic in \mathbb{D} , f_- in $\mathbb{C} \setminus \bar{\mathbb{D}}$, and

$$\sup_{0 < r < 1} \int_{\alpha}^{\beta} |f_+(re^{i\theta})| \frac{d\theta}{2\pi} + \sup_{1 < r < 2} \int_{\alpha}^{\beta} |f_-(re^{i\theta})| \frac{d\theta}{2\pi} < \infty \tag{3.4}$$

and if for a.e. $\theta \in (\alpha, \beta)$,

$$f_+(e^{i\theta}) = f_-(e^{i\theta}) \tag{3.5}$$

then there is G analytic in $\mathbb{C} \setminus [\partial\mathbb{D} \setminus \{e^{i\theta} \mid \alpha < \theta < \beta\}]$ so that $G = f_+$ on $\partial\mathbb{D}$ and f_- on $\mathbb{C} \setminus \bar{\mathbb{D}}$.

Proof of Theorem 1.4. Let F be given by (3.2) and define

$$b_n = \int_{\alpha + \varepsilon}^{\beta - \varepsilon} e^{-in\theta} f(e^{i\theta}) \frac{d\theta}{2\pi}. \tag{3.6}$$

Then, by expanding $(e^{i\theta} - z)^{-1}$ in suitable geometric series, the Taylor expansion of F near zero is $\sum_{n=0}^{\infty} b_n z^n$ and near ∞ is $\sum_{n=-\infty}^{-1} b_n z^n$. Moreover, by the Riemann–Lebesgue lemma and (b) above,

$$\lim_{|n| \rightarrow \infty} b_n = 0. \tag{3.7}$$

Let $c_n = a_n - b_n$ and

$$f_+(z) = \sum_{n=0}^{\infty} c_n z^n. \tag{3.8}$$

Then, by (a), for a.e. $\theta \in (\alpha + \varepsilon, \beta - \varepsilon)$,

$$\lim_{r \uparrow 1} f_+(re^{i\theta}) = f(e^{i\theta}) - \lim_{r \uparrow 1} F(re^{i\theta}) \tag{3.9}$$

and if

$$f_-(z) = - \sum_{n=-\infty}^{-1} b_n z^n \tag{3.10}$$

then, for a.e. $\theta \in (\alpha + \varepsilon, \beta - \varepsilon)$,

$$\lim_{r \downarrow 1} f_-(z) = - \lim_{r \downarrow 1} F(re^{i\theta}). \tag{3.11}$$

It follows that (3.5) holds, so f_+ has a classical analytic continuation across $(\alpha + \varepsilon, \beta - \varepsilon)$. By Theorem 1.3, every right limit of c_n is reflectionless on $(\alpha + \varepsilon, \beta - \varepsilon)$.

But, by (3.7), the right limits of c_n and a_n are the same! Thus, each right limit of a_n is reflectionless on each $(\alpha + \varepsilon, \beta - \varepsilon)$, and so on (α, β) . \square

Theorem 3.1. *If the hypotheses of Theorem 2.3 hold, then $\sum_{n=0}^{\infty} a_n z^n$ has a strong natural boundary on $\partial\mathbb{D}$.*

Example 3.2. The Rudin and Shapiro [39,42] sequence is defined by defining polynomials P_n and Q_n recursively by $P_0(z) = Q_0(z) = 1$, $P_{n+1}(z) = P_n(z) + z^{2^n} Q_n(z)$, $Q_{n+1}(z) = P_n(z) - z^{2^n} Q_n(z)$. As power series with bounded coefficients, $\lim P_n(z)$ exists and defines a series $f(z) = \sum_{n=0}^{\infty} a_n z^n$, where each a_n is ± 1 . By using Szegő’s theorem, Brilhart [7] proved this function had a natural boundary. Here is an elementary direct proof. P_{n+1} is $P_{n-1} Q_{n-1} P_{n-1}(-Q_{n-1})$. Taking right limits at the end of the P_{n-1} in $P_{n-1} Q_{n-1}$ and in $P_{n-1}(-Q_{n-1})$ yields right limits which agree at negative index but have opposite signs at positive index.

4. Gap theorems

In this section, we’ll prove Theorem 1.6 and resolve the apparent contradiction to Pólya [33] and Erdős [15]. The following proof shows the power of reflectionless theorems.

Proof of Theorem 1.6. By compactness, we can find a subsubsequence, n_{j_ℓ} , so $\lim_{\ell \rightarrow \infty} a_{n_{j_\ell} + k} = b_k$ exists for all k and $b_0 \neq 0$, $b_k = 0$ for $k < 0$. For such a right limit, $f_-(z) = 0$, but $f_+(0) = b_0 \neq 0$. Thus, f_- cannot be an analytic continuation of f_+ through any I , and this $\{b_n\}_{n=-\infty}^{\infty}$ is not reflectionless on any I . Theorem 1.4 completes the proof. \square

It is simplest to resolve the apparent contradiction with [33,15] in the context of an example.

Example 4.1. Let $U = \bigcup_{j=2}^{\infty} \{n \mid j! \leq n \leq j! + j\}$. We want to consider bounded power series with $a_n = 0$ if $n \in U$. If $a_{j!+j+1} = 1$ for $j \geq 2$ and a_n is arbitrary but bounded for $n \notin U \cup \{j! + j + 1\}_{j=2}^{\infty}$, then $f(z)$ has a strong natural boundary on $\partial\mathbb{D}$. The zero values may be only on U , which is a set with zero density.

On the other hand, [33,15] say that since U does not have density 1, there must be $\{a_n\}_{n=1}^{\infty}$ with $a_n = 0$ for $n \in U$ so that $\partial\mathbb{D}$ is not a natural boundary.

The resolution is that our natural boundary examples have hard edges, that is, a_n jumps at the edges of U , while the examples of [15] have soft edges. $\mathbb{Z}_+ \setminus U$ has longer and longer nonzero intervals and Erdős’ examples ramp up slowly and down slowly to be 1 in the center of

these intervals. It is easy to see that these examples have right limits which are constant, and so reflectionless!

These examples of Erdős are reminiscent of the sparse potentials of Molchanov [27] and Remling [36], where approximate solitons are placed in between long gaps. In fact, given the chronology, we should say the examples of [27,36] are reminiscent of Erdős [15]!

As noted by Agmon [1,2], it isn't important that $a_{N_j+n} \rightarrow 0$ as $j \rightarrow \infty$ from gaps, only that its lim sup decays exponentially fast. Using right limits, this is easy to see since

Theorem 4.2. *Let $\{b_n\}_{n=-\infty}^{\infty}$ be a two-sided series which is reflectionless on some interval, $I \subset \partial\mathbb{D}$. Suppose that for some $C, D > 0$, we have that*

$$|b_n| \leq C e^{-Dn} \quad \text{for } n > 0. \tag{4.1}$$

Then $b_n \equiv 0$.

Proof. By (4.1), $f_+(z)$ has an analytic continuation to the circle $\{z \mid |z| < e^D\}$. Since $f_+ = -f_-$ in a neighborhood of I , we conclude that f_+ defines an entire function. Since $|f_-(z)| \rightarrow 0$ as $|z| \rightarrow \infty$, $f_+ \equiv 0$, so $f_- \equiv 0$ also, and then $b_n \equiv 0$. \square

This immediately implies the following extension of Theorem 1.6:

Theorem 4.3. *Suppose $\{a_n\}_{n=0}^{\infty}$ obeys (1.4) and there exists $n_j \rightarrow \infty$ so that for some $C, D > 0$ and for all $k < 0$,*

$$\limsup_{j \rightarrow \infty} |a_{n_j+k}| \leq C e^{-D|k|}, \tag{4.2}$$

$$\liminf |a_{n_j}| > 0. \tag{4.3}$$

Then $f(z) = \sum_{n=0}^{\infty} a_n z^n$ has a strong natural boundary on $\partial\mathbb{D}$.

Proof. By compactness, there exists a right limit, so (4.1) holds for $n < 0$ and $b_0 \neq 0$. If $\{b_n\}_{n=-\infty}^{\infty}$ is reflectionless on I , then $\{b_{-n}\}_{n=-\infty}^{\infty}$ is reflectionless on $\bar{I} = \{z \mid \bar{z} \in I\}$, so Theorem 4.2 implies $b_n \equiv 0$ if this right limit is reflectionless. But $b_0 \neq 0$. \square

5. Szegő's theorem

In this section, we'll prove

Theorem 5.1. *Let $\sum_{n=0}^{\infty} a_n z^n$ where the values of $\{a_n\}$ lie in a finite set, F . Then either $f(z) = \sum_{n=0}^{\infty} a_n z^n$ has a strong natural boundary or a_n is eventually periodic, in which case f is a rational function with poles at roots of unity.*

Remark. This result for ordinary natural boundaries is due to Szegő [47]. That f is unbounded on any sector is due to Duffin and Schaeffer [13], and that there is a strong natural boundary is a result of Boas [4]. This is an analogy of spectral theory results of Kotani [22] and Remling [36]. One could use an argument of Kotani and our reflectionless machinery to prove Theorem 5.2,

but instead we'll borrow part of Boas' argument and note one could use that to find an alternate proof of the spectral theory results.

Proof of Theorem 5.1. Let V be the finite set of possible values of a_n . Suppose $\{a_n\}$ is not eventually periodic. Fix $p = 1, 2, \dots$ and consider p blocks, $\{a_j\}_{j=\ell p+1}^{(\ell+1)p}$ for $\ell = 0, 1, 2, \dots$. Since there are only $(\#V)^p$ possible p blocks, some value must recur, that is, there exist $Q_p > P_p$ so that

$$a_{Q_p+j} = a_{P_p+j}, \quad j = 1, \dots, p. \tag{5.1}$$

If (5.1) holds for all $j \geq 1$, then for $k = P_p + j \geq P_p + 1$, we have

$$a_{(Q_p-P_p)+k} = a_k, \quad k = 1, \dots$$

that is, a is eventually periodic. Since we are assuming the contrary, there is $L_p \geq p + 1$, so $a_{Q_p+L_p} \neq a_{P_p+L_p}$.

Let $N_p = P_p + L_p \rightarrow \infty$ as $p \rightarrow \infty$ since $L_p \geq p$ and $M_p = Q_p + L_p > N_p + L$. Then

$$a_{N_p+j} = a_{M_p+j}, \quad j = -p, \dots, -1, \tag{5.2}$$

$$a_{N_p} \neq a_{M_p}. \tag{5.3}$$

By compactness, we get right limits, b, c , obeying (2.16), (2.17), so by Theorem 3.1, $\sum_{n=0}^\infty a_n z^n$ has a strong natural boundary. \square

We note the following extension of Szegő's theorem which appears in Bieberbach [3] (who only proved classical natural boundary):

Theorem 5.2. *Let $\{a_n\}_{n=0}^\infty$ be a bounded sequence with finitely many limit points. Then either $\sum_{n=0}^\infty a_n z^n$ has a strong natural boundary or there is a periodic sequence, c_n , with*

$$|a_n - c_n| \rightarrow 0 \quad \text{as } n \rightarrow \infty. \tag{5.4}$$

Proof. Let V be the finite set of limit points. Let $\gamma = \frac{1}{2} \min_{x,y \in V, x \neq y} |x - y|$. Eventually, for all n , there is $c_n \in V$ with $|a_n - c_n| \leq \gamma$. It follows that (5.4) holds. If c_n is eventually periodic, it can be modified for small n to be periodic and (5.4) still holds.

If c_n is not eventually periodic, by the above, it has a right limit which is not reflectionless. But a_n and c_n have the same right limits. \square

6. Random power series

In this section, we'll prove Theorem 1.7 as well as

Theorem 6.1. *Let $\{a_n(\omega)\}_{n=0}^\infty$ be a sequence of independent random variables so that*

- (i) $\sup_{n,\omega} |a_n(\omega)| = K < \infty$.

(ii) For some sequence $n_j \rightarrow \infty$,

$$\limsup_{j \rightarrow \infty} [\mathbb{E}(|a_{n_j}(\omega)|^2) - |\mathbb{E}(a_{n_j}(\omega))|^2] > 0. \tag{6.1}$$

Then for a.e. ω , $\sum_{n=0}^\infty a_n(\omega)z^n$ has a strong natural boundary.

Remark. \mathbb{E} is expectation. We'll use \mathbb{P} for probability and \mathbb{V} for variation, so (6.1) is $\limsup_{j \rightarrow \infty} \mathbb{V}(a_{n_j}(\omega)) > 0$.

Lemma 6.2.

(i) For any K and m , there exists $K_m > 0$ so that for any random variable f with $\|f\|_\infty \leq K$, there exists $z \in \mathbb{C}$ so that

$$\mathbb{P}\left(|f(\omega) - z| \leq \frac{1}{m}\right) \geq K_m. \tag{6.2}$$

(ii) For any K, m , and $\sigma > 0$, there exists $\tilde{K}_m > 0$ so that for any random variable f with $\|f\|_\infty \leq K$ and $\mathbb{V}(f) \geq \sigma$, there exist $z, w \in \mathbb{C}$ with

$$|z - w| \geq \left(\frac{\sigma}{2}\right)^{1/2} \tag{6.3}$$

so that

$$\mathbb{P}\left(|f(\omega) - z| \leq \frac{1}{m}\right) \geq \tilde{K}_m, \quad \mathbb{P}\left(|f(\omega) - w| \leq \frac{1}{m}\right) \geq \tilde{K}_m. \tag{6.4}$$

Proof. (i) Cover $\{z \mid |z| \leq K\}$ by disks of radius $\frac{1}{m}$. By compactness (or simple geometry), one can arrange for a finite number N_m . One of these disks must have probability at least $K_m = (N_m)^{-1}$. (6.2) holds for the center of that disk.

(ii) For any $c > 0$ and bounded random variable g ,

$$\mathbb{E}(|g|^2) \leq c^2 \mathbb{P}(|g| \leq c) + \|g\|_\infty^2 \mathbb{P}(|g| \geq c). \tag{6.5}$$

Thus, for any α with $|\alpha| \leq \|f\|_\infty$,

$$\text{Var}(f) \leq c^2 + (2\|f\|_\infty)^2 \mathbb{P}(|f - \alpha| > c). \tag{6.6}$$

Picking $c \leq (\frac{\sigma}{2})^{1/2}$ with $\sigma = \text{Var}(f)$, we see that

$$\mathbb{P}(|f - \alpha| > c) \geq \frac{1}{8\|f\|_\infty^2} \text{Var}(f). \tag{6.7}$$

Use (i) to find z so $\mathbb{P}(|f(\omega) - z| \leq \frac{1}{m}) \geq K_m$ where, if necessary, m is increased so $\frac{1}{m} \leq (\frac{\sigma}{2})^{1/2}$. By repeating (i) using $\mathbb{P}(|f - z| \geq (\frac{\sigma}{2})^{1/2}) \geq \frac{1}{8\|f\|_\infty} \sigma$, we get a w outside the disk $\{\zeta \mid |\zeta - z| \leq (\frac{\sigma}{2})^{1/2}\}$ so that $\mathbb{P}(|f(\omega) - w| \leq \frac{1}{m}) \geq \tilde{K}_m$ for some $\tilde{K}_m \leq K_m$. \square

Proof of Theorem 6.1. Pick $\sigma > 0$ and $n_j \rightarrow \infty$ so that $\mathbb{V}(a_{n_j}(\omega)) \geq \sigma$ for all j and so that $n_{j+1} \geq n_j$ and $n_{j+1} - n_j \rightarrow \infty$. For each $n \neq n_j$, for all j , use (i) of Lemma 6.2 to pick $z_n^{(m)}$ so that

$$\mathbb{P}\left(|a_n(\omega) - z_n^{(m)}| \leq \frac{1}{m}\right) \geq K_m. \tag{6.8}$$

By compactness and using the diagonalization trick, one can pass to a subsequence of the n_j , which we'll still denote by n_j , so for all m and all $k \neq 0$, $z_{n_j+k}^{(m)} \rightarrow z_k$ as $j \rightarrow \infty$.

By using (ii) of Lemma 6.2, find $w_j^{(m)}, \zeta_j^{(m)}$ so that

$$\begin{aligned} \mathbb{P}\left(|a_{n_j}(\omega) - w_j^{(m)}| \leq \frac{1}{m}\right) &\geq \tilde{K}_m, \\ \mathbb{P}\left(|a_{n_j}(\omega) - \zeta_j^{(m)}| \leq \frac{1}{m}\right) &\geq \tilde{K}_m \end{aligned} \tag{6.9}$$

and $|w_j^{(m)} - \zeta_j^{(m)}| \geq (\frac{\sigma}{2})^{1/2}$. Again, by passing to a subsequence, we may assume that $w_j^{(m)} \rightarrow w$, $\zeta_j^{(m)} \rightarrow \zeta$ where, of course, $|\zeta - w| \geq (\frac{\sigma}{2})^{1/2}$.

By the Borel–Cantelli lemma, for each m and each Q ,

$$|a_{n_j}(\omega) - w_j^{(m)}| \leq \frac{1}{m}, \quad |a_{n_j+k}(\omega) - z_{n_j+k}^{(m)}| \leq \frac{1}{m} \tag{6.10}$$

for all k with $0 < |k| \leq Q$, occurs infinitely often for a.e. ω , and the same for $\zeta_j^{(m)}$. Thus, for a.e. ω , the right limits include b, c with

$$b_k = c_k = z_k \quad \text{for } k \neq 0, \quad b_0 = w, \quad c_0 = \zeta \neq w. \tag{6.11}$$

It follows from Theorem 3.1 that $a_n(\omega)$ has a strong natural boundary for all ω . \square

Remark. The use of the Borel–Cantelli lemma to get nonreflectionless limits (in the context of spectral theory of CMV matrices) is taken from work of Breuer, Ryckman and Zinchenko [6].

For our proof of Theorem 1.7, we need two lemmas whose proof we defer to the end of the section.

Lemma 6.3. *Let $\alpha < \beta$ in $(-\pi, 2\pi)$ with $|\beta - \alpha| < 2\pi$. Then $\{\{a_n\}_{n=0}^\infty \mid \sup_n |a_n| \leq A \text{ for all } \varepsilon > 0; \sup_{0 < r < 1} (\int_{\alpha+\varepsilon}^{\beta-\varepsilon} |\sum_{n=0}^\infty a_n r^n e^{in\theta}| \frac{d\theta}{2\pi}) < \infty\}$ is a measurable set (in the product topology) invariant under $a_n \rightarrow a_{n+1}$.*

Lemma 6.4. *Let μ be a probability measure on $\{\{a_n\}_{n=-\infty}^\infty \mid \sup_n |a_n| \leq A\}$ that defines an ergodic invariant stochastic process. Let \mathcal{R}_μ be the set of two-sided series $\{b_n\}_{n=-\infty}^\infty$ so that $\{b_n\}_{n=-\infty}^\infty$ is a right limit of $\{a_n\}_{n=0}^\infty$ with positive probability. Then \mathcal{R}_μ is the support of the measure μ .*

Proof of Theorem 1.7. By Lemma 6.3, Theorem 1.4, and ergodicity (letting α, β run through rational multiples of 2π), either $\sum_{n=0}^\infty a_n z^n$ has a strong natural boundary with probability 1 or there is an interval, I , in $\partial\mathbb{D}$ so that with probability 1, all right limits of $\sum_{n=0}^\infty a_n z^n$ are reflectionless across I . In that case, by Lemma 6.4, all $\{b_n\}_{n=-\infty}^\infty$ in the support of μ are reflectionless across I . By Theorem 2.2, on $\text{supp}(\mu)$, b_0 is a continuous, and so measurable, function of $\{b_n\}_{n=-\infty}^{-1}$. Thus, the process is deterministic (in the strong sense of there being a continuous, rather than merely a measurable function). \square

Proof of Lemma 6.3. By Theorem 10.1 of Duren [14], the space E (sector) can be defined by a countable family of approximating curves. Thus, letting $z_1^{(n)} = \exp(i(\alpha + \frac{1}{n}))$, $z_2^{(n)} = \exp(i(\beta - \frac{1}{n}))$, the sup condition can be replaced by

$$\sup_{m,n} \left(\int_{\alpha + \frac{1}{n}}^{\beta - \frac{1}{n}} \left| (z - z_1^{(n)})(z - z_2^{(n)}) \sum_{n=0}^\infty a_n z^n \right| \Big|_{z=(1-\frac{1}{m})e^{i\theta}} \frac{d\theta}{2\pi} \right) < \infty. \tag{6.12}$$

Since it is described by a countable sup of uniformly convergent sums, this set is clearly measurable.

If $f(z) = \sum_{n=0}^\infty a_n z^n$ and $\tilde{f}(z) = \sum_{n=0}^\infty a_{n+1} z^n$, then

$$\tilde{f}(z) = z^{-1}(f(z) - a_0 z) \tag{6.13}$$

and finiteness of the sup in (6.12) implies finiteness of the sup for \tilde{f} replacing f . This proves the claimed invariance. \square

Proof of Lemma 6.4. If b_n is in $\text{supp}(d\mu)$, then for any $\varepsilon, N, \mu(\{a_n \mid |a_{N+n} - b_n| < \varepsilon, n = 0, \pm 1, \dots, \pm N\}) > 0$. By the ergodic theorem for a.e. ω ,

$$\lim_{M \rightarrow \infty} \frac{1}{M} \#\{j < M \mid |a_{n+j+N} - b_j| < \varepsilon, n = 0, \pm 1, \dots, \pm N\} > 0 \tag{6.14}$$

so for a.e. ω , there exists $N_\ell \rightarrow \infty$ with $|a_{n+N_\ell} - b_n| < \varepsilon$ for all n with $|n| < N$. By a diagonalization trick, b_n is a right limit for a.e. ω .

If b_n is not in the support of μ , pick N and ε so that $\mu(\{a_n \mid |a_{N+n} - b_n| \leq \varepsilon, |n| \leq N\}) = 0$. By translation invariance for all k ,

$$\mu(\{a_n \mid |a_{N+k+n} - b_n| \leq \varepsilon, n \leq N\}) = 0 \tag{6.15}$$

which implies b_n is not a right limit with probability 1. \square

7. Hecke’s example

Motivated by a spectral theory result of Damanik and Killip [12], we prove the following that includes Theorem 1.8 and so, Hecke’s example.

Theorem 7.1. *Let $T: \partial\mathbb{D} \rightarrow \partial\mathbb{D}$ be a homeomorphism so that for any $e^{i\theta} \in \partial\mathbb{D}$, $\{T^k(e^{i\theta}) \mid k = 0, 1, \dots\}$ is dense in $\partial\mathbb{D}$. Let $f: \partial\mathbb{D} \rightarrow \mathbb{C}$ be a bounded and piecewise continuous function with a finite number of discontinuities so that at, at least, one discontinuity, the right and left limits exist and are not equal. Then for any $e^{i\theta} \in \partial\mathbb{D}$, $\sum_{n=0}^{\infty} f(T^n(e^{i\theta}))z^n$ has a strong natural boundary on $\partial\mathbb{D}$.*

Remark. For other papers on other extensions of Hecke’s example, see [40,29,28,41,26,34,9].

Proof of Theorem 7.1. By rotating, we suppose $e^{i\theta} = 1$ is a point of a discontinuity with $\lim_{\theta \downarrow 0} f(e^{i\theta}) = r \neq s = \lim_{\theta \uparrow 0} f(e^{i\theta})$. By the density of any orbit for any $e^{i\theta}$, we can find $n_j \rightarrow \infty$, so $T^{n_j}(e^{i\theta}) \rightarrow 1$ with $T^{n_j}(e^{i\theta}) = e^{i\psi_j}$ with $-\pi < \psi_j < 0$ and $m_j \rightarrow \infty$, so $T^{m_j}(e^{i\theta}) \rightarrow 1$ and $T^{m_j}(e^{i\theta}) = e^{i\eta_j}$ with $\pi > \eta_j > 0$ (for find $n_{j+1} > n_j$ so $|T^{n_{j+1}}(e^{i\theta}) - e^{-i/(j+1)}| \leq (j+1)^{-2}$).

Thus, $a_{n_j} \rightarrow s$ and $a_{m_j} \rightarrow r$. On the other hand, look at the orbit $\{T^\ell(1) \mid \ell = 1, 2, \dots\}$. Since this orbit is dense, the $T^\ell(1)$ must be distinct, so for some L and all $\ell > L$, $T^\ell(1)$ must be a point of continuity. Thus, for $\ell > L$, $a_{n_j+\ell} - a_{m_j+\ell} \rightarrow 0$. The hypotheses of Theorem 2.3 hold for the right limits defined by a_{n_j+n} and a_{m_j+n} . So, by Theorem 3.1, we have a strong natural boundary. \square

8. Baire genericity

Proof of Theorem 1.9. Let $\{a_n\}_{n=0}^{\infty} \in \Omega^\infty$. Pick two distinct points, $z_0, z_1 \in \Omega$. For any ℓ , define

$$a_n^{(\ell)} = \begin{cases} a_n, & n \leq \ell, \\ z_0, & n = \ell + k! \text{ for } k = 1, 2, \dots, \\ z_1, & n = \ell + m, \ m \geq 1, \ m \neq k! \text{ for any } k. \end{cases}$$

Then, by the gap theorem, $\{a_n^{(\ell)}\}$ has a natural boundary on $\partial\mathbb{D}$, indeed, by Weierstrass’ original direct arguments. But $\lim_{\ell \rightarrow \infty} a_n^{(\ell)} = a_n$ for each fixed n , so the set in the theorem is dense in the weak topology.

For every rational multiple α, β of 2π with $\alpha < \beta$, and every $K = 1, 2, \dots$, and any $n = 1, \dots$, let $F_{\alpha,\beta,K,n} = \{\{a_n\}_{n=1}^{\infty} \in \Omega^\infty \mid f(z) = \sum_{n=0}^{\infty} a_n z^n \text{ has an analytic continuation to } \{z \mid |z| < 1 + n^{-1}, \alpha < \arg(z) < \beta\} \text{ with } |f(z)| \leq K \text{ there}\}$. It is easy to see that

- (i) Each $F_{\alpha,\beta,K,n}$ is closed in the topology of pointwise convergence of the a_n ’s (by the Vitali theorem).
- (ii) $\bigcup F_{\alpha,\beta,K,n}$ is the set of power series for which $\partial\mathbb{D}$ is not a natural boundary.

Thus, the complement of the set in the theorem is an F_δ , so the set is a G_δ . \square

Acknowledgments

We would like to thank Shmuel Agmon, John Garnett, Jean-Pierre Kahane, Rowan Killip, and Genadi Levin for useful discussions.

References

- [1] S. Agmon, Sur le séries de Dirichlet, *Ann. Sci. École Norm. Sup.* (3) 66 (1949) 263–310.
- [2] S. Agmon, Functions of exponential type in an angle and singularities of Taylor series, *Trans. Amer. Math. Soc.* 70 (1951) 492–508.
- [3] L. Bieberbach, Analytische Fortsetzung, *Ergeb. Math. Grenzgeb.*, vol. 3, Springer-Verlag, Berlin/Göttingen/Heidelberg, 1955.
- [4] R.P. Boas, *Entire Functions*, Academic Press, New York, 1954.
- [5] J. Breuer, E. Ryckman, B. Simon, Equality of the spectral and dynamical definitions of reflection, *Comm. Math. Phys.* 295 (2010) 531–550.
- [6] J. Breuer, E. Ryckman, M. Zinchenko, Right limits and reflectionless measures for CMV matrices, *Comm. Math. Phys.* 292 (2009) 1–28.
- [7] J. Brillhart, On the Rudin–Shapiro polynomials, *Duke Math. J.* 40 (1973) 335–353.
- [8] R. Carmona, J. Lacroix, *Spectral Theory of Random Schrödinger Operators*, *Probab. Appl.*, Birkhäuser, Boston, 1990.
- [9] F.W. Carroll, J.H.B. Kemperman, Noncontinuable analytic functions, *Duke Math. J.* 32 (1965) 65–83.
- [10] S.N. Chandler-Wilde, M. Lindner, Sufficiency of Favard’s condition for a class of band-dominated operators on the axis, *J. Funct. Anal.* 254 (2008) 1146–1159.
- [11] S.N. Chandler-Wilde, M. Lindner, Limit operators, collective compactness, and the spectral theory of infinite matrices, *Mem. Amer. Math. Soc.*, doi:10.1090/S0065-9266-2010-00626-4, in press.
- [12] D. Damanik, R. Killip, Ergodic potentials with a discontinuous sampling function are non-deterministic, *Math. Res. Lett.* 12 (2005) 187–192.
- [13] R.J. Duffin, A.C. Schaeffer, Power series with bounded coefficients, *Amer. J. Math.* 67 (1945) 141–154.
- [14] P.L. Duren, *Theory of H^p Spaces*, *Pure Appl. Math.*, vol. 38, Academic Press, New York/London, 1970.
- [15] P. Erdős, Note on the converse of Fabry’s gap theorem, *Trans. Amer. Math. Soc.* 57 (1945) 102–104.
- [16] G. Faber, Über Potenzreihen mit unendlich vielen verschwindenden Koeffizienten, *Sitz. ber. Konigl. Bayer. Akad. Wiss. Math.-Phys.* 36 (1906) 581–583.
- [17] E. Fabry, Sur les points singuliers d’une fonction donnée par son développement en série et l’impossibilité du prolongement analytique dans les cas très généraux, *Ann. Sci. École Norm. Sup.* (3) 13 (1896) 367–399.
- [18] V. Georgescu, A. Iftimovici, Crossed products of C^* -algebras and spectral analysis of quantum Hamiltonians, *Comm. Math. Phys.* 228 (2002) 519–560.
- [19] J. Hadamard, Essai sur l’étude des fonctions données par leur développement de Taylor, *J. Math. Pures Appl.* (4) 8 (1892) 101–186.
- [20] E. Hecke, Über analytische Funktionen und die Verteilung von Zahlen mod. eins, *Hamburg Abh.* 1 (1921) 54–76.
- [21] J.-P. Kahane, *Some Random Series of Functions*, second ed., *Cambridge Stud. Adv. Math.*, vol. 5, Cambridge University Press, Cambridge, 1985.
- [22] S. Kotani, Jacobi matrices with random potentials taking finitely many values, *Rev. Math. Phys.* 1 (1989) 129–133.
- [23] Y. Last, B. Simon, Eigenfunctions, transfer matrices, and absolutely continuous spectrum of one-dimensional Schrödinger operators, *Invent. Math.* 135 (1999) 329–367.
- [24] Y. Last, B. Simon, The essential spectrum of Schrödinger, Jacobi, and CMV operators, *J. Anal. Math.* 98 (2006) 183–220.
- [25] M. Măntoiu, C^* -algebras, dynamical systems at infinity and the essential spectrum of generalized Schrödinger operators, *J. Reine Angew. Math.* 550 (2002) 211–229.
- [26] H.G. Meijer, Irrational power series, *Indag. Math.* 25 (1963) 682–690.
- [27] S.A. Molchanov, Multiscale averaging for ordinary differential equations. Applications to the spectral theory of one-dimensional Schrödinger operator with sparse potentials, in: *Ser. Adv. Math. Appl. Sci.*, vol. 50, World Sci. Publ., River Edge, NJ, 1999, pp. 316–397.
- [28] L.J. Mordell, Irrational power series, *Proc. Amer. Math. Soc.* 12 (1961) 522–526.
- [29] M. Newman, Irrational power series, *Proc. Amer. Math. Soc.* 11 (1960) 699–702.

- [30] R.E.A.C. Paley, A. Zygmund, A note on analytic functions in the unit circle, *Proc. Cambridge Philos. Soc.* 28 (1932) 266–272.
- [31] L. Pastur, A. Figotin, *Spectra of Random and Almost-Periodic Operators*, Grundlehren Math. Wiss., vol. 297, Springer, Berlin, 1992.
- [32] D.B. Pearson, Singular continuous measures in scattering theory, *Comm. Math. Phys.* 60 (1978) 13–36.
- [33] G. Pólya, On converse gap theorems, *Trans. Amer. Math. Soc.* 52 (1942) 65–71.
- [34] J. Popken, Irrational power series, *Indag. Math.* 25 (1963) 691–694.
- [35] V.S. Rabinovich, Essential spectrum of perturbed pseudodifferential operators. Applications to the Schrödinger, Klein–Gordon, and Dirac operators, *Russ. J. Math. Phys.* 12 (2005) 62–80.
- [36] C. Remling, The absolutely continuous spectrum of Jacobi matrices, *Ann. of Math.*, in press.
- [37] R. Remmert, *Classical Topics in Complex Function Theory*, Grad. Texts in Math., vol. 172, Springer-Verlag, New York, 1998.
- [38] M. Riesz, Satze über Potenzreihen, *Ark. Mat. Astr. Fys.* 11 (1916) 1–16.
- [39] W. Rudin, Some theorems on Fourier coefficients, *Proc. Amer. Math. Soc.* 10 (1959) 855–859.
- [40] R. Salem, Power series with integral coefficients, *Duke Math. J.* 12 (1945) 153–172.
- [41] W. Schwarz, Irrationale Potenzreihen, *Arch. Math.* 13 (1962) 228–240.
- [42] H.S. Shapiro, Extremal problems for polynomials and power series, Master’s thesis, Massachusetts Institute of Technology, 1951.
- [43] B. Simon, Operators with singular continuous spectrum: I. General operators, *Ann. of Math.* 141 (1995) 131–145.
- [44] B. Simon, *Szegő’s Theorem and Its Descendants: Spectral Theory for L^2 Perturbations of Orthogonal Polynomials*, Princeton University Press, Princeton, NJ, 2011.
- [45] M. Sodin, P. Yuditskii, Almost periodic Jacobi matrices with homogeneous spectrum, infinite-dimensional Jacobi inversion, and Hardy spaces of character-automorphic functions, *J. Geom. Anal.* 7 (1997) 387–435.
- [46] H. Steinhaus, Über die Wahrscheinlichkeit dafür, daßder Konvergenzkreis einer Potenzreihe ihre natürliche Grenze ist, *Math. Z.* 31 (1930) 408–416.
- [47] G. Szegő, Tschebyscheffsche Polynome und nichtfortsetzbare Potenzreihen, *Math. Ann.* 87 (1922) 90–111.