

8

Extreme points and the Krein–Milman theorem

The next four chapters will focus on an important geometric aspect of compact sets, namely, the role of extreme points where:

Definition An *extreme point* of a convex set, A , is a point $x \in A$, with the property that if $x = \theta y + (1 - \theta)z$ with $y, z \in A$ and $\theta \in [0, 1]$, then $y = x$ and/or $z = x$. $\mathcal{E}(A)$ will denote the set of extreme points of A .

In other words, an extreme point is a point that is not an interior point of any line segment lying entirely in A . This chapter will prove a point is a limit of convex combinations of extreme points and the following chapters will refine this representation of a general point.

Example 8.1 The ν -simplex, Δ_ν , is given by (5.3) as the convex hull in $\mathbb{R}^{\nu+1}$ of $\{\delta_1, \dots, \delta_{\nu+1}\}$, the coordinate vectors. It is easy to see its extreme points are precisely the $\nu + 1$ points $\{\delta_j\}_{j=1}^{\nu+1}$. The hypercube $C_0 = \{x \in \mathbb{R}^\nu \mid |x_i| \leq 1\}$ has the 2^ν points $(\pm 1, \pm 1, \dots, \pm 1)$ as extreme points. The ball $B^\nu = \{x \in \mathbb{R}^\nu \mid |x| \leq 1\}$ has the entire sphere as extreme points, showing $\mathcal{E}(A)$ can be infinite.

An interesting example (see Figure 8.1) is the set $A \subset \mathbb{R}^3$, which is the convex hull of

$$A = \text{ch}(\{(x, y, 0) \mid x^2 + y^2 = 1\} \cup \{(1, 0, \pm 1)\}) \quad (8.1)$$

Its extreme points are

$$\mathcal{E}(A) = \{(x, y, 0) \mid x^2 + y^2 = 1, x \neq 1\} \cup \{(1, 0, \pm 1)\}$$

$(1, 0, 0) = \frac{1}{2}(1, 0, 1) + \frac{1}{2}(1, 0, -1)$ is not an extreme point. This example shows that even in the finite-dimensional case, the extreme points may not be closed. In the infinite-dimensional case, we will even see that the set of extreme points can be dense! □




Figure 8.1 An example of not closed extreme points

If a point, x , in A is not extreme, it is an interior point of some segment

$$[y, z] = \{\theta y + (1 - \theta)z \mid 0 \leq \theta \leq 1\} \tag{8.2}$$

with $y \neq z$. If y or z is not an extreme point, we can write them as convex combinations and continue. (If A is compact and in \mathbb{R}^ν , and if one extends the line segment to be maximal, one can prove this process will stop in finitely many steps. Indeed, that in essence is the method of proof we will use in Theorem 8.11.) If one thinks about writing y, z as convex combinations, one “expects” that any point in A is a convex linear combination of extreme points of A – and we will prove this when A is compact and finite-dimensional. Indeed, if $A \subset \mathbb{R}^\nu$, we will prove that at most $\nu + 1$ extreme points are needed. This fails in infinite dimension, but we will find a replacement, the Krein–Milman theorem, which says that any point is a limit of convex combinations of extreme points. These are the two main results of this chapter.

Extreme points are a special case of a more general notion:

Definition A *face* of a convex set is a nonempty subset, F , of A with the property that if $x, y \in A$, $\theta \in (0, 1)$, and $\theta x + (1 - \theta)y \in F$, then $x, y \in F$. A face, F , that is strictly smaller than A is called a *proper face*.

Thus, a face is a subset so that any line segment $[xz] \subset A$, with interior points in F must lie in F . Extreme points are precisely one-point faces of A . (Note: See the remark before Proposition 8.6 for a later restriction of this definition.)

Example 8.2 (Example 8.1 continued) Δ_ν has lots of faces; explicitly, it has $2^{\nu+1} - 2$ proper faces, namely, $\nu + 1$ extreme points, $\binom{\nu+1}{2}$ facial lines, \dots , $\binom{\nu+1}{\nu-1}$ faces of dimension $(\nu - 1)$. The hypercube C_ν has $3^\nu - 1$ faces, namely, 2^ν extreme points, $\nu 2^{\nu-1}$ facial lines, $\binom{\nu}{2} 2^{\nu-2}$ facial planes, \dots , $2 \binom{\nu}{\nu-1}$ faces of dimension $(\nu - 1)$. The only faces on the ball are its extreme points. The faces of the set A of (8.1) are its extreme points, the line $\{(1, 0, y) \mid |y| \leq 1\}$, and the lines

$\{\theta(x_0, y_0, 0) + (1-\theta)(1, 0, 1)\}$ and $\{\theta(x_0, y_0, 0) + (1-\theta)(1, 0, -1)\}$, where x_0, y_0 are fixed with $x_0^2 + y_0^2 = 1$ and $x_0 \neq 1$. \square

A canonical way proper faces are constructed is via linear functionals.

Theorem 8.3 *Let A be a convex subset of a real vector space. Let $\ell: A \rightarrow \mathbb{R}$ be a linear functional with*

(i)

$$\sup_{x \in A} \ell(x) = \alpha < \infty \quad (8.3)$$

(ii) $\ell \upharpoonright A$ is not constant.

Then

$$\{y \mid \ell(y) = \alpha\} = F \quad (8.4)$$

if nonempty, is a proper face of A .

Remark If A is compact and ℓ is continuous, of course, F is nonempty.

Proof Since ℓ is linear, F is convex. Moreover, if $y, z \in A$ and $\theta \in (0, 1)$ and $\theta y + (1-\theta)z \in F$, then $\theta\ell(y) + (1-\theta)\ell(z) = \alpha$ and $\ell(y) \leq \alpha, \ell(z) \leq \alpha$ implies $\ell(y) = \ell(z) = \alpha$, that is, $y, z \in F$. By (ii), F is a proper subset of A . \square

The hyperplane $\{y \mid \ell(y) = \alpha\}$ with α given by (8.3) is called a *tangent hyperplane* or *support hyperplane*. The set (8.4) is called an *exposed set*. If F is a single point, we call the point an *exposed point*.

Example 8.4 We have just seen that every exposed set is a face so, in particular, every exposed point is an extreme point. I'll bet if you think through a few simple examples like a disk or triangle in the plane or a convex polyhedron in \mathbb{R}^3 , you'll conjecture the converse is true. But it is not! Here is a counterexample in \mathbb{R}^2 (see Figure 8.2):

$$A = \{(x, y) \mid -1 \leq x \leq 1, -2 \leq y \leq 0\} \cup \{(x, y) \mid x^2 + y^2 \leq 1\}$$

The boundary of A above $y = -2$ is a C^1 curve, so there is a unique supporting hyperplane through each such boundary point. The supporting hyperplane through the extreme point $(1, 0)$ is $x = 1$ so $(1, 0)$ is not an exposed point, but it is an extreme point. \square

Proposition 8.5 *Any proper face F of A lies in the topological boundary of A . Conversely, if $A \subset X$, a locally convex space (and, in particular, in \mathbb{R}^ν), and A^{int} is nonempty, then any point $x \in A \cap \partial A$ lies in a proper face.*




Figure 8.2 A nonexposed extreme point

Proof Let $x \in F$ and pick $y \in A \setminus F$. The set of $\theta \in \mathbb{R}$ so $z(\theta) \equiv \theta x + (1 - \theta)y \in A$ includes $[0, 1]$, but it cannot include any $\theta > 1$ for if it did, $\theta = 1$ (i.e., x) would be an interior point of a line in A with at least one endpoint in $A \setminus F$. Thus, $x = \lim_{n \downarrow 0} z(1 + n^{-1})$ is a limit point of points not in A , that is, $x \in \bar{A} \cap \overline{X \setminus A} = \partial A$.

For the converse, let $x \in A \cap \partial A$ and let $B = A^{\text{int}}$. Since B is open, Theorem 4.1 implies there exists a continuous $L \neq 0$ with $\alpha = \sup_{y \in B} L(y) \leq L(x)$. Since $x \in A$, $L(x) = \alpha$. Since B is open, $L[B]$ is an open set (Lemma 4.2), so the supporting hyperplane $H = \{y \mid L(y) = \alpha\}$ is disjoint from B and so $H \cap A$ is a proper face. □

To have lots of extreme points, we will need lots of boundary points, so it is natural to restrict ourselves to closed convex sets. The convex set $\mathbb{R}_+^n = \{x \in \mathbb{R}^n \mid x_i \geq 0 \text{ all } i\}$ has a single extreme point, so we will also restrict to bounded sets. Indeed, except for some examples, we will restrict ourselves to compact convex sets in the infinite-dimensional case. Convex cones are interesting but can normally be treated as suspensions of compact convex sets; see the discussion in Chapter 11. So we will suppose A is a compact convex subset of a locally convex space. As noted in Corollary 4.9, A is weakly compact, so we will suppose henceforth that we are dealing with the weak topology.

Remark Henceforth, we will also restrict the term “face” to indicate a closed set.

Proposition 8.6 *Let $F \subset A$ with A a compact convex set and F a face of A . Let $B \subset F$. Then B is a face of F if and only if it is a face of A . In particular, $x \in F$ is in $\mathcal{E}(F)$ if and only if it is also in $\mathcal{E}(A)$, that is,*

$$\mathcal{E}(F) = F \cap \mathcal{E}(A)$$

Proof If B is a face of A , $x \in B$, and x is an interior point of $[y, z] \subset F$, it is an interior point of $[y, z] \subset A$. Thus, $y, z \in A$, so $y, z \in B$, and thus, B is a face of F .

Conversely, if B is a face of F , $x \in B$, and $x \in [y, z] \subset A$, since $x \in F$, the fact that F is a face implies $y, z \in F$ so $[y, z] \subset F$. Thus, since B is a face of F , $y, z \in B$ and so B is a face of A . □

We turn next to a detailed study of the finite-dimensional case. We begin with some notions that involve finite dimension but which are useful in the infinite-dimensional case also. Since we will be discussing affine subspaces, affine spaces, affine independence, etc., we will temporarily use vector subspaces, etc. to denote the usual notions in a vector space where we don't normally include "vector."

Let X be a vector space. An *affine subspace* is a set of the form $a + W$ where $a \in X$ and W is a vector subspace. The *affine span* of a subset $A \subset X$ is the smallest affine subspace containing A . If $A = \{e_1, \dots, e_n\}$, then its affine span is just

$$S(e_1, \dots, e_n) = \left\{ \theta_1 e_1 + \dots + \theta_n e_n \mid \theta \in \mathbb{R}^n, \sum_{i=1}^n \theta_i = 1 \right\} \quad (8.5)$$

as is easy to see since $\sum_{i=1}^n \theta_i = 1$ implies that

$$\theta_1 e_1 + \dots + \theta_n e_n = e_1 + \sum_{j=2}^n \theta_j (e_j - e_1) \quad (8.6)$$

so the right-hand side of (8.5) is e_1 plus the vector span of $\{e_j - e_1\}_{j=2}^n$. The convex hull of $\{e_1, \dots, e_n\}$ is, of course,

$$\text{ch}(e_1, \dots, e_n) = \left\{ \theta_1 e_1 + \dots + \theta_n e_n \mid \theta \in \mathbb{R}^n, \sum_{i=1}^n \theta_i = 1, \theta_i \geq 0 \right\} \quad (8.7)$$

We call $\{e_1, \dots, e_n\}$ *affinely independent* if and only if $\sum_{i=1}^n \theta_i e_i = 0$ and $\sum_{i=1}^n \theta_i = 0$ implies $\theta \equiv 0$. By (8.6) this is true if and only if $\{e_j - e_1\}_{j=2}^n$ are vector independent.

Proposition 8.7 $\text{ch}(e_1, \dots, e_n)$ always has a nonempty interior as a subset of $S(e_1, \dots, e_n)$.

Proof By successively throwing out dependent vectors from $P = \{e_j - e_1\}_{j=2}^n$, find a maximal independent subset of P . By relabeling, suppose it is $P' \equiv \{e_j - e_1\}_{j=1}^k$ so $\{e_1, \dots, e_k\}$ are affinely independent, and each $e_\ell - e_1$ with $\ell > k$ is a linear combination of P' . Then $S(e_1, \dots, e_n) = S(e_1, \dots, e_k)$.

Since $\text{ch}(e_1, \dots, e_k) \subset \text{ch}(e_1, \dots, e_n)$, it suffices to prove the result when e_1, \dots, e_n are affinely independent. In that case, $\varphi: \Delta_{n-1} \rightarrow \text{ch}(e_1, \dots, e_k)$ is a bijection and continuous, so a homeomorphism. Since Δ_{n-1} has a nonempty interior ($\{(\theta_1, \dots, \theta_n) \mid \sum_{i=1}^n \theta_i = 1, 0 < \theta_i\}$), so does $\text{ch}(e_1, \dots, e_k)$. \square

Remark The θ 's are called *barycentric coordinates* for $S(e_1, \dots, e_\ell)$ and $\text{ch}(e_1, \dots, e_\ell)$.

Theorem 8.8 Let $A \subset \mathbb{R}^v$ be a convex set. Then there is a unique affine subspace W of \mathbb{R}^v so that $A \subset W$, and as a subset of W , A has a nonempty interior.

Proof Pick $e_1 \in A$ and consider $B = A - e_1 \ni 0$. Let W be the subspace generated by B , that is, let $f_1, \dots, f_{\ell-1}$ be a maximal linear independent subset of B , and let X be the vector span of $\{f_j\}_{j=1}^{\ell-1}$. Let $e_j = f_{j-1} + e_1$ for $j = 2, \dots, \ell$ so $e_1 + X \equiv W$ is the affine span of $\{e_j\}_{j=1}^{\ell}$. By construction $B \subset X$ so $A \subset W = S(e_1, \dots, e_{\ell})$. By Proposition 8.7, $\text{ch}(e_1, \dots, e_{\ell}) \subset A$ is open in S , so A has no nonempty interior as a subset of S .

W is unique because any affine subspace containing A must contain e_1, \dots, e_{ℓ} and so $S(e_1, \dots, e_{\ell})$. If its dimension were larger than W , W would have empty interior in it and so would A . Thus, the condition that A have nonempty interior uniquely determines W . \square

Definition The *dimension* of a convex set $A \subset \mathbb{R}^{\nu}$ is the dimension of the unique affine subspace given by Theorem 8.8. The interior of A as a subset of W is written A^{int} and called the *intrinsic interior* of A . $\partial^i A$, the *intrinsic boundary* of $A = \bar{A} \setminus A^{\text{int}}$.

Proposition 8.9 *Let A be a compact convex subset of \mathbb{R}^{ν} . Then*

- (i) $\partial^i A$ is the union of all the proper faces of A .
- (ii) If $x \in \partial^i A$ and y is any point in A^{int} , $\{\theta \mid (1 - \theta)x + \theta y \in A\} = [0, \alpha]$ for some $\alpha > 1$.
- (iii) If $x \in A^{\text{int}}$ and $y \in A$, $\{\theta \mid (1 - \theta)x + \theta y \in A\} \cap (-\infty, 0) \neq \emptyset$.

Remark This gives us an intrinsic definition of A^{int} . $x \in A^{\text{int}}$ if and only if for any $y \in A$, the line $[y, x]$ continued past x lies in A for at least a while. Similarly, $\partial^i A$ is determined by the condition that any line that intersects A in more than one point enters and leaves A at points in $\partial^i A$ and any $x \in \partial^i A$ lies on such a line as an endpoint.

Proof (i) This follows from Proposition 8.5 if we view A as a subset of W .

(ii) We know $x \in \partial^i A$ lies in some face F . Since A^{int} , viewed as a subset of W , is disjoint from the boundary, $y \notin F$. As in the proof of Proposition 8.5, $\{\theta \mid (1 - \theta)x + \theta y \in A\} \cap (-\infty, 0) = \emptyset$. Since this set is connected and compact and contains $[0, 1]$, it must be the requisite form. That $\alpha > 1$ and $\alpha \neq 1$ follow from (iii).

(iii) $[x, y]$ lies in A , so in W , so since A^{int} is open in W , $\{\theta \mid (1 - \theta)x + \theta y \in A^{\text{int}}\}$ is open. Since it contains 0, it must contain an interval $(-\varepsilon, \varepsilon)$ about 0. \square

Proposition 8.10 *Let $A \subset \mathbb{R}^{\nu}$ be a compact convex set. Let $\ell = \dim(A)$ and let F be a proper face of F . Then $\dim(F) < \ell$.*

Proof Let $A \subset W$ where W is the unique ℓ -dimensional space containing A . If $\dim(F) = \ell$, then W must also be the unique ℓ -dimensional space containing F , and so F has not empty interior. But as a set in W , $F \subset \partial A$, which contradicts $F^{\text{int}} \neq \emptyset$. Thus, $\dim(F) < \ell$. \square

We are now ready for the main finite-dimensional result:

Theorem 8.11 (Minkowski–Carathéodory Theorem) *Let A be a compact convex subset of \mathbb{R}^ν of dimension n . Then any point in A is a convex combination of at most $n + 1$ extreme points. In fact, for any x , one can fix $e_0 \in \mathcal{E}(A)$ and find $e_1, \dots, e_n \in \mathcal{E}(A)$ so x is a convex combination of $\{e_j\}_{j=0}^n$. If $x \in A^{\text{int}}$, then $x = \sum_{j=0}^n \theta_j e_j$ with $\theta_0 > 0$. In particular,*

$$A = \text{ch}(\mathcal{E}(A)) \quad (8.8)$$

Remarks 1. It pays to think of the square in \mathbb{R}^2 which has four extreme points, but where any point is in the convex hull of three points (indeed, for most interior points in exactly two ways).

2. The example of the n simplex Δ_n shows that for general A 's, one cannot do better than $n + 1$ points. Of course, for some sets, one can do better. No matter what value of ν , the ball B^ν has the property that any point is a convex combination of at most two extreme points.

Proof We use induction on n . $n = 0$, that is, single-point sets, is trivial. Suppose we have the result for all sets, B , with $\dim(B) \leq n - 1$. Let A have dimension n and $x \in A$ and $e_0 \in \mathcal{E}(A)$. Take the line segment $[e_0, x]$ and extend it – $\{\theta \mid (1 - \theta)e_0 + \theta x \in A\} = [0, \alpha]$ for some α by Proposition 8.9. Let $y = (1 - \alpha)e_0 + \alpha x$. Since $\alpha \geq 1$,

$$x = \theta_0 e_0 + (1 - \theta_0)y \quad (8.9)$$

where $\theta_0 = 1 - \alpha^{-1} \geq 0$.

By construction, $y \in \partial^i A$ and so, by Proposition 8.9, $y \in F$, some proper face of A . By Proposition 8.10, $\dim(F) \leq n - 1$, so by the induction hypothesis, $y = \sum_{j=1}^n \varphi_j e_j$ where $\varphi_j \geq 0$, $\sum_{j=1}^n \varphi_j = 1$, and $\{e_1, \dots, e_n\} \subset \mathcal{E}(F)$. By Proposition 8.6, $\mathcal{E}(F) \subset \mathcal{E}(A)$. Thus,

$$x = \sum_{j=0}^n \theta_j e_j$$

where $\theta_j = (1 - \theta_0)\varphi_j$ for $j = 1, \dots, n$.

If $\theta_0 = 0$, by (8.9), $x = y$ and $x \in \partial^i A$. Thus, if $x \in A^{\text{int}}$, $\theta_0 \neq 0$. □

We will have more to say about extreme points of finite-dimensional convex sets in Chapter 15 when we discuss a particular convex set, the set of all doubly stochastic matrices. In particular, we will show that a compact, convex set, K , in \mathbb{R}^ν has finitely many extreme points if and only if it is a finite intersection of closed half-spaces (Corollary 15.3).

In the infinite-dimensional case, it is not clear that $\mathcal{E}(A)$ is nonempty – we will go through the main construction in two phases. We will first show that $\mathcal{E}(A) \neq \emptyset$ for A a compact convex subset of a locally convex space and then, fairly easily, we will be able to show that

$$A = \text{cch}(\mathcal{E}(A))$$

which is the Krein–Milman theorem. The following illustrates that the infinite-dimensional case is subtle.

Example 8.12 Let A be the closed unit ball in $L^1(0, 1)$. Let $f \in A$ with $f \neq 0$. Then $H_f(s) = \int_0^s |f(t)| dt$ is a continuous function with $H_f(0) = 0$ and $H_f(1) = \alpha \leq 1$. Thus, there exists s_0 with $H_f(s_0) = \alpha/2$. Let

$$g = 2f\chi_{(0, s_0)}$$

$$h = 2f\chi_{(s_0, 1)}$$

Then $\|g\|_1 = \|h\|_1 = \|f\|_1 = \alpha \leq 1$ and $f = \frac{1}{2}h + \frac{1}{2}g$. Since $h \neq g$, f is not an extreme point. Clearly, $0 = \frac{1}{2}(f - f)$ is not extreme either. Thus, A has no extreme points!

We will show below that any compact convex subset, A , of a locally convex space has $\mathcal{E}(A) \neq \emptyset$. This means that the unit ball in $L^1(0, 1)$ cannot be compact in any topology making it into a locally convex space. In particular, because of the Bourbaki–Alaoglu theorem, $L^1(0, 1)$ cannot be the dual of any Banach space. This is subtle because $\ell^1(\mathbb{Z})$ is a dual (of $c_0(\mathbb{Z})$, the bounded sequences vanishing at infinity). Of course, the unit ball in $\ell^1(\mathbb{Z})$ has lots of extreme points in each $\pm\delta_n$. □

Proposition 8.13 *Let A be a compact convex subset of a locally convex space, X . Then $\mathcal{E}(A) \neq \emptyset$.*

Proof Extreme points are one-point faces. We will find them as minimal faces. So let \mathcal{F} be the family of proper faces of A with $F_1 > F_2$ if $F_1 \subset F_2$. This is a partially ordered set and it has the chain property, that is, if $\{F_\alpha\}_{\alpha \in I}$ is linearly ordered, then it has an “upper” bound (“upper” here means small since a “larger than” means contained in), namely, $\bigcap_{\alpha \in I} F_\alpha$. This is closed, a face (by a simple argument), and nonempty because of the intersection property for compact sets.

Thus, by Zorn’s lemma, there exist minimal faces. Suppose F is such a minimal face and F has at least two distinct points x and y . By Corollary 4.6, there is a linear functional on X and so on F with $\ell(x) \neq \ell(y)$. Since F is compact,

$$\tilde{F} = \{z \in F \mid \ell(z) = \sup_{w \in F} \ell(w)\}$$

is nonempty. It is a face of F and so, by Proposition 8.6, \tilde{F} is a face of A . Since $\ell(x) \neq \ell(y)$, it cannot be that both x and y lie in \tilde{F} , so $\tilde{F} \subsetneq F$, violating minimality. It follows that F has a single point and that point must be an extreme point. \square

Remark In $L^1(0, 1)$, $F_\alpha = \{f \in L^1 \mid \|f\|_1 = 1, f \geq 0, \text{ and } f(x) = 0 \text{ on } (0, \alpha)\}$ is a face and it is linearly ordered (since $\alpha > \beta \Rightarrow F_\alpha \subset F_\beta$), but $\bigcap_\alpha F_\alpha$ is empty. This proves the lack of compactness directly.

Theorem 8.14 (The Krein–Milman Theorem) *Let A be a compact convex subset of a locally convex vector space, X . Then*

$$A = \text{cch}(\mathcal{E}(A)) \quad (8.10)$$

Proof Since $\mathcal{E}(A) \subset A$ and A is closed and convex, $B \equiv \text{cch}(\mathcal{E}(A)) \subset A$. Suppose $B \neq A$ so there exists $x_0 \in A \setminus B$. Since B is closed and convex, by Theorem 4.5, there exists $\ell \in X^*$ so

$$\ell(x_0) > \sup_{y \in B} \ell(y) \quad (8.11)$$

Let $F = \{x \in A \mid \ell(x) = \sup_{z \in A} \ell(z)\}$. Then F is nonempty since A is compact, a face, and by (8.11),

$$F \cap B = \emptyset \quad (8.12)$$

By Proposition 8.13, F has an extreme point, y_0 , and then, by Proposition 8.6, $y_0 \in \mathcal{E}(A)$. Thus, $y_0 \in B$, contradicting (8.12). \square

Remark In the next chapter (see Theorem 9.4), we will prove a sort of converse of this theorem.

Example 8.15 Let $X = C_{\mathbb{R}}([0, 1])$ and let A be the unit ball in $\|\cdot\|_\infty$. If $|f(x)| < 1$ for some x_0 in $[0, 1]$, then by continuity for some ε , $|f(y)| < 1$ for $|y - x_0| < \varepsilon$ and we can find $g \neq 0$ supported in $(x_0 - \varepsilon, x_0 + \varepsilon)$, so both $f + g$ and $f - g$ lie in A . Since $f = \frac{1}{2}(f + g) + \frac{1}{2}(f - g)$, f is not an extreme point. Thus, extreme points have $|f(x)| = 1$. By continuity and reality, A has precisely two extreme points $f \equiv \pm 1$. $\text{cch}(\mathcal{E}(A))$ is the constant functions in A so $A \neq \text{cch}(\mathcal{E}(A))$. Thus, $C_{\mathbb{R}}([0, 1])$ is not a dual space. \square

Example 8.16 This is an important example. Let X be a compact Hausdorff space and let $A = \mathcal{M}_{+1}(X)$ be the set of regular Borel probability measures on X . The extreme points of A are precisely the single-point pure points, δ_x , since if $C \subset X$ has $0 < \mu(C) < 1$ and

$$\begin{aligned} \mu_C(B) &= \mu(C)^{-1} \mu(B \cap C) \\ \mu_{X \setminus C} &= \mu(X \setminus C)^{-1} \mu(B \setminus C) \end{aligned}$$

then with $\theta = \mu(C)$, $\mu = \theta\mu_C + (1 - \theta)\mu_{X \setminus C}$ so μ is not an extreme point.

Suppose μ has the property that $\mu(A)$ is 0 or 1 for each $A \subset X$. If $x \neq y$ are both in $\text{supp}(\mu)$, we can find disjoint open sets B, C with $x \in B$ and $y \in C$. By the 0, 1 law, either $\mu(B) = 0$ or $\mu(C) = 0$ or both. But that would mean x and y cannot both be in $\text{supp}(\mu)$. Thus, $\text{supp}(\mu)$ is a single point and $\mu = \delta_x$ for some x , that is, the only extreme points are among the $\{\delta_x\}$. But each δ_x is an extreme point since $\delta_x = \frac{1}{2}\mu + \frac{1}{2}\nu$ implies $\text{supp}(\mu) \subset \{x\}$ so $\mu = \delta_x$. Thus, $\mathcal{E}(A) = \{\delta_x \mid x \in X\}$.

$\text{ch}(\mathcal{E}(A))$ is the pure point measures. A is compact in the $\sigma(\mathcal{M}(X), C(X))$ -topology and so the Krein–Milman theorem says that the pure point measures are weakly dense – something that is easy to prove directly. \square

Example 8.17 In some ways, this is an extension of the last example. Let X be a compact Hausdorff space and let $T: X \rightarrow X$ be a continuous bijection. A regular Borel probability measure μ on X is called *invariant* if and only if $\mu(T^{-1}[A]) = \mu(A)$ for all $A \subset X$. This is equivalent to

$$\int f(Tx) d\mu(x) = \int f(x) d\mu(x) \tag{8.13}$$

for all $f \in C(X)$. An invariant measure, μ , is called *ergodic* if and only if $\mu(A \Delta T[A]) = 0$ (i.e., $A = T[A]$ μ a.e.) implies $\mu(A)$ is 0 or 1.

Let T^* map $\mathcal{M}_{+,1}(X) \rightarrow \mathcal{M}_{+,1}(X)$ by

$$\int f(x) d(T^*\mu)(x) = \int f(Tx) d\mu(x)$$

Pick any $\mu \in \mathcal{M}_{+,1}(X)$ and let

$$\mu_n = \frac{1}{n} \sum_{j=0}^{n-1} (T^*)^j(\mu)$$

Then for any $f \in C(X)$,

$$|\mu_n(f) - \mu_n(Tf)| = \left| \frac{1}{n} [((T^*)^n \mu)(f) - \mu(f)] \right| \leq \frac{2}{n} \|f\|_\infty \tag{8.14}$$

Thus, if μ_∞ is any weak-* limit point of μ_n , $\mu_\infty(Tf) = \mu_\infty(f)$ for all f , that is, $T^*\mu_\infty = \mu_\infty$. Since $\mathcal{M}_{+,1}(X)$ is compact in the weak-* topology, we conclude

$$\mathcal{M}_{+,1}^I(T) = \{\mu \in \mathcal{M}_{+,1} \mid T^*\mu = \mu\}$$

is not empty.

We claim $\mu \in \mathcal{M}_{+,1}^I(T)$ is ergodic if and only if $\mu \in \mathcal{E}(\mathcal{M}_{+,1}^I(T))$. Suppose μ is not ergodic. Then there exists an almost invariant set A with $0 < \mu(A) < 1$. μ can be decomposed $\mu = \theta\mu_A + (1 - \theta)\mu_{X \setminus A}$ with $\theta = \mu(A)$ and $\mu_C(B) = \mu(C)^{-1}\mu(B \cap C)$.

Conversely, suppose μ is ergodic. Then in $L^2(X, d\mu)$, define $(Uf)(x) = f(Tx)$. Then U is unitary. Since as functions on ∂D ,

$$\frac{1}{n} \sum_{j=0}^{n-1} e^{inj\theta} \rightarrow \begin{cases} 1, & \theta = 0 \\ 0, & \theta \in (0, 2\pi) \end{cases}$$

the continuity of the functional calculus (see [303, Thm. VIII.20]) implies

$$\frac{1}{n} \sum_{j=1}^{n-1} U^j f \xrightarrow{L^2} P_{\{1\}} f \quad (8.15)$$

where $P_{\{1\}}$ is the projection onto the invariant functions, that is, those g with $Ug = g$. (This is essentially a version of the von Neumann ergodic theorem.) We claim that, since μ is ergodic, any such g is constant. For clearly, $\operatorname{Re} g$ and $\operatorname{Im} g$ obey $Ug = g$ so we can suppose g is real. But then, for all rational (α, β) , $\{x \mid \alpha < g(x) < \beta\}$ is almost T -invariant and so it has measure 0 or 1. This implies g is a.e. constant. Since $\langle 1, U^n f \rangle = \langle 1, f \rangle = \mu(f)$, we see the constant must be $\mu(f) = \int f(x) d\mu(x)$.

We have thus shown that if μ is ergodic, then

$$\int \left| \frac{1}{n} \sum_{j=0}^{n-1} f(T^{n-1}x) - \mu(f) \right|^2 d\mu(x) = 0 \quad (8.16)$$

Suppose now $\mu = \theta\nu + (1 - \theta)\eta$ with $0 < \theta < 1$. Since (8.16) has a positive integrand, we see that (8.16) holds if $d\mu$ is replaced by $d\nu$ or $d\eta$ (but $\mu(f)$ is left unchanged). Thus,

$$\int \frac{1}{n} \sum_{j=0}^{n-1} f(T^{n-1}x) d\nu(x) \rightarrow \mu(f) \quad (8.17)$$

But since ν is invariant, the left side of (8.17) is $\nu(f)$ for any n . Thus, $\nu(f) = \mu(f)$, and similarly, $\eta(f) = \mu(f)$. It follows that $\nu = \eta = \mu$, that is, μ is an extreme point.

We have therefore shown that ergodic measures are precisely the extreme points of $\mathcal{M}_{+,1}^I(T)$. The Krein–Milman theorem therefore implies the existence of ergodic measures. If $\mathcal{M}_{+,1}^I(T)$ has more than one point, there must be multiple extreme points.

Now suppose that $\{T_\alpha\}_{\alpha \in I}$ is an arbitrary family of commuting maps of X to X . Invariant measures for all the T_α 's at once are defined in the obvious way, and μ is called ergodic if $\mu(A \Delta T_\alpha[A]) = 0$ for all α implies $\mu(A)$ is 0 or 1. Since the T 's commute, T_α^* maps each $\mathcal{M}_{+,1}^I(T_\beta)$ to itself, and so by repeating the proof that $\mathcal{M}_{+,1}(X)$ has invariant measures, we see $\mathcal{M}_{+,1}^I(T_\beta)$ has a T_α^* -invariant point. By induction, there are invariant measures for any finite set $\{T_{\alpha_i}^*\}_{i=1}^\ell$, and then by compactness and the fact that invariant measures are closed, invariant measures for

all $\{T_\alpha\}_{\alpha \in I}$. We summarize in the following theorem. This example is discussed further in Example 9.7. \square

Theorem 8.18 *Let X be a compact Hausdorff space and let $\{T_\alpha\}_{\alpha \in I}$ be a family of commuting bijections of X to itself. Then $\mathcal{M}_{+,1}^I(\{T_\alpha\})$, the set of common invariant measures, is nonempty. The ergodic measures are precisely $\mathcal{E}(\mathcal{M}_{+,1}^I(\{T_\alpha\}))$, the extreme points, and are therefore also nonempty.*

As an example, if X is a compact abelian group, and for each $x \in X$, $T_x : X \rightarrow X$ by $T_x(y) = xy$, then there is an invariant measure. We have therefore constructed a Haar measure in this case, which is known to be unique. Similar ideas can be used to construct what are invariant means on noncompact abelian groups. See the Notes.

(8.15) provides a useful criterion for ergodicity.

Theorem 8.19 *Let μ be an invariant measure for a continuous bijection T on a compact Hausdorff space. For any function $f \in L^2(X, d\mu)$ and $n = 0, 1, \dots$, define*

$$(Av_n f)(x) = \frac{1}{2n+1} \sum_{j=-n}^n f(T^j x) \tag{8.18}$$

Then μ is ergodic if and only if

$$\lim_{n \rightarrow \infty} \mu(|Av_n f|^2) = |\mu(f)|^2 \tag{8.19}$$

For (8.19) to hold, it suffices that it holds for a dense set, S , in $L^2(X, d\mu)$.

Proof (8.19) is equivalent to weak operator convergence as operators on $L^2(X, d\mu)$,

$$(Av_n)^*(Av_n) \xrightarrow{w} (1, \cdot)1$$

the projection onto 1, so since $\|Av_n\| \leq 1$, it suffices to prove it for a dense set.

If T is ergodic, then (8.15) implies (8.19). Conversely, if (8.19) holds, A is an invariant set, and χ_A is its characteristic function, then $Av_n(\chi_A) = \chi_A$ so (8.19) implies $\mu(A) = \mu(A)^2$, that is, $\mu(A)$ is 0 or 1. Thus, μ is ergodic. \square

Example 8.20 Let $X = \partial D$, the unit circle. Let α be an irrational number and let

$$T(e^{i\theta}) = e^{i(\theta+2\pi\alpha)}$$

Let $d\mu = d\theta/2\pi$ and $f_m = e^{im\theta} \in L^2(\partial D, d\mu)$. Then, for $m \neq 0$,

$$\begin{aligned} Av_n(f_m) &= (2n+1)^{-1} \left(\sum_{j=-n}^n e^{2\pi i j \alpha m} \right) f_m \\ &= (2n+1)^{-1} \frac{\sin(2\pi(n + \frac{1}{2})m\alpha)}{\sin(\pi m\alpha)} f_m \end{aligned}$$

hence $\|Av_n(f_m)\| \rightarrow 0$ if $m \neq 0$. Since $\{f_m\}_{m=0,\pm 1,\dots}$ are a basis of $L^2(\partial D, d\mu)$, (8.19) holds, so μ is ergodic. Notice $A = \{e^{2\pi i\alpha m}\}_{m=-\infty}^{\infty}$ is an invariant set but it has measure 0. It can be shown that μ is the only invariant measure in this case. \square

Example 8.21 Given a locally compact group, G , a unitary representation is a continuous map U taking G to the unitary operators on a Hilbert space, \mathcal{H} . Given such a representation, one can form the functions $F_{\varphi,U}(g) = \langle \varphi, U(g)\varphi \rangle$ for each $U \in \mathcal{H}$. One can show that as φ runs over all unit vectors and U over all representations, $\{F_{\varphi,U}\}$ forms a compact convex subset in $C(G)$ in the $\|\cdot\|_{\infty}$ -topology. Its extreme points will correspond to what are called irreducible representations, and one can use the Krein–Milman theorem to prove the existence of such representations. \square

Just the existence of extreme points in compact convex sets is powerful. The penultimate topic in this chapter provides proofs of two analytic results that would seem to have no direct connection to the Krein–Milman theorem. First, we provide a proof of the Stone–Weierstrass theorem; see, for example, [303, Appendix to Sect. IV.3] for the “usual” proof.

Theorem 8.22 (Stone–Weierstrass Theorem) *Let X be a compact Hausdorff space. Let A be a subalgebra of $\mathbb{C}_{\mathbb{R}}(X)$, the real-valued function on X , so that for any $x, y \in X$ and $\alpha, \beta \in \mathbb{R}$, there exists $f \in A$ so $f(x) = \alpha$ and $f(y) = \beta$. Then A is dense in $\mathbb{C}_{\mathbb{R}}(X)$ in $\|\cdot\|_{\infty}$.*

Proof $\mathcal{M}(X) = \mathbb{C}_{\mathbb{R}}(X)^*$ is the space of real signed measures on X with the total variation norm, that is, for any μ , there is a set, unique up to μ -measure zero sets, $B \subset X$ so $\mu \upharpoonright B \geq 0$, $\mu \upharpoonright X \setminus B \leq 0$, and $\|\mu\| = \mu(B) + |\mu(X \setminus B)|$.

Define

$$L = \{\mu \in \mathcal{M}(X) \mid \|\mu\| \leq 1; \mu(f) = 0 \text{ for all } f \in A\}$$

Then, since the unit ball in $\mathcal{M}(X)$ is compact in the weak-* topology, L is a compact convex set. If L is larger than $\{0\}$, L has an extreme point which necessarily has $\|\mu\| = 1$ since, if $0 < \|\mu\| \leq 1$, μ is a nontrivial concave combination of $\mu/\|\mu\|$ and 0.

If $g \in A$ and $\|g\|_{\infty} \leq 1$, then $g d\mu \in L$ for any $\mu \in L$ since $\int f(g d\mu) = \int (fg) d\mu = 0$ and $\|g d\mu\| \leq \|g\|_{\infty} \|\mu\|$. If $0 \leq g \leq 1$, then

$$\begin{aligned} & \|g d\mu\| + \|(1-g) d\mu\| \\ &= \int_B g d\mu + \int_B (1-g) d\mu - \int_{X \setminus B} g d\mu - \int_{X \setminus B} (1-g) d\mu = \|\mu\| \end{aligned}$$

so

$$\mu = \frac{g d\mu}{\|g d\mu\|} + \frac{(1-g) d\mu}{\|(1-g) d\mu\|}$$

is a convex combination of elements in L . Thus,

$$\begin{aligned} &\mu \text{ extreme and } 0 \leq g \leq 1 \\ &\text{with } g \in A \Rightarrow g = 0 \text{ a.e. } d\mu \text{ or } (1 - g) = 0 \text{ a.e. } d\mu \end{aligned} \tag{8.20}$$

If $\text{supp}(d\mu)$ has two points x, y , we can pick $f \in A$ with $f(x) = 1, f(y) = 2$. Thus, $g = f^2 / \|f\|_\infty^2$ has $0 \leq g \leq 1$ and $0 < g(x) < \frac{1}{4}$, and so $g \in (0, \frac{1}{4})$ in a neighborhood U of x . Since $\mu(U) \neq 0$, (8.20) fails. We conclude $\text{supp}(d\mu)$ is a single point, x . But then $\mu \in L$ implies $f(x) = 0$ for all $f \in A$, violating the assumption about $f(x) = \alpha$ can have any real value α .

This contradiction implies $L = \{0\}$ which, by the Hahn–Banach theorem, implies that A is dense. □

Remark The Stone–Weierstrass theorem does not hold if $\mathbb{C}_\mathbb{R}(X)$ is replaced by $\mathbb{C}(X)$, the complex-valued function. The canonical example of a nondense subalgebra of $\mathbb{C}(X)$ with the α, β property is the analytic functions on \mathbb{D} . It is a useful exercise to understand why the above proof breaks down in this case.

The second application concerns vector-valued measures, that is, measures with values in \mathbb{R}^ν , equivalently, n -tuples of signed real measures. Given such a measure, $\vec{\mu}$, one can form the scalar measure $\tilde{\mu} = \sum_{i=1}^N |\mu_i|$, which we suppose is finite. Then $d\mu_i = f_i d\tilde{\mu}$ with $f_i \in L^1$.

Definition A scalar measure, $d\mu$, is called *weakly nonatomic* if and only if for any A with $\mu(A) > 0$, there exists $B \subset A$ so $\mu(B) > 0$ and $\mu(A \setminus B) > 0$.

In much of the literature, what we have called weakly nonatomic is called nonatomic, but we defined nonatomic in Chapter 2 as a measure obeying Corollary 8.24 below. That corollary shows the definitions are equivalent, so one can drop “weakly” once one has the theorem.

Theorem 8.23 (Lyapunov’s Theorem) *Let $\vec{\mu}$ be a weakly nonatomic finite measure on (M, Σ) , a space with countably generated sigma algebra, and $\vec{f} \in L^1(M, \vec{\mu}; \mathbb{R}^\nu)$ fixed. Then*

$$\left\{ \int_A \vec{f} d\mu \mid A \subset M, \text{ measurable} \right\} \subset \mathbb{R}^\nu$$

is a compact convex subset of \mathbb{R}^ν .

Before proving this, we note a corollary and make some remarks:

Corollary 8.24 *Let μ be a σ -finite scalar positive measure which is weakly nonatomic. Then μ is nonatomic, that is, for any A and any $\alpha \in (0, \mu(A))$, there is $B \subset A$ with $\mu(B) = \alpha$.*

Proof By a simple approximation argument, we can suppose $\mu(A) < \infty$. Applying the theorem to $\mu \upharpoonright A$, we see $\{\mu(B) \mid B \subset A, \text{measurable}\} \subset \mathbb{R}$ is convex. Since $\mu(\emptyset) = 0$ and $\mu(A) = \mu(A)$, we see this convex set must be $[0, \mu(A)]$. \square

The main remark that helps us understand the proof is that the extreme points of $\{f \in L^\infty(M, d\mu) \mid 0 \leq f \leq 1\}$ are precisely the characteristic functions.

Proof of Theorem 8.23 Let $Q = \{g \in L^\infty \mid 0 \leq g \leq 1\}$. Then Q is a convex set, compact in the $\sigma(L^\infty, L^1)$ -topology, and $F: Q \rightarrow \mathbb{R}^\nu$ by

$$F(g) = \int g \vec{f} d\mu$$

is a continuous linear function, so $\{F(g) \mid g \in Q\}$ is a compact convex set, S . We will show that for any $\vec{\alpha} \in S$, there is $g = \chi_A \in Q$ with $F(g) = \vec{\alpha}$, so $\{\int_A \vec{f} d\mu \mid A \subset M\}$ is S , and so convex.

Let $Q_{\vec{\alpha}} = \{g \in Q \mid F(g) = \vec{\alpha}\}$. $Q_{\vec{\alpha}}$ is a closed subset of S and so a compact convex subset. By the Krein–Milman theorem, $Q_{\vec{\alpha}}$ has an extreme point g . We will prove $g = \chi_A$ using the fact that μ is weakly nonatomic.

Suppose for some $\varepsilon > 0$, $A = \{x \mid \varepsilon < g < 1 - \varepsilon\}$ has $\mu(A) > 0$. By induction, we can find B_1, \dots, B_{n+1} disjoint, so $\mu(B_j) > 0$ and $\cup_{j=1}^{n+1} B_j = A$. Let $\vec{\alpha}_j = \int_{B_j} \vec{f} d\mu$. Since \mathbb{R}^n has dimension n , we can find $(\beta_1, \dots, \beta_{n+1}) \in \mathbb{R}^{n+1}$, so that $\sum_{j=1}^{n+1} \beta_j \vec{\alpha}_j = 0$, some $\beta_j \neq 0$ and $|\beta_j| < \varepsilon$ for all j . Let

$$g_{\pm} = g \pm \sum \beta_j \chi_{B_j}$$

Since $|\beta_j| < \varepsilon$ and $\varepsilon < g < 1 - \varepsilon$ on B_j , we have that $0 \leq g_{\pm} \leq 1$. Since some $\beta_j \neq 0$, $g_+ \neq g_-$. Since $\sum \beta_j \vec{\alpha}_j = 0$, $g_{\pm} \in Q_{\vec{\alpha}}$. Clearly, $g = \frac{1}{2}g_+ + \frac{1}{2}g_-$, violating the fact that g is an extreme point of $Q_{\vec{\alpha}}$. It follows that g is 0 or 1 for a.e. x , that is, $g = \chi_A$ for some A . Thus, $\int_A \vec{f} d\mu = \vec{\alpha}$. \square

We end this chapter with a few results relating extreme points and linear or affine maps between spaces and sets. These will be needed in the next chapter.

Proposition 8.25 *Let X and Y be locally convex spaces and let A, B be compact convex subsets of X and Y , respectively. Let $T: X \rightarrow Y$ be a continuous linear map. Then if $T[\mathcal{E}(A)] \subset B$, we have that $T[A] \subset B$.*

Proof Since T is linear and B is convex, each $T(\sum_{i=1}^n \theta_i x_i)$ with $\sum_{i=1}^n \theta_i = 1$ and $x_i \in \mathcal{E}(A)$ lies in B . Then, since B is closed and T is continuous, the same is true of limits. Since $A = \text{ch}(\mathcal{E}(A))$, we see $T[A] \subset B$. \square

Definition Let X and Y be locally convex spaces and let A, B be convex subsets of X and Y , respectively. A map $T: A \rightarrow B$ is called *affine* if and only if for all $x, y \in A$ and $\theta \in [0, 1]$, $T(\theta x + (1 - \theta)y) = \theta T(x) + (1 - \theta)T(y)$.

Proposition 8.26 *Let A and B be compact convex subsets of locally convex spaces and let $T: A \rightarrow B$ be a continuous affine map. Then for any face, F , of B , $G \equiv T^{-1}[F]$, if nonempty, is a face of A .*

Proof G is closed since F is closed and T is continuous. If $x \in G$, $y, z \in A$, and $x = \theta y + (1 - \theta)z$ with $\theta \in (0, 1)$, then $T(x) \in F$, $T(y), T(z) \in B$, and $T(x) = \theta T(y) + (1 - \theta)T(z)$. Since F is a face, $T(y), T(z) \in F$, that is, $y, z \in G$. Thus, G is a face. \square