

# RANK ONE PERTURBATIONS WITH INFINITESIMAL COUPLING

A. KISELEV AND B. SIMON\*

Division of Physics, Mathematics, and Astronomy  
California Institute of Technology  
Pasadena, CA 91125

ABSTRACT. We consider a positive self-adjoint operator  $A$  and formal rank one perturbations

$$B = A + \alpha(\varphi, \cdot)\varphi$$

where  $\varphi \in \mathcal{H}_{-2}(A)$  but  $\varphi \notin \mathcal{H}_{-1}(A)$ , with  $\mathcal{H}_s(A)$  the usual scale of spaces. We show that  $B$  can be defined for such  $\varphi$  and what are essentially negative infinitesimal values of  $\alpha$ . In a sense we'll make precise, every rank one perturbation is one of three forms: (i)  $\varphi \in \mathcal{H}_{-1}(A)$ ,  $\alpha \in \mathbb{R}$ ; (ii)  $\varphi \in \mathcal{H}_{-1}$ ,  $\alpha = \infty$ ; or (iii) the new type we consider here.

## §1. Introduction

There has recently been considerable interest in the study of rank one perturbations of positive self-adjoint operators (see [11] and references therein). Let  $A \geq 0$  on a Hilbert space  $\mathcal{H}$  and consider

$$B = A + \alpha(\varphi, \cdot)\varphi. \tag{1.1}$$

Simon-Wolff [12] first pointed out that a natural framework for this was to consider  $\varphi \in \mathcal{H}_{-1}(A)$  where  $\mathcal{H}_s(A)$  is the usual scale of spaces associated to  $A$ ; that is, if  $s \geq 0$ ,  $\mathcal{H}_s(A) = D(|A|^{s/2})$  with the norm  $\|\cdot\|$  given by

$$\|\varphi\|_s^2 = \langle \varphi, (A+1)^s \varphi \rangle,$$

and if  $s < 0$ ,  $\mathcal{H}_s(A)$  is the completion of  $\mathcal{H}$  in the  $\|\cdot\|_s$  norm.  $\mathcal{H}_s \subset \mathcal{H}_t$  if  $s > t$  and one can define  $\mathcal{H}_\infty(A) = \bigcap_s \mathcal{H}_s(A)$  and  $\mathcal{H}_{-\infty}(A) = \bigcup_s \mathcal{H}_s(A)$ .  $\mathcal{H}_s^* = \mathcal{H}_{-s}$  in a natural way.

When  $\varphi \in \mathcal{H}_{-1}(A)$ ,  $\psi \mapsto |(\psi, \varphi)|^2$  defines a quadratic form on  $Q(A) = \mathcal{H}_{+1}(A)$ , which is  $A$ -bounded with relative bound zero. So the standard form perturbation theory ([7,10]) lets one define (1.1) for any  $\alpha \in \mathbb{R}$ .

---

\* This material is based upon work supported by the National Science Foundation under Grant No. DMS-9101715. The Government has certain rights in this material.

To appear in *J. Funct. Anal.*

Define

$$F_\alpha(z) = (\varphi, (A_\alpha - z)^{-1}\varphi) \quad (1.2a)$$

$$F(z) = F_{\alpha=0}(z). \quad (1.2b)$$

One easily proves the formulae (going back to Krein and Aronszajn):

$$F_\alpha(z) = F(z)/1 + \alpha F(z) \quad (1.3)$$

$$(A_\alpha - z)^{-1}\varphi = (1 + \alpha F(z))^{-1}(A - z)^{-1}\varphi \quad (1.4a)$$

$$(A_\alpha - z)^{-1} = (A - z)^{-1} - \alpha(1 + \alpha F(z))^{-1}((A - \bar{z})^{-1}\varphi, \cdot)(A - z)^{-1}\varphi. \quad (1.4b)$$

From (1.4) one sees  $\text{s-lim}_{\alpha \rightarrow \infty} (A_\alpha - z)^{-1}$  exists. If  $\varphi \notin \mathcal{H}_0(A) = \mathcal{H}$ , it defines an operator  $A_\infty$  on  $\mathcal{H}$ . This is studied in Gesztesy-Simon [5].

Our primary goal here is two-fold:

(a) To construct a family of rank one perturbations  $A + \alpha(\varphi, \cdot)\varphi$  where  $\varphi \notin \mathcal{H}_{-1}(A)$  but only in  $\mathcal{H}_{-2}(A)$ . Here  $\alpha$  is infinitesimal.

(b) Every pair of semibounded operators with  $(A + i)^{-1} - (B + i)^{-1}$  rank one can be written using the  $\alpha(\varphi, \cdot)\varphi$  construction with  $\varphi \in \mathcal{H}_{-1}$  and  $\alpha$  finite or infinite.

These two apparently paradoxical statements are not paradoxical because in (b) we did not specify if  $B$  is a perturbation of  $A$  or vice-versa. In fact, one can always label them so that  $A \leq B$ . Then we will show that  $B = A + \alpha(\varphi, \cdot)\varphi$  with  $\varphi \in \mathcal{H}_{-1}(A)$  with  $\alpha \in [0, \infty]$ . If  $\alpha < \infty$ , then  $A$  can be obtained from  $B$  by a rank one perturbation with  $\varphi \in \mathcal{H}_{-1}(B)$ . But if  $\alpha = \infty$ , it is necessary to use the  $\mathcal{H}_{-2}(B)$  construction to recover  $A$  from  $B$ .

At first, it is comforting that infinitesimal coupling is needed to undo infinite coupling, but that feeling is unfounded. For multiplicative perturbations, infinitesimal should undo infinite, but these perturbations are additive. In fact,  $(\eta, \cdot)\eta$  with  $\eta \in \mathcal{H}_{-2}(B)/\mathcal{H}_{-1}(B)$  is so infinite we need infinitesimal coupling to undo  $\infty(\varphi, \cdot)\varphi$  with  $\varphi \in \mathcal{H}_{-1}(A)$ .

A theme that we will explore in this paper is that if  $A, B$  have resolvents that differ by a rank one, then there exists a symmetric operator  $C$  with deficiency indices  $(1, 1)$  so that  $A$  and  $B$  are both self-adjoint extensions of  $C$ . To say that  $B$  is  $A + \alpha(\varphi, \cdot)\varphi$  with  $\alpha = \infty$  and  $\varphi \in \mathcal{H}_{-1}(A)$  (equivalently that  $A$  is  $B + \alpha(\varphi, \cdot)\varphi$  with  $\varphi \in \mathcal{H}_{-2}(B)/\mathcal{H}_{-1}(A)$  and  $\alpha$  infinitesimal) is equivalent to saying that  $B$  is the Friedrich's extension. From this point of view, our assertion (b) above is a special case of the Birman-Krein-Vishik theory of quadratic forms of positive self-adjoint extensions [3,8,13,6,2].

In §2, we present the construction of rank one perturbations with  $\varphi \in \mathcal{H}_{-2}$ . In §3, we use resolvent ordering to prove assertion (b). In §4, we explain the relation of infinite and infinitesimal coupling. In §5, we consider fairly general situations  $A_n = A + \alpha_n(\varphi_n, \cdot)\varphi_n$  with  $\varphi_n$  a cutoff of  $\varphi \in \mathcal{H}_{-\infty}(A)$  and show that as  $n \rightarrow \infty$ ,  $A_n$  converges to  $A$  in strong resolvent sense unless  $\varphi \in \mathcal{H}_{-1}(A)$  or  $\varphi \in \mathcal{H}_{-2}(A)$ ,  $\alpha_n < 0$  and  $\alpha_n \rightarrow 0$  at a suitable rate. This provides another view of the fact that the only rank one perturbations are the  $\mathcal{H}_{-1}(A)$  and  $\mathcal{H}_{-2}(A)$  constructions. In §6, we discuss the connection to the theory of self-adjoint extensions of deficiency indices  $(1, 1)$ . Finally, §7 presents some simple examples.

## §2. The Basic $\mathcal{H}_{-2}(A)$ Construction

Let  $\varphi \in \mathcal{H}_{-2}(A)$  so  $(A - z)^{-1}\varphi$  makes sense for any  $z \notin \text{spec}(A)$  and in particular, for  $\text{Im } z \neq 0$ . Motivated by (1.4), we try to construct a self-adjoint operator whose resolvent  $R(z)$  obeys

$$R(z) = (A - z)^{-1} - \sigma(z)K(z) \quad (2.1a)$$

where

$$K(z) = ((A - \bar{z})^{-1}\varphi, \cdot)(A - z)^{-1}\varphi. \quad (2.1b)$$

The idea is to define  $R(z)$  by (2.1) and then to pick the unknown function  $\sigma(z)$  in order that  $R$  obey the equation obeyed by any resolvent:

$$\frac{dR}{dz} = R(z)^2. \quad (2.2)$$

Since  $\frac{dK}{dz} = (A - z)^{-1}K + K(A - z)^{-1}$  and  $\frac{d}{dz}(A - z)^{-1} \equiv (A - z)^{-2}$ , (2.2) is equivalent to

$$\frac{d\sigma}{dz} K(z) = -\sigma(z)^2 K(z)^2. \quad (2.3)$$

But  $K(z)^2 = K(z)(\varphi, (A - z)^{-2}\varphi)$ . Thus (2.2) is equivalent to

$$\frac{d}{dz} \sigma^{-1}(z) = (\varphi, (A - z)^{-2}\varphi). \quad (2.4)$$

Supposing that  $A \geq 0$ , we note that (2.4) shows that  $\sigma^{-1}$ , originally defined for  $\text{Im } z \neq 0$ , can be continued through  $(-\infty, 0)$ . Self-adjointness for  $R$ , that is,  $R^*(z) = R(\bar{z})$  requires  $\sigma^{-1}$  be real there; and thus the solutions can be written

$$\sigma^{-1}(z) = \beta + (\varphi, [(A - z)^{-1} - (A + 1)^{-1}]\varphi) \quad (2.5)$$

with  $\beta$  real and equal to  $\sigma^{-1}(-1)$ . This motivates:

**Theorem 2.1.** *Fix  $\beta \in \mathbb{R}$ . Suppose  $A \geq 0$  and  $\varphi \in \mathcal{H}_{-2}(A)$ . For  $\text{Im } z \neq 0$ , define  $R_\beta(z)$  by (2.1) with  $\sigma(z)$  given by (2.5). Then there is a self-adjoint operator  $\tilde{A}_\beta$  with  $R_\beta(z) = (\tilde{A}_\beta - z)^{-1}$ .*

*Proof.* Let

$$G(z) \equiv (\varphi, [(A - z)^{-1} - (A + 1)^{-1}]\varphi). \quad (2.6)$$

Then for  $y \in (-\infty, 0)$ ,  $\frac{dG}{dy} = (\varphi, (A - y)^{-2}\varphi) > 0$ . Thus, there is at most one  $y < 0$ , call it  $y_0$ , so  $\sigma(y)^{-1} = 0$ . Therefore,  $R_\beta(z)$  extends to  $\mathbb{C} \setminus [0, \infty) \cup \{y_0\}$  with  $R_\beta(y)$  self-adjoint if  $y \in \mathbb{R} \setminus [0, \infty) \cup \{y_0\}$ . Fix any  $y_1 < 0$  with  $y_1 \neq y_0$  and define  $A_\beta \equiv R_\beta(y_1)^{-1} - y_1$ . Then  $R_\beta(z)$  and  $(A_\beta - z)^{-1}$  obey the same differential equation (1.2) and same initial conditions at  $y = y_1$ , and so they are equal on  $\text{Im } z \neq 0$ .

*Remark.* One can think of (2.1) in the form

$$\begin{aligned} (\tilde{A}_\beta - z)^{-1} &= (A - z)^{-1} - \sigma_\beta(z)K(z) \\ \sigma_\beta(z)^{-1} &= \beta + (\varphi, ((A - z)^{-1} - (A + 1)^{-1}\varphi) \end{aligned} \quad (2.1c)$$

as a renormalized form of (1.4), which can be written

$$\begin{aligned} (A_\alpha - z)^{-1} &= (A - z)^{-1} - \hat{\sigma}_\alpha(z)K(z) \\ \hat{\sigma}_\alpha(z)^{-1} &= \alpha^{-1} + (\varphi, (A - z)^{-1}\varphi). \end{aligned}$$

If  $\varphi \in \mathcal{H}_{-1}(A)$ , then  $\tilde{A}_\beta = A_\alpha$  where  $\beta$  and  $\alpha$  are related by

$$\beta = \alpha^{-1} + (\varphi, (A + 1)^{-1}\varphi). \quad (2.7)$$

If  $\varphi \notin \mathcal{H}_{-1}$ , in essence we need to take  $\alpha^{-1} = -\infty$  to undo the divergence of  $(\varphi, (A+1)^{-1}\varphi)$ , and  $\alpha$  is infinitesimal and negative. The condition  $\varphi \in \mathcal{H}_{-2}(A)$  is required for the single renormalization to work.

**Theorem 2.2.** *If  $\varphi \notin \mathcal{H}_{-1}(A)$ , then each operator  $A_\beta$  defined in Theorem 2.1 obeys  $\tilde{A}_\beta \leq A$  with  $\tilde{A}_\beta \neq A$ . If  $\varphi \in \mathcal{H}_{-1}(A)$ , there exist  $\tilde{A}_\beta$ 's with  $\tilde{A}_\beta \geq A$  with  $\tilde{A}_\beta \neq A$ .*

*Remark.* Recall ([7]) that we say  $A, B$  obey  $A \geq B$  if and only if there is  $a \in \mathbb{R}$  with  $A \geq a1$ ,  $B \geq a1$ ; and for  $z < a$  real, we have  $(B - z)^{-1} \geq (A - z)^{-1}$  as bounded operators.

*Proof.* If  $\varphi \in \mathcal{H}_{-1}(A)$ , we have seen above that  $\{\tilde{A}_\beta\}$  is the same as  $\{A_\alpha\}$  using (2.7). Since  $A_\alpha \geq A$  if  $\alpha > 0$ , that proves the  $\mathcal{H}_{-1}$  result.

If  $\varphi \notin \mathcal{H}_{-1}$ , then  $G(-y) \rightarrow -\infty$  as  $y \rightarrow \infty$ . Thus, there is some  $y_0 \in (-\infty, 0)$ , so  $G(y) + \beta < 0$  for all  $y \leq y_0$ . By (2.5) and (2.1c),  $(\tilde{A}_\beta - y)^{-1} \geq (A - y)^{-1} > 0$  for such  $y$ , so  $\tilde{A}_\beta \geq y_0$ ,  $A \geq y_0$ , and  $\tilde{A}_\beta \leq A$ .

### §3. Every Rank One Perturbation Is $\mathcal{H}_{-1}(A)$ -bounded

In this section, we want to consider pairs of operators  $A, B$  so that  $(A + i)^{-1} - (B + i)^{-1}$  is rank one. We start with two results that illuminate the notion:

**Proposition 3.1.** *Let  $A, B$  be self-adjoint operators. Then  $Q(z) = (A - z)^{-1} - (B - z)^{-1}$  is rank one for one  $z$  with  $\text{Im } z \neq 0$  if and only if it is rank one for all such  $z$ .*

*Proof.*

$$(A - z)^{-1} = (1 + (w - z)(A - z)^{-1})(A - w)^{-1}, \quad (3.1)$$

so using the fact that

$$(\varphi, (A - z)^{-1} - (B - z)^{-1}\psi) = ((A - \bar{z})^{-1}\varphi, B(B - z)^{-1}\psi) - (A(A - \bar{z})^{-1}\varphi, (B - z)^{-1}\psi),$$

we see that

$$Q(z) = (1 + (w - z)(A - z)^{-1})Q(w)(1 + (w - z)(B - z)^{-1})$$

and so  $\text{Rank } Q(z) \leq \text{Rank } Q(w)$ .

**Proposition 3.2.** *Suppose that  $A, B$  are self-adjoint,  $A \geq 0$ , and  $(A + i)^{-1} - (B + i)^{-1}$  is rank one. Then  $B$  is bounded from below.*

*Proof.* By (3.1) for  $B$ ,  $w \in (-\infty, 0)$  is in  $\text{spec}(B)$  if and only if  $1 + (w - i)(B - i)^{-1}$  is not invertible. But

$$\begin{aligned} L(w) &= 1 + (w - i)(B - i)^{-1} = 1 + (w - i)(A - i)^{-1} + (w - i)((B - i)^{-1} - (A - i)^{-1}) \\ &= L_1(w) + L_2(w), \end{aligned}$$

where  $L_1 = 1 + (w - i)(A - i)^{-1} = (A - w)(A - i)^{-1}$  is invertible for  $w \in (-\infty, 0)$  and  $L_2 = (w - i)((B - i)^{-1} - (A - i)^{-1})$  is rank one.

Thus,  $L(w)$  is invertible if and only if  $1 + L_1(w)^{-1}L_2(w)$  is invertible. By (3.1),  $w \in \text{spec}(B)$  if and only if  $1 + L_1(w)^{-1}L_2(w)$  is not invertible. Thus, since  $L_2$  is rank one,  $w \in \text{spec}(B)$  if and only if  $F(w) \equiv \text{Tr}(L_1(w)^{-1}L_2(w)) = -1$ .  $F$  is an entire analytic function with  $F(w) \neq -1$  if  $\text{Im } w \neq 0$ . We conclude  $B$  has isolated point spectrum on  $(-\infty, 0)$ .

Thus, there exist real  $w_0$  with  $F(w_0) \neq -1$  and so  $(B - w_0)^{-1} - (A - w_0)^{-1}$  is rank one. For rank one perturbations of self-adjoint operators, eigenvalues intertwine. Since  $A$  has no eigenvalues in  $(-\infty, 0)$ ,  $B$  can have only one eigenvalue in  $(-\infty, 0)$ ; that is,  $B$  is bounded from below.

**Corollary 3.3.** *If  $A \geq 0$  and  $(A + i)^{-1} - (B + i)^{-1}$  is rank one, then either  $A \geq B$  or  $B \geq A$ .*

*Proof.* Pick  $w$  below  $\text{spec}(A) \cup \text{spec}(B)$ . Then  $(A - w)^{-1} \geq 0$ ,  $(B - w)^{-1} \geq 0$ , and since  $(A - w)^{-1} - (B - w)^{-1}$  is rank one and self-adjoint, either  $(A - w)^{-1} \geq (B - w)^{-1}$  or  $(B - w)^{-1} \geq (A - w)^{-1}$ . It follows that either  $A \geq B$  or  $B \geq A$ .

**Theorem 3.4.** *Let  $A, B$  be self-adjoint operators with  $B \geq A \geq 0$ . Suppose that  $(A + 1)^{-1} - (B + 1)^{-1}$  is rank one. Then  $B = A + \alpha(\varphi, \cdot)\varphi$  with  $\varphi \in \mathcal{H}_{-1}(A)$  and  $\alpha \in [0, \infty]$  (with  $\alpha = \infty$  allowed).*

*Proof.* Write

$$(A + 1)^{-1} = (B + 1)^{-1} + (\eta, \cdot)\eta, \quad (3.2)$$

which we can do because  $(A + 1)^{-1} \geq (B + 1)^{-1}$ .

We claim that  $\eta \in \mathcal{H}_{+1}(A)$  with  $(\eta, (A + 1)\eta) \leq 1$ ; see Lemma 3.5 below. Define  $\varphi = (A + 1)\eta$  so (3.2) becomes

$$(B + 1)^{-1} = (A + 1)^{-1} - ((A + 1)^{-1}\varphi, \cdot)(A + 1)^{-1}\varphi,$$

which is just (1.4) if

$$\frac{\alpha}{1 + \alpha(\varphi, (A + 1)^{-1}\varphi)} = 1$$

or

$$\alpha = \frac{1}{1 - (\eta, (A + 1)\eta)} \quad (3.3)$$

where  $(\eta, (A + 1)\eta) = 1$  corresponds to  $\alpha = \infty$ . (1.4) at  $z = -1$  implies the general relation for all  $z$ .

**Lemma 3.5.** *Let  $A \geq 0$  be self-adjoint. Suppose  $\eta \in \mathcal{H}$  with  $(\eta, \cdot)\eta \leq (A + 1)^{-1}$ . Then  $\eta \in \mathcal{H}_{+1}(A)$  with  $(\eta, (A + 1)\eta) \leq 1$ .*

*Proof.* Let  $E_k$  be the spectral projection  $E_{[0, k]}(A)$ . Let  $\varphi_k = (A + 1)E_k\eta$ . Then, by hypothesis,

$$|(\eta, \varphi_k)|^2 \leq (\varphi_k, (A + 1)^{-1}\varphi_k). \quad (3.4)$$

(3.4) is equivalent to

$$(\eta, E_k(A + 1)\eta)^2 \leq (\eta, E_k(A + 1)\eta)$$

or

$$(\eta, E_k(A + 1)\eta) \leq 1.$$

Taking  $k \rightarrow \infty$ , we see  $\eta \in \mathcal{H}_{+1}(A)$  and  $(\eta, (A + 1)\eta) \leq 1$ .

*Remark.* It may seem puzzling that the  $\alpha$  in (3.3) obeys  $1 < \alpha \leq \infty$ . How about  $B = A + \alpha(\varphi, \cdot)\varphi$  with  $\alpha < 1$ ? The resolution is that until we normalize  $\varphi$  in some way, the scale of  $\alpha$  is irrelevant. If we demand  $\tilde{\varphi}$  obey  $(\tilde{\varphi}, (A + 1)^{-1}\tilde{\varphi}) = 1$ , then we take  $\tilde{\varphi} = \varphi/(\eta, (A + 1)\eta)^{1/2}$  and  $\alpha(\varphi, \cdot)\varphi = \tilde{\alpha}(\tilde{\varphi}, \cdot)\tilde{\varphi}$  where now

$$\tilde{\alpha} = \frac{(\eta, (A + 1)\eta)}{1 - (\eta, (A + 1)\eta)}.$$

As  $(\eta, (A + 1)\eta)$  runs from 0 to 1,  $\tilde{\alpha}$  runs from 0 to infinity.

As an application of Lemma 3.5, we return to the construction of §2:

**Theorem 3.6.** *Suppose  $A \geq 0$ ,  $\varphi \in \mathcal{H}_{-2}(A)$  but  $\varphi \notin \mathcal{H}_{-1}(A)$ , and that  $\tilde{A}_\beta$  is the operator of Theorem 2.1. Then*

- (i)  $\mathcal{H}_{+1}(\tilde{A}_\beta) \supset \mathcal{H}_{+1}(A)$
- (ii)  $\mathcal{H}_{+1}(\tilde{A}_\beta) \neq \mathcal{H}_{+1}(A)$ .

*Remark.* We'll see later in §6 that  $\mathcal{H}_{+1}(A)$  has codimension 1 in  $\mathcal{H}_{+1}(\tilde{A}_\beta)$ .

*Proof.* By Theorem 2.2,  $\tilde{A}_\beta \leq A$  which implies (i). To see (ii), note that by the construction in §2 for all sufficiently large  $c > 0$ ,

$$(A_\beta + c)^{-1} = (A + c)^{-1} - \sigma(c)((A + c)^{-1}\varphi, \cdot)(A + c)^{-1}\varphi$$

with  $\sigma(c) < 0$ . Thus by Lemma 3.5,  $(A + c)^{-1}\varphi \in \mathcal{H}_{+1}(\tilde{A}_\beta)$ . Since  $\varphi \notin \mathcal{H}_{-1}(A)$ , we have that  $(A + c)^{-1}\varphi \notin \mathcal{H}_{+1}(A)$ .

#### §4. Relation to Infinite Coupling

Suppose  $B = A + \alpha(\varphi, \cdot)\varphi$  with  $\varphi \in \mathcal{H}_{-1}(A)$ . If  $\alpha < \infty$ , then  $\mathcal{H}_{+1}(B) = \mathcal{H}_{+1}(A)$  and  $A = B - \alpha(\varphi, \cdot)\varphi$  so  $A$  can be recovered from  $B$  by the  $\mathcal{H}_{-1}$  construction. Our goal here is to show that when  $\alpha = \infty$ ,  $A$  can be recovered from  $B$  by the  $\mathcal{H}_{-2}(B)$  construction of §2, and vice-versa that the  $A \rightarrow \tilde{A}_\beta$  construction can be undone with infinite coupling.

Recall ([5]) if  $\varphi \in \mathcal{H}_{-1}(A)$  but  $\varphi \notin \mathcal{H}$  and  $A_\infty = A + \infty(\varphi, \cdot)\varphi$ , then there exists a natural  $\eta \in \mathcal{H}_{-2}(A_\infty)$  which obeys

$$(A_\infty - z)^{-1}\eta = F(z)^{-1}(A - z)^{-1}\varphi \quad (4.1)$$

with  $F$  given by (1.2b).

**Proposition 4.1.** *Suppose  $A \geq 0$ ,  $\varphi \in \mathcal{H}_{-1}(A)$  but  $\varphi \notin \mathcal{H}$ , and  $\eta$  is given by (4.1). Then  $\eta \notin \mathcal{H}_{-1}(A_\infty)$ .*

*Proof.*  $\eta \in \mathcal{H}_{-1}(A_\infty)$  if and only if  $\lim_{c \rightarrow \infty} \left( \eta, \frac{c}{A_\infty + c} \frac{1}{A_\infty + 1} \eta \right)$  is finite. But by (4.1)

$$\left( \eta, \frac{c}{A_\infty + c} \frac{1}{A_\infty + 1} \eta \right) = \frac{1}{F(-1)F(-c)} \left( \varphi, \frac{c}{A + c} \frac{1}{A + 1} \varphi \right).$$

The expectation on the right side of this equation has a non-zero limit as  $c \rightarrow \infty$  since  $\varphi \in \mathcal{H}_{-1}(A)$ . But  $F(-c) \rightarrow 0$  as  $c \rightarrow \infty$  so the limit is infinity; that is,  $\eta \notin \mathcal{H}_{-1}(A_\infty)$ .

**Theorem 4.2.** *Suppose  $A \geq 0$  and  $\varphi \in \mathcal{H}_{-1}(A)$  but  $\varphi \notin \mathcal{H}$ . Let  $B \equiv A_\infty = A + \alpha(\varphi, \cdot)\varphi$ . Then for some  $\beta$  and the perturbation  $\eta$ ,  $\tilde{B}_\beta = A$ ; that is,  $A$  can be recovered from  $B$  by the construction of §2.*

*Proof.* By (1.4b) in the limit

$$(B - z)^{-1} = (A - z)^{-1} - F(z)^{-1}((A - \bar{z})^{-1}\varphi, \cdot)(A - z)^{-1}\varphi.$$

By (4.1)

$$(A - z)^{-1} = (B - z)^{-1} + F(z)((B - \bar{z})^{-1}\eta, \cdot)(B - z)^{-1}\eta$$

which shows that  $(A + 1)^{-1}$  is a  $(\tilde{B}_\beta + 1)^{-1}$ .

*Remark.* By §2, the coefficient in front of  $((B - \bar{z})^{-1}\varphi, \cdot)(B - z)^{-1}\varphi$  should be  $(\beta + G(z))^{-1}$  where  $G(z) = (\eta, [(A_\infty - z)^{-1} - (A_\infty + 1)^{-1}]\eta)$ . The resulting relation of  $\text{Im } F(z)^{-1}$  and  $\text{Im } (G(z))$  is exactly what was found in [5].

## §5. Limits

We've shown in the last two sections that if  $(A - z)^{-1} - (B - z)^{-1}$  is rank one (and both are bounded below), then  $B$  can be recovered from  $A$  via either a  $\varphi \in \mathcal{H}_{-1}(A)$  construction with  $\alpha \in (-\infty, \infty]$  or else by the  $\varphi \in \mathcal{H}_{-2}(A) \setminus \mathcal{H}_{-1}(A)$  construction with  $\alpha$  infinitesimal. Thus it should be impossible to define  $A + \alpha(\varphi, \cdot)\varphi$  if  $\varphi \notin \mathcal{H}_{-2}(A)$ . That is what we'll prove in this section.

**Theorem 5.1.** *Let  $A \geq 0$  and  $\varphi \in \mathcal{H}_{-\infty}(A)$ . Let  $\varphi_n = E_{[0, n]}(A)\varphi$  and*

$$A_n = A + \alpha_n(\varphi_n, \cdot)\varphi_n.$$

*Then:*

- (i) *If  $\varphi \notin \mathcal{H}_{-2}(A)$ , then for any choice of  $\alpha_n$ ,  $(A_n - z)^{-1}$  converges to  $(A - z)^{-1}$  strongly as  $n \rightarrow \infty$  for any  $z \in \mathbb{C} \setminus \mathbb{R}$ .*
- (ii) *If  $\varphi \notin \mathcal{H}_{-1}(A)$  and  $\alpha_n \geq 0$ , then for any choice of  $\alpha_n$  (subject to  $\alpha_n \geq 0$ ),  $(A_n - z)^{-1}$  converges to  $(A - z)^{-1}$  strongly as  $n \rightarrow \infty$  for any  $z \in \mathbb{C} \setminus \mathbb{R}$ .*
- (iii) *If  $\varphi \notin \mathcal{H}_{-1}(A)$  and  $\alpha_n \rightarrow \alpha_\infty \neq 0$ , then for any choice of  $\alpha_n$  (subject to  $\alpha_n \rightarrow \alpha_\infty$ ),  $(A_n - z)^{-1}$  strongly to  $(A - z)^{-1}$  as  $n \rightarrow \infty$  for any  $z \in \mathbb{C} \setminus \mathbb{R}$ .*

*Remarks.* 1. Thus to get a non-trivial limit, we either need  $\varphi \in \mathcal{H}_{-1}(A)$  or else  $\varphi \in \mathcal{H}_{-2}(A)$  and  $\alpha_n$  negative and infinitesimal.

2. In cases (ii) and (iii), if  $\varphi \in \mathcal{H}_{-2}(A)$ , our proof shows norm convergence.

*Proof.* By general principles [9], weak convergence of resolvents implies strong convergence. Since the  $\{(A_n - z)^{-1}\}$  are uniformly bounded for fixed  $z \in \mathbb{C} \setminus \mathbb{R}$ , it suffices to prove convergence of  $(\psi_1, (A_n - z)^{-1}\psi_2)$  for  $\psi_i \in \mathcal{H}_\infty$ .

By (1.4b),

$$(A_n - z)^{-1} = (A - z)^{-1} - [\alpha_n^{-1} + (\varphi_n(A - z)^{-1}\varphi_n)]^{-1}((A - \bar{z})^{-1}\varphi_n, \cdot)(A - z)^{-1}\varphi_n. \quad (5.1)$$

Since  $(\psi, (A - z)^{-1}\varphi_n)$  is uniformly bounded if  $\psi \in \mathcal{H}_{+\infty}(A)$  (since  $\varphi \in \mathcal{H}_{-\infty}(A)$ ), strong convergence is equivalent to

$$|\gamma_n| \equiv |\alpha_n^{-1} + (\varphi_n, (A - z)^{-1}\varphi_n)| \rightarrow \infty.$$

Now

$$\operatorname{Im} \gamma_n = (\operatorname{Im} z) \|(A - z)^{-1}\varphi_n\|^2$$

goes to infinity as  $n \rightarrow \infty$  if  $\varphi \notin \mathcal{H}_{-2}$ , so (i) is proven.

Suppose now  $\varphi \in \mathcal{H}_{-2}$ . Since

$$\operatorname{Re} \gamma_n = \alpha_n^{-1} + (\varphi_n, A[(A - \operatorname{Re} z)^2 + (\operatorname{Im} z)^2]^{-1}\varphi_n) - \operatorname{Re} z \|(A - z)^{-1}\varphi_n\|^2,$$

we see that if  $\alpha_n > 0$  and  $\varphi_n \notin \mathcal{H}_{-1}(A)$ , then  $\operatorname{Re} \gamma_n \rightarrow \infty$ , and similarly if  $\alpha_n^{-1}$  has a finite limit  $\operatorname{Re} \gamma_n \rightarrow \infty$ .

*Remark.* Friedman [4] has shown that if  $V_n$  are functions on  $\mathbb{R}^\nu$  with  $\operatorname{supp} V_n \subset \{x \mid |x| < n^{-1}\}$  and  $H_n = -\Delta + V_n$ , then if  $\nu \geq 2$ ,  $H_n \rightarrow H$  in strong resolvent sense if  $V_n \geq 0$  (irrespective of how big  $V_n$  is); and if  $\nu \geq 4$ ,  $H_n \rightarrow H$  with no positivity assumption. Notice that  $\delta_0 \in \mathcal{H}_{-\alpha}(-\Delta)$  if and only if  $2\alpha > \nu$ . Thus  $\delta_0 \in \mathcal{H}_{-1}$  only if  $\nu < 2$  and  $\delta_0 \in \mathcal{H}_{-2}$  if and only if  $\nu < 4$ . We can therefore regard Theorem 5.1 as a kind of analog of Friedman's results.

## §6. Self-Adjoint Extensions

The punchline of this section is that rank one perturbations of  $A \geq 0$  is really the same as the theory of self-adjoint extensions of deficiency indices  $(1, 1)$  of a positive operator. From this point of view, the  $\alpha = \infty$  operator found by Gesztesy-Simon [5] is exactly the Friedrich's extension.

Let  $A \geq 0$  and  $\varphi \in \mathcal{H}_{-2}(A)$ . Whatever  $A_\alpha = A + \alpha(\varphi, \cdot)\varphi$  is to mean  $A_\alpha\psi$  should equal  $A\psi$  if  $(\varphi, \psi) = 0$ . Thus, define

$$D_\varphi = \{\psi \in D(A) \mid (\varphi, \psi) = 0\}.$$

Since  $\varphi \in \mathcal{H}_{-2}(A)$ ,  $(\varphi, \psi)$  is defined for  $\psi \in D(A) = \mathcal{H}_{+2}(A)$ .



**Lemma.** *Let  $A_0 = A \upharpoonright D_\varphi$  with domain  $D_\varphi$ . Then  $A_0$  has deficiency indices  $(1, 1)$ .*

*Proof.* It suffices to prove that  $\text{Ran}(A_0 + 1)$  has codimension 1. But by definition,  $\psi \in D_\varphi$  if and only if  $(A + 1)\psi$  is orthogonal to  $(A + 1)^{-1}\varphi$ ; that is,  $\text{Ran}(A_0 + 1) = \{(A + 1)^{-1}\varphi\}^\perp$  has codimension 1.

The rank one perturbations are thus the self-adjoint extensions of  $A_0$ . Deficiency one extension of semibounded operators (and generally semibounded extensions of semibounded operators) have been studied extensively [3,8,13,6,2]. The result of this theory is that these are parametrized by a single parameter  $\gamma$  which runs in  $(-\infty, \infty]$  with  $+\infty$  allowed. They are best described in terms of quadratic forms. The operator  $A^{(\infty)}$  is the Friedrich's extension and has form domain  $Q(A^{(\infty)})$ . There is a vector  $\xi$  defined by  $(A_0 + 1)^*\xi = 0$  and for  $\gamma \neq \infty$ ,

$$Q(A^{(\gamma)}) = Q(A^{(\infty)}) \cup \{\lambda\xi\}_{\lambda \in \mathbb{C}},$$

where  $\cup$  means disjoint sums and

$$((\psi + \lambda\xi), A^{(\gamma)}(\psi + \lambda\xi)) = (\psi, A^{(\infty)}\psi) + \lambda^2\gamma.$$

$\xi$  is easily seen to be  $(A + 1)^{-1}\varphi$ .

The original operator  $A$  is some  $A^{(\gamma_0)}$ . If  $A = A^{(\gamma_0)}$  with  $\gamma_0 \neq \infty$ , then the  $A^{(\gamma)}$  are precisely  $\{A + c(\gamma - \gamma_0)(\varphi, \cdot)\varphi\}$  for a suitable constant  $c$  ( $= (\varphi, (A + 1)^{-1}\varphi)$ ). The  $\gamma = \infty$  operator is exactly a Friedrich's extension.

If  $\gamma_0 = \infty$ , we see in this situation where the other  $A^{(\gamma)}$ 's are gotten by the construction in §2.

## §7. Examples

**Example 1.** Take  $A = -\Delta$  on  $L^2(\mathbb{R}^\nu)$ . We want to see what  $\varphi$  can be used for rank one perturbations defined at a single point 0. Since  $\varphi$  is supported at 0,  $\varphi \in \mathcal{H}_{-\infty}(A)$  means  $\varphi$  is a distribution, so its Fourier transform is a polynomial  $P$  in  $p$ . For  $\varphi \in \mathcal{H}_{-1}(A)$ , we need

$$\int \frac{d^\nu p |P(p)|^2}{(p^2 + 1)} < \infty. \quad (7.1)$$

This can only happen if  $\nu = 1$  and  $P$  has degree 0, that is,  $\varphi = \delta(x)$ . For  $\varphi$  to be in  $\mathcal{H}_{-2}(A)$ , we need the analog of (7.1) with  $(p^2 + 1)$  replaced by  $(p^2 + 1)^2$ . This allows  $P$  of degree 0 if  $\nu = 2, 3$  and degree 1 if  $\nu = 1$ . Thus, the rank one theory works exactly for  $\delta(x)$  in  $\nu = 1, 2, 3$  and  $\delta'(x)$  in  $\nu = 1$ . The  $\mathcal{H}_{-2}(A)$  construction exactly corresponds to point interactions as discussed extensively (see [1] and references therein.) Of course, our construction specialized to this case is just the standard one for point interactions; so our construction in §2 can be viewed as an abstraction of that method. One thing one can look at is undoing the point interaction in dimension 2 and 3. For concreteness, take  $\nu = 3$ . Then  $\mathcal{H}_{+1}(\tilde{A}_\beta)$  is strictly bigger than  $\mathcal{H}_{+1}(A)$ . The extra functions have a Coulomb singularity at  $x = 0$ ; that is,  $\psi \in \mathcal{H}_{+1}(\tilde{A}_\beta)$  has the form

$$\psi(x) = ce^{-\mu|x|}|x|^{-1} + \tilde{\psi}$$

with  $\tilde{\psi} \in \mathcal{H}_{+1}(-\Delta)$ .  $\mu$  is a convenient parameter;  $c$  is independent of  $\mu$ . One can think of  $c$  is formally given by  $\lim_{|x| \rightarrow 0} |x|\psi(x)$ . Since  $\psi$  is not bounded, we can't use that definition but can use

$$c(\psi) = \lim_{r \rightarrow 0} r \frac{3}{4\pi r^3} \int_{|x| \leq r} \psi(x) d^3x.$$

So  $c$  defines a vector  $\varphi \in \mathcal{H}_{-1}(\tilde{A}_\beta)$  and the various  $\tilde{A}_\beta$ 's are just  $\tilde{A}_{\beta_0} + \alpha(\varphi, \cdot)\varphi$  for  $\alpha \in (-\infty, \infty)$ .  $\alpha = \infty$  recovers the original Laplacian.

**Example 2.** Let  $A$  be  $-\frac{d^2}{dx^2}$  on  $L^2(0, \infty)$  with Neumann boundary condition at zero. Let  $\varphi(x) = \delta(x) \in \mathcal{H}_{-1}(A)$ . Then  $A + \alpha(\varphi, \cdot)\varphi$  precisely corresponds to the boundary conditions

$$\sin(\theta)u'(0) + \cos(\theta)u(0) = 0$$

where  $\alpha = -\cot(\theta)$ .  $\alpha = \infty$  corresponds to Dirichlet boundary condition. The corresponding  $\eta$  as discussed in [5] is just  $\delta'(x)$ ; that is,  $\delta' \in \mathcal{H}_{-2}(A_\infty)$ . The construction in §2 tells us how to reconstruct  $A_\theta$  from  $A_\infty$ .

## REFERENCES

- [1] S. Albeverio, F. Gesztesy, R. Høegh-Krohn, and H. Holden, *Solvable Models in Quantum Mechanics*, Springer, 1988.
- [2] A. Alonso and B. Simon, *The Birman-Krein-Vishik theory of self-adjoint extensions of semibounded operators*, J. Operator Theory **4** (1980), 251–270.
- [3] M.S. Birman, *On the self-adjoint extensions of positive definite operators*, Math. Sb. **38** (1956), 431–450. (Russian)
- [4] C.N. Friedman, *Perturbations of the Schrödinger equation by potentials with small support*, J. Funct. Anal. **10** (1972), 346–360.
- [5] F. Gesztesy and B. Simon, *Rank one perturbations at infinite coupling*, J. Funct. Anal. **128** (1995), 245–252.
- [6] G. Grubb, *A characterization of the non-local boundary value problems associated with an elliptic operator*, Ann. Scuola Norm. Sup. Pisa **22** (1968), 425–513.
- [7] T. Kato, *Perturbation Theory for Linear Operators*, 2nd edition, Springer, 1980.
- [8] M. Krein, *The theory of self-adjoint extensions of semibounded Hermitian transformations and its applications, I.*, Rec. Math. (Math. Sb.) **20** (1947), 431–495.
- [9] M. Reed and B. Simon, *Methods of Modern Mathematical Physics, I. Functional Analysis*, Academic Press, 1972.
- [10] ———, *Methods of Modern Mathematical Physics, II. Fourier Analysis, Self-Adjointness*, Academic Press, 1975.
- [11] B. Simon, *Spectral analysis of rank one perturbations and applications*, Proc. 1993 Vancouver Summer School in Mathematical Physics (to appear).
- [12] B. Simon and T. Wolff, *Singular continuous spectrum under rank one perturbations and localization for random Hamiltonians*, Commun. Pure Appl. Math. **39** (1986), 75–90.
- [13] M. Vishik, *On general boundary conditions for elliptic differential equations*, Trudy Moskov. Mat. Obšč. **1** (1952), 187–246. (Russian)