

Weak containment in the space of actions of a free group

Alexander S. Kechris

(A) We consider measure preserving actions of an infinite, countable (discrete) group Γ on non-atomic standard measure spaces (X, μ) , i.e., standard Borel spaces equipped with a non-atomic probability Borel measure. (All such measure spaces are isomorphic to $([0, 1], \lambda)$, where λ is Lebesgue measure.) We denote by $A(\Gamma, X, \mu)$ the space of such actions. If $a \in A(\Gamma, X, \mu)$ and $\gamma \in \Gamma$, we denote by $\gamma^a(x) = a(\gamma, x)$, the corresponding automorphism of the space (X, μ) . The group $\text{Aut}(X, \mu)$ admits a canonical Polish topology, called the *weak topology*, which is the topology generated by the maps $T \mapsto T(A)$ (A a Borel subset of X) from $\text{Aut}(X, \mu)$ to the measure algebra $\text{MALG}(X, \mu)$ of (X, μ) , equipped with the metric $d_\mu(A, B) = \mu(A \Delta B)$ and the corresponding topology. Since $A(\Gamma, X, \mu)$ can be viewed as a subspace of the product space $\text{Aut}(X, \mu)^\Gamma$, it inherits the product of the weak topology which we also call the *weak topology* on $A(\Gamma, X, \mu)$. Note that $\text{Aut}(X, \mu)$ acts continuously via conjugation on $A(\Gamma, X, \mu)$: Given $S \in \text{Aut}(X, \mu)$ and $a \in A(\Gamma, X, \mu)$, we let $S \cdot a = SaS^{-1}$ be the action of Γ for which $\gamma^{SaS^{-1}} = S\gamma^aS^{-1}$, $\forall \gamma \in \Gamma$. Then $a, b \in A(\Gamma, X, \mu)$ are conjugate iff they are isomorphic, in symbols $a \cong b$.

Motivated by the concept of weak containment of unitary representations, we can consider an analogous concept of weak containment of actions (see Kechris [Ke10], Section 10, (C)). We say, for $a \in A(\Gamma, X, \mu), b \in A(\Gamma, Y, \nu)$, that a is *weakly contained* in b , in symbols

$$a \prec b,$$

if for any Borel sets $A_1, \dots, A_n \subseteq X, \gamma_1 \dots \gamma_m \in \Gamma$ and $\epsilon > 0$, there are Borel

sets $B_1, \dots, B_n \subseteq X$ such that

$$|\mu(\gamma_i^a(A_j) \cap A_k) - \nu(\gamma_i^b(B_j) \cap B_k)| < \epsilon,$$

$\forall i \leq m, \forall j, k \leq n$.

Alternatively one can see that the following are equivalent for any $a \in A(\Gamma, X, \mu), b \in A(\Gamma, Y, \nu)$:

- (i) $a \prec b$,
- (ii) a is in the weak closure of $\{c \in A(\Gamma, X, \mu) : c \cong b\}$.

If $(Y, \nu) = (X, \mu)$, these are also clearly equivalent to:

- (iii) a is in the weak closure of the conjugacy class of b .

(See Kechris [Ke10], Section 10 (C).)

We say that a, b are *weakly equivalent* if $a \prec b$ and $b \prec a$. It is easy to see that \prec is a partial (pre-)order on $A(\Gamma, X, \mu)$. It was shown independently by Hjorth (unpublished) and Glasner-Thouvenot-Weiss [GTW] that \prec has a largest element (unique up to weak equivalence), which we denote by a_∞ . This means that a_∞ has dense conjugacy class in $A(\Gamma, X, \mu)$. The action a_∞ is obtained by taking the (diagonal) product of a countable dense set $\{a_n\}$ of actions in $A(\Gamma, X, \mu)$, but this hardly gives a concrete representation of a_∞ . It is thus of interest to “compute” explicitly such maximum actions for various groups Γ . We do that below for the free groups \mathbb{F}_n . An additional motivation for this goal is the connection with the theory of costs, which will be described below.

If Γ is an infinite, residually finite group, then Γ is a dense subgroup of its profinite completion $\hat{\Gamma}$ which is defined as the inverse limit of the groups $\Gamma/N, N \triangleleft \Gamma, [\Gamma : N] < \infty$. Clearly $\hat{\Gamma}$ is a compact Polish group and we let $\hat{\eta}_\Gamma$ be its (normalized) Haar measure, which is clearly non-atomic. The (left-) translation action of Γ on $\hat{\Gamma}$ is a measure preserving action of Γ on $(\hat{\Gamma}, \hat{\eta}_\Gamma)$, which we denote by p_Γ . We now have:

Theorem 1 *Let $1 \leq n \leq \infty$ and let \mathbb{F}_n be the free group with n generators. Then the action $p_{\mathbb{F}_n}$ is maximum in the order \prec of weak containment of measure preserving actions of \mathbb{F}_n .*

We now discuss an application to the theory of costs by showing how Theorem 1 together with the result in Abért-Nikolov [AN] gives a new method

for showing that the cost of any free, measure preserving action of \mathbb{F}_n ($1 \leq n < \infty$) is equal to n , a result originally proved by Gaboriau [G]. (Recall that an action $a \in A(\Gamma, X, \mu)$ is *free* if $\gamma^a(x) \neq x, \forall \gamma \neq 1, \mu$ -a.e. (x) .)

Denote by $\text{FR}(\Gamma, X, \mu)$ the subspace of $A(\Gamma, X, \mu)$ consisting of the free actions. For any $a \in A(\Gamma, X, \mu)$, we write

$$xE_a y \Leftrightarrow \exists \gamma (\gamma^a(x) = y),$$

for the equivalence relation induced by a . We denote by $C_\mu(E_a) \equiv C(E_a)$ the cost of the equivalence relation E_a , see Gaboriau [G]. Finally, we put

$$C_\mu(a) \equiv C(a) = C(E_a).$$

Then $C: A(\Gamma, X, \mu) \rightarrow [0, \infty]$. We now have:

Theorem 2 (Kechris [Ke10], 10.13) *If Γ is infinite and finitely generated, then $C|_{\text{FR}(\Gamma, X, \mu)}$ is upper semicontinuous.*

Corollary 3 *For such Γ , if $a, b \in \text{FR}(\Gamma, X, \mu)$, then*

$$a \prec b \Rightarrow C(b) \leq C(a).$$

The *cost* of the group Γ is defined by

$$C(\Gamma) = \inf\{C(a): a \in \text{FR}(\Gamma, X, \mu)\}.$$

It thus follows that if a_∞ is maximum in the order \prec , then

$$C(\Gamma) = C(a_\infty),$$

which gives an additional reason for “computing” explicitly a_∞ . For the case of the free groups, we now have:

Corollary 4 *For $1 \leq n < \infty, C(\mathbb{F}_n) = C(p_{\mathbb{F}_n})$.*

Now Abért-Nikolov [AN] had already found an explicit calculation of the cost of p_Γ , which is as follows.

Theorem 5 (Abért-Nikolov [AN]) *Let Γ be an infinite, finitely generated, residually finite group and let $\text{RG}(\Gamma)$ be its (absolute) rank gradient:*

$$\text{RG}(\Gamma) = \inf_{[\Gamma: H] < \infty} \frac{d(H) - 1}{[\Gamma: H]},$$

where $d(H)$, the rank of H , is the smallest number of generators of H . Then

$$C(p_\Gamma) = \text{RG}(\Gamma) + 1.$$

Now in the case $\Gamma = \mathbb{F}_n$ ($1 \leq n < \infty$), it is a standard fact in group theory that for any $H \leq \Gamma$ with $[\Gamma : H] < \infty$, we have

$$\frac{d(H) - 1}{[\Gamma : H]} = n - 1,$$

so

$$C(\mathbb{F}_n) = C(p_{\mathbb{F}_n}) = n.$$

Also, since clearly for every $a \in \text{FR}(\mathbb{F}_n, X, \mu)$, we have $C(a) \leq n$, it follows that $C(a) = n$, for *every* $a \in \text{FR}(\mathbb{F}_n, X, \mu)$, i.e., \mathbb{F}_n has *fixed price*.

(B) It is now of some interest to investigate for which residually finite groups Γ the analog of Theorem 1 goes through, i.e., p_Γ is maximum in the order of weak containment. It is also of some interest to investigate the (a priori) weaker condition that the (diagonal) product action $i_\Gamma \times p_\Gamma$, where i_Γ is the trivial action of Γ (on a non-atomic standard measure space) is maximum in \prec . Note that $p_\Gamma \prec i_\Gamma \times p_\Gamma$ and $i_\Gamma \times p_\Gamma$ is the direct sum of continuum many copies of p_Γ . Moreover $C(p_\Gamma) = C(i_\Gamma \times p_\Gamma)$ (see Kechris-Miller [KM], 18.14). We introduce the following terminology (whose choice will be explained below).

An infinite, residually finite group Γ has the *property MD* if $i_\Gamma \times p_\Gamma$ is maximum in the order \prec of weak containment and has the *property EMD* if p_Γ is maximum.

It turns out that each one of these properties is equivalent to an appropriate density condition in the space $A(\Gamma, X, \mu)$.

An action $a \in A(\Gamma, X, \mu)$ is called *finitely modular* or *profinite* if there is a decreasing sequence of finite Borel partitions $\{X\} = \mathcal{P}_0 \geq \mathcal{P}_1 \geq \dots$ such that each \mathcal{P}_n is Γ -invariant and $\{\mathcal{P}_n\}$ separates points. Up to isomorphism, these can be also equivalently described as the actions of the following form: Given an infinite, finite splitting, rooted tree T with no finite branches, and an action of Γ by automorphisms on T , let ∂T be the boundary of T and consider the induced action of Γ on ∂T and a (non-atomic) invariant measure μ on ∂T for this action. Then the finitely modular actions, up to isomorphism, are exactly the actions of Γ on $(\partial T, \mu)$ as above. The finitely modular actions which are ergodic correspond exactly to the actions of Γ on trees which are *level transitive*, i.e., act transitively on each level of the tree (see §1 below for details).

Finally, let us call $a \in A(\Gamma, X, \mu)$ *finite* iff it factors through an action of a finite group, i.e., there is a finite group Δ , an action $b \in A(\Delta, X, \mu)$, and a surjective homomorphism $\pi: \Gamma \rightarrow \Delta$, such that $\gamma^a = \pi(\gamma)^b, \forall \gamma \in \Gamma$.

We now have

Proposition 6 *The following are equivalent for each infinite, residually finite group Γ :*

- i) Γ satisfies EMD,*
- ii) The ergodic, finitely modular actions are dense in $A(\Gamma, X, \mu)$.*

Also the following are equivalent:

- a) Γ satisfies MD,*
- b) The finitely modular actions are dense in $A(\Gamma, X, \mu)$,*
- c) The finite actions are dense in $A(\Gamma, X, \mu)$.*

Clearly

$$\text{EMD} \Rightarrow \text{MD}$$

and any group that satisfies EMD cannot have *property* (τ) (which states that the trivial representation 1_Γ is weakly contained in the Koopman representation associated to p_Γ , restricted to the orthogonal of the constants). Moreover, from a recent result of Abért-Weiss [AW], it follows that when Γ does not have property (τ) , then Γ satisfies EMD iff it satisfies MD.

It is not known if EMD and MD are equivalent but by the above remarks this question is equivalent to the problem of whether the property MD and (τ) are incompatible. It is also unknown whether MD and property (T) are incompatible.

The property MD is an ergodic theoretic analog of the property FD discussed in Lubotzky-Shalom [LS] (see also Lubotzky-Zuk [LZ]). This asserts that the finite unitary representations of Γ on an infinite-dimensional separable Hilbert space \mathcal{H} are dense in the space $\text{Rep}(\Gamma, \mathcal{H})$ of unitary representations of Γ in \mathcal{H} . Here $\text{Rep}(\Gamma, \mathcal{H}) \subseteq U(\mathcal{H})^\Gamma$, where $U(\mathcal{H})$ is the unitary group of \mathcal{H} , is equipped with the product topology, with $U(\mathcal{H})$ having the weak topology. (A *finite representation* is again one that factors through a representation of a finite group.) One can see that

$$\text{MD} \Rightarrow \text{FD}$$

but the converse is not known. It is also not known if FD is incompatible with property (T) or property (τ) .

The extent of the class of infinite, residually finite groups that satisfy EMD or MD is rather unclear. We have seen that the free groups \mathbb{F}_n satisfy EMD and so do all amenable groups. Moreover the property MD is stable under going to subgroups (but this is not clear for EMD) and both EMD, MD are stable under going to supergroups of finite index. Thus, in particular, $\mathrm{SL}_2(\mathbb{Z})$ and $A * B$, for A, B finite non-trivial groups, have property EMD. On the other hand Lubotzky-Shalom [LS] and Lubotzky-Zuk [LZ] discuss various examples, including $\mathrm{SL}_n(\mathbb{Z})$, for $n \geq 3$, that fail to have property FD and thus also fail to have property MD.

(C) It turns out that there is also another description of the maximum, under weak containment, action of a free group.

Theorem 7 *Let $\Gamma = \mathbb{F}_n$ ($1 \leq n \leq \infty$). Then there is a subgroup $H \leq \mathbb{F}_n$ of infinite index such that the generalized shift action $s_{\Gamma, \Gamma/H}$ of Γ on $2^{\Gamma/H}$ is maximum in the order \prec of weak containment.*

A similar result holds for representations: For $\Gamma = \mathbb{F}_n$, there is an infinite index subgroup $H \leq \Gamma$ such that the quasi-regular representation $\lambda_{\Gamma/H}$ of Γ on $\ell^2(\Gamma/H)$ weakly contains *every* unitary representation of Γ .

It follows that for such H the action of $\Gamma = \mathbb{F}_n$ on $I = \Gamma/H$ is transitive, faithful and amenable. (Amenability means that $1_\Gamma \prec \lambda_{\Gamma/H}$.) The existence of such actions was first proved by van Douwen and later other examples were found by Glasner-Monod and Grigorchuk-Nekrashevych (see Glasner-Monod [GM] for more details). The above result provides an alternative such construction, which has the additional property that $\pi \prec \lambda_{\Gamma/H}$ for *every* unitary representation of $\Gamma = \mathbb{F}_n$ (instead of just $1_\Gamma \prec \lambda_{\Gamma/H}$). Moreover, for $n \geq 2$, one can find such H for which in addition $\lambda_{\Gamma/H}$ is irreducible. The existence of irreducible representations of \mathbb{F}_n that weakly contain any unitary representation was originally proved in Yoshizawa [Y]. (For another proof see Kechris [Ke10], Appendix H, **(C)**.)

(D) This paper is organized as follows. Section 1 reviews various preliminaries and establishes general notation. In Section 2 we discuss actions on trees and in Section 3 we prove Theorem 1 (see Theorem 3.1) and discuss the application to costs. In Section 4 we discuss the properties EMD and MD and some related facts and questions. Section 5 contains the proof of Theorem 7 (see Theorem 5.1). In Section 6 we discuss various results concerning density and meagerness conditions for various sets of actions (both within the space of all actions and also within the space of all actions “included”

in the full group of an equivalence relation). Finally, there is an appendix reviewing the concept of co-induced action and some facts concerning this notion that are used in the paper.

Addendum. Lewis Bowen informed me that he has earlier studied a property of groups which he called the *periodic approximation property* (PA), expressed in terms of approximations of invariant measures for the shift actions of the group by periodic ones. It turns out that PA is equivalent to MD. Bowen had shown, among other things, that free groups have property PA and that $PA \Rightarrow FD$ (and also raised the question of whether $PA \Leftrightarrow FD$). The result that the free groups have property PA is included in L. Bowen, Periodicity and packings of the hyperbolic plane, *Geom. Dedicata*, **102** (2003), 213–236. See also L. Bowen, Free groups in lattices, *arXiv:0802.0185v5*. Other results concerning PA are included in the preprints: L. Bowen, Notes on the periodic approximation property, December 18, 2007 and Section 5 of L. Bowen, Sphere packings in hyperbolic space: periodicity and continuity, December, 2003. The last paper contains a proof of a special case of the Conjecture 4.14 below (which includes surface groups and lattices in $SO(3,1)$) that he can generalize to obtain all of 4.14.

Acknowledgements. I would like to thank M. Abért, L. Bowen, D. Gaboriau, A. Ioana, N. Monod, Y. Shalom and T. Tsankov for many useful discussions or comments concerning this paper and G. Hjorth for allowing me to include 6.3 below. The research of the author was partially supported by NSF Grant DMS-0455285.

1 Preliminaries

(A) We work throughout in standard Borel spaces X , i.e., sets equipped with a σ -algebra of subsets (called *Borel sets*), which is isomorphic to the σ -algebra of Borel sets on a *Polish* (separable, completely metrizable) space. A (*Borel*) *measure* μ on such a space X is a measure on the σ -algebra of Borel sets. It is a *probability measure* if $\mu(X) = 1$. *Unless otherwise indicated, we consider only probability measures in the sequel.*

If $a: G \times S \rightarrow S$ is an action of a group G on a set S , we let

$$g^a(s) = a(g, s),$$

for $g \in G, s \in S$. We also put

$$g \cdot s = a(g, s),$$

if there is no danger of confusion. Given a countable (discrete) group Γ , a Borel action of Γ on (X, μ) is *measure preserving* if $\mu(\gamma \cdot A) = \mu(A)$, for every $\gamma \in \Gamma$ and Borel $A \subseteq X$.

We let

$$A(\Gamma, X, \mu)$$

be the space of all measure preserving actions of Γ on (X, μ) , where the actions a, b are identified if $\gamma^a = \gamma^b, \mu$ -a.e., $\forall \gamma \in \Gamma$. An action $a \in A(\Gamma, X, \mu)$ is *free* if $\forall \gamma \neq 1 (\gamma \cdot x \neq x, \mu$ -a.e.). We denote by

$$\text{FR}(\Gamma, X, \mu)$$

the set of free actions. An action $a \in A(\Gamma, X, \mu)$ is *ergodic* if there are no non-trivial invariant sets. Again we denote by

$$\text{ERG}(\Gamma, X, \mu)$$

the set of such actions.

If $a_i \in A(\Gamma, X_i, \mu_i), i \in I$, are actions, where I is countable, the (*diagonal*) *product action*

$$\prod_{i \in I} a_i \in A(\Gamma, X^I, \mu^I)$$

is defined by

$$\gamma \cdot (x_i)_{i \in I} = (\gamma \cdot x_i)_{i \in I}.$$

A particular case is the product

$$a \times b$$

of two actions.

(B) We use the following notation for some particular actions:

(a) If $\Gamma \leq \Delta$ are countable groups, $a_{\Delta/\Gamma}$ is the canonical action of Δ on Δ/Γ (= the set of left cosets of Γ in Δ):

$$\delta \cdot \delta' \Gamma = \delta \delta' \Gamma.$$

For any set X , $s_{\Gamma, X}$ is the *shift* action of Γ on X^Γ given by

$$\gamma \cdot f(\delta) = f(\gamma^{-1}\delta).$$

When $X = \{0, 1\} = 2$, we write

$$s_\Gamma \equiv s_{\Gamma, 2}.$$

If $\Gamma \leq \Delta$, we write $s_{\Delta, \Delta/\Gamma, X}$ for the *generalized shift action* of Δ on $X^{\Delta/\Gamma}$ given by

$$\delta \cdot f(\delta'\Gamma) = f(\delta^{-1}\delta'\Gamma).$$

Again when $X = 2$, we write

$$s_{\Delta, \Delta/\Gamma} \equiv s_{\Delta, \Delta/\Gamma, 2}.$$

More generally, if Γ acts on a countable set I , we define the *generalized shift action* of $s_{\Gamma, I, X}$ on X^I by

$$\delta \cdot f(i) = f(\delta^{-1} \cdot i),$$

and again we let $s_{\Gamma, I} \equiv s_{\Gamma, I, 2}$.

(C) We denote by

$$\text{Aut}(X, \mu)$$

the group of measure preserving automorphisms of (X, μ) , where two such automorphisms S, T are identified if $S = T$, μ -a.e. Thus $A(\Gamma, X, \mu)$ can be viewed as the set of all homomorphisms of Γ into $\text{Aut}(X, \mu)$.

The group $\text{Aut}(X, \mu)$ has two canonical topologies, the weak and the uniform. The *weak topology* is the topology generated by the maps

$$T \mapsto T(A),$$

where $T \in \text{Aut}(X, \mu)$, $A \in \text{MALG}(X, \mu)$ = the measure algebra of (X, μ) , and $\text{MALG}(X, \mu)$ is equipped with the metric $d_\mu(A, B) = \mu(A \Delta B)$ (and the corresponding topology). The group $\text{Aut}(X, \mu)$ is a Polish group in the weak topology.

The *uniform topology* on $\text{Aut}(X, \mu)$ is the one induced by the following complete metric

$$\delta_\mu(S, T) = \mu(\{x: S(x) \neq T(x)\}).$$

It is not separable, if μ is non-atomic.

We equip the product space $\text{Aut}(X, \mu)^\Gamma$ with the product topology (in either one of the two topologies on $\text{Aut}(X, \mu)$). Then $A(\Gamma, X, \mu) \subseteq \text{Aut}(X, \mu)^\Gamma$ is a closed subspace (in either one of these topologies), and we equip it with the relative topology called, respectively, the *weak and the uniform topology* on $A(\Gamma, X, \mu)$. The weak topology is Polish and the uniform topology is completely metrizable.

(D) Two actions $a \in A(\Gamma, X, \mu), b \in A(\Gamma, Y, \nu)$ are *isomorphic*, in symbols

$$a \cong b,$$

if there is an isomorphism $\varphi: (X, \mu) \rightarrow (Y, \nu)$ such the $\varphi\gamma^a\varphi^{-1} = \gamma^b, \forall \gamma \in \Gamma$. If $(X, \mu) = (Y, \nu)$ and we consider the action of $\text{Aut}(X, \mu)$ on $A(\Gamma, X, \mu)$ by conjugation, $S \cdot a \equiv SaS^{-1}$, where

$$\gamma^{SaS^{-1}} = S\gamma^aS^{-1}, \forall \gamma \in \Gamma,$$

then $a, b \in A(\Gamma, X, \mu)$ are isomorphic iff they are conjugate.

If $a \in A(\Gamma, X, \mu), b \in A(\Gamma, Y, \nu)$, then b is a *factor* of a , in symbols

$$b \sqsubseteq a,$$

if there is a Borel map $\varphi: X \rightarrow Y$ such that $\varphi_*\mu = \nu$ (i.e., $\nu(A) = \mu(\varphi^{-1}(A))$) and $\varphi(\gamma^a(x)) = \gamma^b(\varphi(x)), \forall \gamma \in \Gamma, \mu$ -a.e. For example, $a \sqsubseteq a \times b$.

Occasionally we need to talk about continuous actions of Γ on compact spaces. In this situation, we say that a continuous action b of Γ on L is a *factor* of a continuous action a of Γ on K if there is a continuous map $\varphi: K \rightarrow L$ such that $\varphi(\gamma^a(x)) = \gamma^b(\varphi(x))$.

(E) For $a \in A(\Gamma, X, \mu), b \in A(\Gamma, Y, \nu)$ we say that a is *weakly contained* in b , in symbols

$$a \prec b,$$

if for any Borel sets $A_1, \dots, A_n \subseteq X$, any $F \subseteq \Gamma$ finite, and any $\epsilon > 0$, there are $B_1, \dots, B_n \subseteq Y$ such that

$$|\mu(\gamma^a(A_i) \cap A_j) - \nu(\gamma^b(B_i) \cap B_j)| < \epsilon,$$

$\forall \gamma \in F, \forall i, j \leq n$. See Kechris [Ke10], Section 10, **(C)** for more information about this concept. When $(X, \mu), (Y, \nu)$ are non-atomic, then $a \prec b$ iff a is

in the weak closure of the set of isomorphic copies of b in $A(\Gamma, X, \mu)$. In particular, if also $(X, \mu) = (Y, \nu)$, then $a \prec b$ iff a is in the weak closure of the conjugacy class of b . It is easy to verify that

$$a \sqsubseteq b \Rightarrow a \prec b.$$

It is also easy to see that \prec is a partial (pre-)order. The associated equivalence relation

$$a \sim b \Leftrightarrow a \prec b \ \& \ b \prec a,$$

is called *weak equivalence*.

(F) Given a standard Borel space X , an equivalence relation E on X is called *countable* if every equivalence class $[x]_E, x \in X$, is countable. By a result of Feldman-Moore [FM], for every such E there is a Borel action a of Γ on X such that

$$E = E_a,$$

where E_a is the equivalence relation induced by a ,

$$xE_a y \Leftrightarrow \exists \gamma \in \Gamma (\gamma \cdot x = y).$$

If now μ is a measure on X , then E is *measure preserving* if $E = E_a$, for some action $a \in A(\Gamma, X, \mu)$ (this notion is independent of the action that induces E).

For a countable, measure preserving E on (X, μ) we let

$$[E] = \{T \in \text{Aut}(X, \mu) : T(x)Ex, \mu\text{-a.e.}\}$$

be the *full group* of E . It is a closed, *separable* subgroup of $\text{Aut}(X, \mu)$ in the uniform topology. Letting below φ vary over all *partial*, measure preserving bijections $\varphi : A \rightarrow B$ between Borel subsets of X , we let

$$[[E]] = \{\varphi : \varphi(x)Ex, \mu\text{-a.e.}\}$$

Thus $[E] = [[E]] \cap \text{Aut}(X, \mu)$.

(G) For any countable, measure preserving E on (X, μ) we denote by

$$C_\mu(E) \equiv C(E)$$

the *cost* of E (see Gaboriau [G] or Kechris-Miller [KM]). If $a \in A(\Gamma, X, \mu)$, let

$$C_\mu(a) \equiv C(a) = C(E_a).$$

The *cost* of a group Γ is defined by

$$C(\Gamma) = \inf\{C_\mu(a) : a \in \text{FR}(\Gamma, X, \mu), \mu \text{ non-atomic}\}.$$

We say that Γ has *fixed price* if $C_\mu(a) = C(\Gamma), \forall a \in \text{FR}(\Gamma, X, \mu), \mu$ non-atomic.

(H) For each (separable) Hilbert space \mathcal{H} , we denote by $\text{Rep}(\Gamma, \mathcal{H})$ the space of unitary representations of Γ on \mathcal{H} , i.e., homomorphisms from Γ into $U(\mathcal{H})$, the unitary group of \mathcal{H} . We equip $U(\mathcal{H})$ with the weak (equivalently the strong) topology and $\text{Rep}(\Gamma, \mathcal{H})$ with the product topology (viewing $\text{Rep}(\Gamma, \mathcal{H})$ as a closed subspace of $U(\mathcal{H})^\Gamma$). As usual, if $\pi \in \text{Rep}(\Gamma, \mathcal{H}_1), \rho \in \text{Rep}(\Gamma, \mathcal{H}_2)$, we let

$$\pi \prec \rho,$$

if π is *weakly contained* in ρ . We also write

$$\pi \leq \rho$$

if π is (isomorphic to) a subrepresentation of ρ , and let

$$\pi \cong \rho$$

if π, ρ are isomorphic.

If $\Gamma \leq \Delta$, then we denote by $\lambda_{\Delta/\Gamma}$ the *quasi-regular representation* of Δ on $\ell^2(\Delta/\Gamma)$ given by

$$\lambda_{\Delta/\Gamma}(\delta)(f)(\delta'\Gamma) = f(\delta^{-1}\delta'\Gamma).$$

Finally 1_Γ is the trivial 1-dimensional representation of Γ .

For each $a \in A(\Gamma, X, \mu)$, we let κ^a be the *Koopman (unitary) representation* associated to a , which is the unitary representation of Γ on $L^2(X, \mu)$ given by

$$\kappa^a(\gamma)(f)(x) = f(\gamma^{-1} \cdot x).$$

We also let κ_0^a be the restriction of κ^a to the orthogonal of the constant functions in $L^2(X, \mu)$. It can be shown that if $a, b \in A(\Gamma, X, \mu), \mu$ non-atomic, then

$$a \prec b \Rightarrow \kappa^a \prec \kappa^b, \kappa_0^a \prec \kappa_0^b,$$

(see [Ke10], Section 10, **(C)**).

(I) Conventions

(a) Throughout the paper, when we work in a measure theoretic context, we neglect null sets, unless there is a danger of confusion. In particular, we do not distinguish between saying that a certain property of $x \in X$, where (X, μ) is a measure space, is true for all x or for μ -almost all x .

(b) The measure spaces (X, μ) in the sequel will be always assumed to be non-atomic, unless otherwise explicitly stated or obviously understood from the context (e.g., when X is finite).

2 Actions on trees

For our purposes in this paper, a *tree* is an acyclic, connected (simple, undirected) rooted graph $T = (V, E, v_0)$ with vertex set V , edge set E and root $v_0 \in V$. For every $v \in V$ there is a unique path $v_0, v_1, \dots, v_n = v$ of distinct vertices with $v_{i+1}Ev_i$ from the root v_0 to v . The *children* of v are all vertices adjacent to v different from v_{n-1} and the *parent* of v is v_{n-1} . The *n th level* ($n \geq 0$) of T , in symbols T_n , consists of all $v \in T$ for which the unique path from v_0 to v has $n + 1$ vertices as above. If $v \in T_n$, we write $|v| = n$. (Thus $T_0 = \{v_0\}$ and $|v_0| = 0$.)

We say that T is *finite splitting* if every $v \in V$ has finitely many children. A *terminal node* of T is a vertex v with no children.

From now on we will assume (unless otherwise explicitly indicated) that all trees are finitely splitting and have no terminal nodes. For such a tree T , the *boundary* ∂T of T consists of all infinite sequences (v_0, v_1, v_2, \dots) of distinct vertices with v_{n+1} adjacent to v_n . It is clearly nonempty and compact metrizable in the topology generated by the basic open sets

$$N_v = \{(v_0, v_1, \dots) \in \partial T : v = v_n\},$$

for $v \in T_n, n \geq 0$. (Thus $N_{v_0} = \partial T$.) Note that these are actually clopen, so ∂T is 0-dimensional.

We will consider actions of infinite, countable (discrete) groups Γ on trees. An *action of Γ on T* is an action of Γ by automorphisms of T . In particular, Γ fixes v_0 and acts on each T_n . Moreover, if $v \in T_n, w \in T_{n+1}$ and w is a child of v , then $\gamma \cdot v \in T_n, \gamma \cdot w \in T_{n+1}$ and $\gamma \cdot w$ is a child of $\gamma \cdot v$. The action of Γ induces an action of Γ on ∂T via

$$\gamma \cdot (v_0, v_1, \dots) = (\gamma \cdot v_0, \gamma \cdot v_1, \dots).$$

This is clearly an action by homeomorphisms on ∂T . Given a probability measure μ on ∂T , let $\mu(v) = \mu(N_v)$. Then μ is invariant under the Γ -action iff $\mu(v) = \mu(\gamma \cdot v), \forall \gamma \in \Gamma$.

The action of Γ on T is *level transitive* if Γ acts transitively on each T_n . In this case the associated action of Γ on ∂T is uniquely ergodic, i.e., has a unique invariant (thus necessarily ergodic) probability measure μ_T defined by

$$\mu_T(v) = \frac{1}{\text{card}(T_n)},$$

for $v \in T_n$. We will always consider this measure when we study the Γ -action on ∂T . Finally, a level transitive action is minimal, i.e., every orbit in ∂T is dense.

We now define the *orbit tree* $O(T)$ associated to the Γ -action on T . The n th level of $O(T)$ consists of all the orbits of Γ on T_n . If o is an n th level orbit and o' an $(n+1)$ th level orbit, then o' is a child of o if for any $v' \in o'$, if v is the parent of v' , then $v \in o$.

If $e \in \partial(O(T))$, say $e = (o_0 = \{v_0\}, o_1, o_2, \dots)$, then let T_e be the subtree of T determined by

$$v \in T_e \Leftrightarrow v \in o_n,$$

for $v \in T_n, n \geq 0$. Clearly T_e is Γ -invariant and the action of Γ on T_e is level transitive. Put $X_e = \partial T_e$. Then $\{X_e\}_{e \in \partial(O(T))}$ is a decomposition of ∂T into closed Γ -invariant sets on which Γ acts minimally and with a unique invariant, ergodic probability measure μ_e .

Proposition 2.1 *Every Γ -invariant, probability measure on ∂T is of the form μ_e .*

Proof. Fix such a measure ν . Now $\partial T = \bigsqcup_{i=1}^k X_{o_i}$, is a decomposition into Γ -invariant sets, where o_1, \dots, o_k are the Γ -orbits on T_1 , and we let $X_o = \bigcup_{v \in o} N_v$. Thus there is (a unique) o_i , say o_1 , such that $\nu(X_{o_1}) = 1$. Next X_{o_1} decomposes into the Γ -invariant sets $X_{o'_1}, \dots, X_{o'_\ell}$, where o'_1, \dots, o'_ℓ are the Γ -orbits of T_2 which are children of o_1 . Again there is unique o' , say o'_1 , with $\nu(X_{o'_1}) = 1$. Proceed this way to define $e = (o_0 = \{v_0\}, o_1, o'_1, o''_1, \dots) \in \partial(O(T))$. Then $X_e = X_{o_1} \cap X_{o'_1} \cap X_{o''_1} \cap \dots$, so $\nu(X_e) = 1$, and as ν is Γ -invariant, $\nu = \mu_e$. \dashv

Thus

$$\mathcal{E} = \{\mu_e : e \in \partial(O(T))\}$$

is the space of all Γ -invariant, ergodic probability measures on ∂T . For $x \in \partial T$, put

$$\mu_x = \mu_e, \text{ where } x \in X_e.$$

Proposition 2.2 *If μ is a Γ -invariant probability measure on ∂T , then*

$$\mu = \int \mu_x d\mu(x).$$

Proof. It is enough to show for any $v \in V$, that $\mu(N_v) = \int \mu_x(N_v) d\mu(x)$. Let $v \in T_n$ and let o be the Γ -orbit of v . Then $\mu(N_v) = \frac{\mu(X_o)}{\text{card}(o)}$. Clearly $\mu_x(N_v) = 0$, unless $x \in X_o$, so

$$\begin{aligned} \int \mu_x(N_v) d\mu(x) &= \int_{X_o} \mu_x(N_v) d\mu(x) \\ &= \frac{\mu(X_o)}{\text{card}(o)} = \mu(N_v) \end{aligned}$$

□

Thus the map $\pi(x) = \mu_x$ has the following properties:

- (i) It is a Borel surjection of ∂T onto the standard Borel space \mathcal{E} .
- (ii) It is Γ -invariant.
- (iii) For any $\rho \in \mathcal{E}$, $\pi^{-1}(\{\rho\})$ is Γ -invariant, $\rho(\pi^{-1}(\{\rho\})) = 1$ and the action of Γ on $\pi^{-1}(\{\rho\})$ is uniquely ergodic as witnessed by ρ .
- (iv) If μ is a Γ -invariant probability measure on ∂T , then $\mu = \int \pi(x) d\mu(x)$.

Thus π is the Ergodic Decomposition of the Γ -action on ∂T (in the strong sense of Farrell, Varadarajan—see, e.g., Kechris-Miller [KM], 3.3, where the context is that of countable equivalence relations but holds equally well in the context of actions by countable groups).

Now consider a level transitive action of Γ on T . Fix $x \in \partial T$, $x = (v_0, v_1, v_2, \dots)$ and let Γ_n be the stabilizer of v_n . Then $\Gamma_0 = \Gamma \geq \Gamma_1 \geq \Gamma_2 \geq \dots$ and $[\Gamma : \Gamma_n] < \infty$ (in fact $[\Gamma : \Gamma_n] = \text{card}(T_n)$). We call such a sequence $\{\Gamma_n\}$ a *chain* in Γ . Conversely, for any chain $\{\Gamma_n\}$ we can define a tree $T(\Gamma, \{\Gamma_n\})$, where the n th level $T_n = T_n(\Gamma, \{\Gamma_n\})$ consists of the cosets $g\Gamma_n$ of Γ_n , i.e., $T_n = \Gamma/\Gamma_n$, and $h\Gamma_{n+1}$ is a child of $g\Gamma_n$ if $h\Gamma_{n+1} \subseteq g\Gamma_n$.

The group Γ acts on $T(\Gamma, \{\Gamma_n\})$ in the obvious way: $\gamma \cdot g\Gamma_n = \gamma g\Gamma_n$. If $T_n = T(\Gamma, \{\Gamma_n\})$, clearly $\Gamma_n \in T_n$ and the stabilizer of T_n in the Γ -action is equal to Γ_n . It is clear that in the particular case that Γ_n is the stabilizer of v_n , in the preceding notation, the map $g\Gamma_n \mapsto g \cdot v_n$ is an isomorphism of the Γ -action on T and the Γ -action on $T(\Gamma, \{\Gamma_n\})$. Thus all level transitive T -action on trees are actions of Γ on trees of the form $T(\Gamma, \{\Gamma_n\})$.

Now assume that Γ is a *residually finite* group, i.e., $\bigcap \{H \leq \Gamma : [\Gamma : H] < \infty\} = \{1\}$. Equivalently this means that there is a chain $\Gamma = \Gamma_0 \geq \Gamma_1 \geq \dots$, with $\bigcap_n \Gamma_n = \{1\}$. Such a chain can always be taken to be normal, i.e., $\Gamma_n \triangleleft \Gamma$, since any subgroup of finite index contains a normal one with the same property, i.e., the intersection of its conjugates. If $\{\Gamma_n\}$ is a normal chain with $\bigcap_n \Gamma_n = \{1\}$, then $\partial T(\Gamma, \{\Gamma_n\})$ is also a group with multiplication defined as follows: If $x = (x_0, x_1, \dots)$, $y = (y_0, y_1, \dots)$, where $x_n, y_n \in \Gamma/\Gamma_n$, then $xy = (x_0y_0, x_1y_1, \dots)$. This turns $\partial T(\Gamma, \{\Gamma_n\})$ into a compact, metrizable, 0-dimensional topological group. Next note that Γ can be identified with a dense subgroup of $\partial T(\Gamma, \{\Gamma_n\})$ by identifying $\gamma \in \Gamma$ with $(\Gamma, \gamma\Gamma_1, \gamma\Gamma_2, \dots)$. With this identification the action of Γ on $\partial T(\Gamma, \{\Gamma_n\})$ is simply the left-translation action of Γ on the group $\partial T(\Gamma, \{\Gamma_n\})$. In particular, it is free.

Consider now the special case of a normal chain $\Gamma = \Gamma_0 \geq \Gamma_1 \geq \dots$ which is *cofinal*, i.e., $\forall H \leq \Gamma ([\Gamma : H] < \infty \Rightarrow \exists n, H \geq \Gamma_n)$ (in particular $\bigcap_n \Gamma_n = \{1\}$). Then the group $\hat{\Gamma} = \partial T(\Gamma, \{\Gamma_n\})$ is called the *profinite completion* of Γ . It is independent of $\{\Gamma_n\}$ as it is isomorphic to the inverse limit given by $\varprojlim \{\Gamma/N : N \triangleleft \Gamma, [\Gamma : N] < \infty\}$. We denote the action of Γ on $\hat{\Gamma}$ by p_Γ . The canonical invariant measure for this action is of course the (normalized) Haar measure on $\hat{\Gamma}$. It is clearly non-atomic, since Γ is infinite.

Proposition 2.3 *For any chain $\Gamma = \Gamma_0 \geq \Gamma_1 \geq \dots$, the action of Γ on $\partial T(\Gamma, \{\Gamma_n\})$ is a factor of the action p_Γ , both in the topological and measure theoretic sense.*

Proof. Fix a normal cofinal chain $\Gamma = N_0 \geq N_1 \geq \dots$. We will find a surjective continuous map $\pi : \partial T(\Gamma, \{N_n\}) \rightarrow \partial T(\Gamma, \{\Gamma_n\})$ that preserves the action of Γ .

Let $k_1 < k_2 < \dots$ be such that $k_1 = (\text{least } k \text{ with } N_k \subseteq \Gamma_1)$ and $k_{n+1} = (\text{least } k > k_n, \text{ with } N_k \subseteq \Gamma_{n+1})$. Then $N_{k_n} \subseteq \Gamma_n$. Given $x \in \partial T(\Gamma, \{N_n\})$, $x = (\Gamma, g_1N_1, g_2N_2, \dots)$, let $\pi(x) = (\Gamma, g_{k_1}\Gamma_1, g_{k_2}\Gamma_2, \dots)$. Note that $g_{k_n}\Gamma_n$ is the unique Γ_n -coset containing the N_{k_n} -coset $g_{k_n}N_{k_n}$. Clearly π is continuous and preserves the Γ -actions.

We will next verify that it is surjective. Fix $y = (\Gamma, g_1\Gamma_1, g_2\Gamma_2, \dots) \in \partial T(\Gamma, \{\Gamma_n\})$. Call a sequence $s = (\Gamma, h_1N_{k_1}, \dots, h_nN_{k_n})$ *good* if $h_1N_{k_1} \supseteq \dots \supseteq h_nN_{k_n}$ and $h_iN_{k_i} \subseteq g_i\Gamma_i$, $1 \leq i \leq n$. Clearly every good s as above has only finitely many good extensions $(\Gamma, h_1N_{k_1}, \dots, h_nN_{k_n}, h_{n+1}N_{k_{n+1}})$. We now claim that for each n , there is some good $s = (\Gamma, h_1N_{k_1}, \dots, h_nN_{k_n})$. Indeed, consider $g_n\Gamma_n$, and let $h_n = g_n$, so that $h_nN_{k_n} \subseteq g_n\Gamma_n$. Then let $h_iN_{k_i}$ be the unique coset containing $h_nN_{k_n}$ ($1 \leq i < n$). We will show that $h_iN_{k_i} \subseteq g_i\Gamma_i$. We have $h_nN_{k_n} \subseteq h_iN_{k_i} \cap g_n\Gamma_n \subseteq h_iN_{k_i} \cap g_i\Gamma_i$, so $h_iN_{k_i} \cap g_i\Gamma_i \neq \emptyset$, thus $h_iN_{k_i} \subseteq g_i\Gamma_i$.

So, by König's Lemma, there is an infinite sequence $(\Gamma, h_1N_{k_1}, h_2N_{k_2}, \dots)$ such that for each i , $h_{i+1}N_{k_{i+1}} \subseteq h_iN_{k_i} \subseteq g_i\Gamma_i$. Let $x = (\Gamma, g_1N_1, g_2N_2, \dots)$ be the unique element of $\partial T(\Gamma, \{N_n\})$ such that $g_{k_i}N_{k_i} = h_iN_{k_i}$, $i = 1, 2, \dots$. Clearly then $\pi(x) = y$.

If μ is the unique invariant probability measure for p_Γ , then $\pi_*\mu$ is invariant for the Γ -action on $\partial T(\Gamma, \{\Gamma_n\})$, so it is equal to the unique Γ -invariant probability measure on $\partial T(\Gamma, \{\Gamma_n\})$. Thus π preserves the measures as well. \dashv

Thus for any level transitive action of Γ on a tree T , the corresponding action a of Γ on ∂T is a factor of p_Γ . In particular, a is weakly contained in p_Γ , $a \prec p_\Gamma$, i.e., p_Γ is the maximum, in the sense of weak containment, level transitive action of Γ on the boundary of a tree.

A *finitely modular action* or *profinite action* of Γ on a standard Borel space X is a Borel action for which there is a decreasing sequence of finite Borel partitions $\{X\} = \mathcal{P}_0 \geq \mathcal{P}_1 \geq \dots$ (i.e., \mathcal{P}_{n+1} refines \mathcal{P}_n) such that each \mathcal{P}_n is Γ -invariant (setwise) and $\{\mathcal{P}_n\}$ separate points. (See Hjorth [Hj02] or Kechris [Ke07]). If moreover for any $A_i \in \mathcal{P}_i$ with $A_0 \supseteq A_1 \supseteq \dots$, we have that $\bigcap_n A_n$ is a singleton, we call this a *special modular action* (thus special modular \Rightarrow finitely modular).

If Γ is an action on a tree T and $\mathcal{P}_n = \{N_v : v \in T_n\}$, clearly $\{\mathcal{P}_n\}$ shows that the action of Γ on ∂T is a special modular action. Conversely, if we have a special modular action of Γ on X with witness $\{\mathcal{P}_n\}$, consider the tree $T_{\{\mathcal{P}_n\}}$ whose n th level is equal to \mathcal{P}_n and $B \in \mathcal{P}_{n+1}$ is a child of $A \in \mathcal{P}_n$ if $B \subseteq A$. Then Γ acts on $T_{\{\mathcal{P}_n\}}$ and it is clear that the action of Γ on X is Borel isomorphic to the action of Γ on $\partial T_{\{\mathcal{P}_n\}}$ via the map $x \in X \mapsto (A_0 = X, A_1, A_2, \dots)$, where $x \in A_n \in \mathcal{P}_n$.

Even if a finitely modular action of Γ on X is not special, the map $x \in X \mapsto (A_0 = X, A_1, A_2, \dots)$ as above gives a Borel embedding π of the Γ -

action on X into the Γ -action on $\partial T_{\{\mathcal{P}_n\}}$ and thus if μ is a Γ -invariant measure on X , then $\nu = \pi_*\mu$ is a Γ -invariant measure on $\partial T_{\{\mathcal{P}_n\}}$ and the Γ -action on (X, μ) is (measure theoretically) isomorphic to the Γ -action on $(\partial T_{\{\mathcal{P}_n\}}, \nu)$. In other words, up to isomorphism, the measure preserving finitely modular actions are the same as the measure preserving actions induced on the boundaries of trees. The ergodic, finitely modular actions correspond to level transitive actions.

Consider now a sequence of actions of Γ on trees T^1, T^2, \dots , the corresponding actions of Γ on $\partial T^1, \partial T^2, \dots$ and the product action of Γ on $\prod_{m=1}^{\infty} \partial T^m$:

$$\gamma \cdot (x^1, x^2, \dots) = (\gamma \cdot x^1, \gamma \cdot x^2, \dots).$$

Proposition 2.4 *Given actions of Γ on trees T^1, T^2, \dots , there is a tree T and an action of Γ on T such that the Γ -action on ∂T is Borel isomorphic to the product action of Γ on $\prod_{m=1}^{\infty} \partial T^m$.*

Proof. It is enough to show that the action of Γ on $\prod_{m=1}^{\infty} \partial T^m$ is a special modular action. Let \mathcal{P}_n be the partition of $\prod_{m=1}^{\infty} \partial T^m$ given by the clopen sets

$$R_{v_1, \dots, v_n} = \{(x_m) \in \prod_{m=1}^{\infty} \partial T^m : \\ x_m(n) = v_m, 1 \leq m \leq n\},$$

where $(v_1, v_2, \dots, v_n) \in \prod_{m=1}^n (T^m)_n$. Clearly $\{\mathcal{P}_n\}$ witnesses that the product action is a special modular action. \dashv

Actually, since the sets $R_{v_1 \dots v_m}$ are clopen, this argument shows that the action of Γ on $\prod_{m=1}^{\infty} \partial T^m$ is isomorphic to the action of Γ on $\partial T_{\{\mathcal{P}_n\}}$ via a homeomorphism of the two spaces.

3 Maximality of the profinite action of \mathbb{F}_n

It is a general fact, proved independently by Glasner-Thouvenot-Weiss [GTW] and Hjorth (unpublished) (see also Kechris [Ke10], 10.7), that for each countable group Γ , the space $A(\Gamma, X, \mu)$ has a weakly dense conjugacy class, which in the language of weak containment can be stated as the existence of a

maximum (unique up to weak equivalence) action a_∞ in the order \prec of weak containment:

$$\exists a_\infty \in A(\Gamma, X, \mu) \forall b \in A(\Gamma, X, \mu) (b \prec a_\infty).$$

Such an a_∞ can always be assumed to be free (by replacing it if necessary by $a_\infty \times a$ for any free action a).

In Kechris [Ke10], 10.13, it is shown that for infinite, finitely generated Γ , the cost function

$$a \in \text{FR}(\Gamma, X, \mu) \mapsto C(a),$$

on the space of free actions with the weak topology, is upper semicontinuous. Since clearly C is invariant under conjugacy, it follows that for any $a, b \in \text{FR}(\Gamma, X, \mu)$, we have

$$b \prec a \Rightarrow C(b) \geq C(a)$$

(see Kechris [Ke10], 10.14) and thus for any free b ,

$$C(a_\infty) \leq C(b),$$

and therefore

$$C(a_\infty) = C(\Gamma) = \text{the cost of } \Gamma.$$

It is therefore of interest to be able to explicitly “compute” a_∞ for various groups Γ . We will do that below for $\Gamma = \mathbb{F}_n$, the free group with n generators ($1 \leq n \leq \infty$).

Since \mathbb{F}_n ($1 \leq n \leq \infty$) is a residually finite group, let $p_{\mathbb{F}_n}$ be the canonical action of \mathbb{F}_n on its profinite completion (see §2).

Theorem 3.1 *Let $1 \leq n \leq \infty$. The action $p_{\mathbb{F}_n}$ is maximum in the order \prec of weak containment of actions of \mathbb{F}_n .*

Proof. We consider for notational simplicity the case $n < \infty$, the argument in the other case being similar. We can work with $X = 2^\mathbb{N}$, μ the usual product measure. Clearly $A(\mathbb{F}_n, 2^\mathbb{N}, \mu)$ can be identified with $(\text{Aut}(2^\mathbb{N}, \mu))^n$, where $\text{Aut}(2^\mathbb{N}, \mu)$ is the group of measure preserving automorphisms of $(2^\mathbb{N}, \mu)$ with the weak topology. If $\mathbb{F}_n = \langle \gamma_1, \dots, \gamma_n \rangle$, with γ_i free generators, then $(S_1, \dots, S_n) \in (\text{Aut}(2^\mathbb{N}, \mu))^n$ is identified with the action $a \in A(\mathbb{F}_n, 2^\mathbb{N}, \mu)$ for which $\gamma_i^a = S_i$, $1 \leq i \leq n$. We will show that given any $a \in A(\mathbb{F}_n, 2^\mathbb{N}, \mu)$ and a weak open nbhd U of a , there is an action ρ of \mathbb{F}_n on a tree T , with ∂T uncountable, which is level transitive, such that if c is the associated action

of \mathbb{F}_n on the boundary ∂T , then U contains an isomorphic copy b of c . Then by §2, we have that c , and thus b , is a factor of $p_{\mathbb{F}_n}$, in particular $b \prec p_{\mathbb{F}_n}$. Thus a is the weak limit of actions $b \prec p_{\mathbb{F}_n}$, and so $a \prec p_{\mathbb{F}_n}$.

Fix an automorphism σ of the binary tree $2^{<\mathbb{N}}$ which is level transitive and denote by S the corresponding automorphism of $\partial 2^{<\mathbb{N}} = 2^{\mathbb{N}}$ (e.g., S could be the odometer). Identify the action a with (S_1, \dots, S_n) , where $\gamma_i^a = S_i$. Then we can assume that $U = U_1 \times \dots \times U_n$, where U_1, \dots, U_n are weak open nbhds of S_1, \dots, S_n , resp., in $\text{Aut}(2^{\mathbb{N}}, \mu)$.

By the Conjugacy Lemma (see, e.g., Kechris [Ke10], 2.4) there is $P \in \text{Aut}(2^{\mathbb{N}}, \mu)$ with $P^{-1}SP \in U_1$. Consider now $PS_2P^{-1}, \dots, PS_nP^{-1}$ and their open nbhds $PU_2P^{-1}, \dots, PU_nP^{-1}$. By the Weak Approximation Theorem (see, e.g., Kechris [Ke10], 2.1), there is large enough N and permutations π_2, \dots, π_n of 2^N (the set of binary sequences of length N) such that if S_{π_i} on $2^{\mathbb{N}}$ is defined by $S_{\pi_i}(s \hat{x}) = \pi_i(s) \hat{x}$, for $s \in 2^N, x \in 2^{\mathbb{N}}$, then $S_{\pi_i} \in PU_iP^{-1}$, $2 \leq i \leq n$. Consider now the action d of \mathbb{F}_n given by $(S, S_{\pi_2}, \dots, S_{\pi_n})$.

Let T be the tree consisting of all finite sequences $s = (s_0, s_1, \dots, s_{n-1})$ (including the empty sequence), where $s_0 \in 2^N, s_i \in \{0, 1\}$, if $1 \leq i < n$. The root of T is the empty sequence and the children of s are all sequences $t = (s_0, s_1, \dots, s_{n-1}, s_n), s_n \in \{0, 1\}$. We can define an action ρ of \mathbb{F}_n on T as follows: γ_1 acts as σ does; $\gamma_2, \dots, \gamma_n$ act via $\gamma_i \cdot (s_0, s_1, \dots, s_{n-1}) = \pi_i(s_0), s_1, \dots, s_{n-1}$. Since σ is level transitive, so is ρ . Let c be the associated action on ∂T . Clearly c is isomorphic to d via the isomorphism $(s_0, s_1, \dots) \in \partial T \mapsto s_0 \hat{s}_1 \hat{\dots}$. Now if $b = P^{-1}dP = (P^{-1}SP, P^{-1}S_{\pi_2}P, \dots, P^{-1}S_{\pi_n}P)$, then $b \in U_1 \times U_2 \times \dots \times U_n$, thus U contains an isomorphic copy b of c . \dashv

We now combine the preceding result with the main theorem in Abért-Nikolov [AN] to give a new proof that the cost of \mathbb{F}_n is equal to n ($1 \leq n < \infty$).

Let Γ be a finitely generated group and $\Gamma = \Gamma_0 \geq \Gamma_1 \geq \dots$ a chain. Let $T(\Gamma, \{\Gamma_n\})$ be the corresponding tree and let $a_{\Gamma, \{\Gamma_n\}}$ the associated action of Γ on $\partial T(\Gamma, \{\Gamma_n\})$. Note that if each Γ_n is normal and $\bigcap_n \Gamma_n = \{1\}$, then $a_{\Gamma, \{\Gamma_n\}}$ is free.

Theorem 3.2 (Abért-Nikolov [AN]) *If the action $a_{\Gamma, \{\Gamma_n\}}$ is free, then*

$$C(a_{\Gamma, \{\Gamma_n\}}) = \lim_{n \rightarrow \infty} \frac{d(\Gamma_n) - 1}{[\Gamma : \Gamma_n]} + 1,$$

where $d(\Gamma_n) = \text{rank}(\Gamma_n) = \text{minimum number of generators of } \Gamma_n$.

Remark. If Γ is finitely generated and $\Delta \leq \Gamma$ has finite index, then $d(\Delta) - 1 \leq [\Gamma : \Delta](d(\Gamma) - 1)$, with equality if Γ is free (see, Lyndon-Schupp [LS]). So if $H \leq \Delta \leq \Gamma$ have finite index, $\frac{d(H)-1}{[\Gamma : H]} \leq \frac{d(\Delta)-1}{[\Gamma : \Delta]}$, thus the sequence $\frac{d(\Gamma_n)-1}{[\Gamma : \Gamma_n]}$ is decreasing, and the above limit exists.

Now for $\Gamma = \mathbb{F}_n$ and any $H \leq \mathbb{F}_n$ of finite index, $\frac{d(H)-1}{[\Gamma : H]} = d(\mathbb{F}_n) - 1 = n - 1$. Therefore, since p_Γ is the action $a_{\Gamma, \{\Gamma_n\}}$, where Γ_n is a decreasing sequence of normal, finite index subgroups which is cofinal, we have that $C(p_{\mathbb{F}_n}) = n - 1$. Moreover, for any action a of \mathbb{F}_n , $C(a) \leq n$, so for any free action a of \mathbb{F}_n , $C(a) = n$. Thus we have shown:

Corollary 3.3 (Gaboriau [G]) *The cost of \mathbb{F}_n ($1 \leq n < \infty$) is equal to n and \mathbb{F}_n has fixed price.*

Remark. In a recent paper, Abért and Weiss [AW] have shown that if for an infinite group Γ , we let s_Γ be the shift action of Γ on 2^Γ (with the usual product measure), then s_Γ is weakly contained in any free action of Γ :

$$\forall b \in \text{FR}(\Gamma, X, \mu) (s_\Gamma \prec b).$$

Thus among the free actions of Γ there is a minimum in the order \prec . It follows that $C(s_\Gamma) \geq C(b), \forall b \in \text{FR}(\Gamma, X, \mu)$, i.e., s_Γ realizes the maximum cost of a free action of Γ .

Note now that there is a minimum, in the order \prec of weak containment, action in $A(\Gamma, X, \mu)$ (where we now consider arbitrary, not necessarily free, actions) iff the group Γ is amenable. Indeed, if $a \in A(\Gamma, X, \mu)$ is minimum in \prec , and $i_\Gamma \in A(\Gamma, X, \mu)$ is the trivial action of Γ , then $a \prec i_\Gamma$, so $a = i_\Gamma$, i.e., i_Γ is minimum. Then $i_\Gamma \prec s_\Gamma$, so (see Kechris [Ke10], 13.2) Γ is amenable. Conversely, if Γ is amenable, then $i_\Gamma \prec a$, for every ergodic $a \in A(\Gamma, X, \mu)$; see Kechris [Ke10], 10.6. Using the ergodic decomposition, this implies that $i_\Gamma \prec a$ for every $a \in A(\Gamma, X, \mu)$, so i_Γ is minimum.

In particular, since, for Γ amenable, s_Γ weakly contains any action (see Kechris [Ke10], 13.2), it follows that for amenable Γ , i_Γ is the minimum and s_Γ the maximum in the order \prec of weak containment in $A(\Gamma, X, \mu)$ and by Abért-Weiss so is every free action of Γ .

4 Density conditions in the space of actions

It is unclear for what infinite, residually finite groups Γ the analog of 3.1 goes through, i.e., the action p_Γ is maximum in the order \prec . Similarly, in the

finitely generated case, concerning the (weaker) condition that $C(p_\Gamma) = C(\Gamma)$. Recall that by Abért-Nikolov [AN], for finitely generated such Γ , $C(p_\Gamma) = \lim_{n \rightarrow \infty} \frac{d(\Gamma_n) - 1}{[\Gamma : \Gamma_n]} + 1 = \text{RG}(\Gamma) + 1$, where $\{\Gamma_n\}$ is a decreasing sequence of normal finite index subgroups which is cofinal, and $\text{RG}(\Gamma)$ is the *absolute rank gradient* of Γ ,

$$\text{RG}(\Gamma) = \inf_H \frac{d(H) - 1}{[\Gamma : H]},$$

the inf taken over all finite index $H \leq \Gamma$.

Denote by $i_\Gamma \times p_\Gamma$ the product of the trivial action i_Γ with p_Γ . This is an action whose ergodic decomposition consists of continuum many copies of p_Γ and thus (see Kechris-Miller [KM], 10.6) $C(i_\Gamma \times p_\Gamma) = C(p_\Gamma)$. So if $i_\Gamma \times p_\Gamma$ is maximum in the order \prec , we also have $C(p_\Gamma) = C(\Gamma)$, so it is also worth considering the (weaker) condition that $i_\Gamma \times p_\Gamma$ is such a maximum.

We will provide below some equivalent reformulations of these maximality conditions, relate them to some other properties considered in the literature, discuss some closure properties and raise some questions.

Definition 4.1 *Let Γ be an infinite, residually finite group. We say that Γ has the property EMD if the ergodic, finitely modular actions $a \in A(\Gamma, X, \mu)$ are weakly dense in $A(\Gamma, X, \mu)$.*

Recall from §2 that the ergodic, finitely modular actions in $A(\Gamma, X, \mu)$, up to isomorphism, can be viewed as actions on boundaries of trees on which the group acts in a level transitive way. Again by §2, up to isomorphism, these can be viewed as actions on $\partial T(\Gamma, \{\Gamma_n\})$, for some chain $\Gamma = \Gamma_0 \geq \Gamma_1 \geq \dots$, which is *proper* in the sense that the Γ_n do not eventually stabilize. (Recall here our convention that $A(\Gamma, X, \mu)$ always denotes the space of measure preserving actions of Γ on a *non-atomic* space (X, μ) , thus $\partial T(\Gamma, \{\Gamma_n\})$ is required to be uncountable, which means that $\{\Gamma_n\}$ is proper.)

Proposition 4.2 *Let Γ be an infinite, residually finite group. Then Γ has property EMD iff p_Γ is maximum in the order \prec of weak containment.*

Proof. If Γ has property EMD and $a \in A(\Gamma, X, \mu)$, then in any weakly open nbhd U of a there is an isomorphic copy of an action b of Γ on $\partial T(\Gamma, \{\Gamma_n\})$, for some proper chain $\Gamma = \Gamma_0 \geq \Gamma_1 \geq \dots$. By 2.3, $b \prec p_\Gamma$ and thus there is an isomorphic copy of p_Γ in U , so $a \prec p_\Gamma$ and p_Γ is maximum in \prec .

Conversely, if p_Γ is maximum in \prec , then the isomorphic copies of p_Γ are weakly dense in $A(\Gamma, X, \mu)$. But clearly p_Γ is an ergodic finitely modular action and we are done. \dashv

Recall that an infinite, residually finite group Γ satisfies *property* (τ) if $1_\Gamma \not\prec \kappa_0^{p_\Gamma}$, where $\kappa_0^{p_\Gamma}$ is the Koopman representation of p_Γ , restricted to the orthogonal of the constant functions, and 1_Γ is the trivial 1-dimensional representation.

Proposition 4.3 *Let Γ be an infinite residually finite group. Then if Γ has property EMD, Γ does not satisfy property (τ) .*

Proof. By 4.2, we have that if Γ has property EMD, p_Γ is maximum in the order \prec , so $i_\Gamma \prec p_\Gamma$. It follows that $1_\Gamma \prec \kappa_0^{p_\Gamma}$ (see Kechris [Ke10], Section 10, (C)), thus Γ does not have property (τ) . \dashv

Definition 4.4 *Let Γ be an infinite, residually finite group. Then Γ has property EMD* if the ergodic, finitely modular actions are weakly dense in the ergodic actions in $A(\Gamma, X, \mu)$.*

Clearly $\text{EMD} \Rightarrow \text{EMD}^*$. If Γ does not have property (T), then, by a result of Glasner-Weiss (see, e.g., Kechris [Ke10], 12.2), the ergodic actions are dense in $A(\Gamma, X, \mu)$, so $\text{EMD} \Leftrightarrow \text{EMD}^*$ for such Γ . The following is proved as in 4.2.

Proposition 4.5 *Let Γ be an infinite, residually finite group. Then Γ has property EMD* iff p_Γ is maximum in the order \prec among the ergodic actions in $A(\Gamma, X, \mu)$.*

We will now consider the final weakening of these properties.

Definition 4.6 *Let Γ be an infinite, residually finite group. Then Γ has property MD if the finitely modular actions are dense in $A(\Gamma, X, \mu)$.*

Definition 4.7 *An action $a \in A(\Gamma, X, \mu)$ is finite if it factors through a finite group, i.e., there is a finite group Δ , a homomorphism $\pi: \Gamma \rightarrow \Delta$ and an action $b \in A(\Delta, X, \mu)$ such that $\gamma^a = \pi(\gamma)^b$. Equivalently a is finite if $\{\gamma^a: \gamma \in \Gamma\}$ is a finite subgroup of $\text{Aut}(X, \mu)$.*

Proposition 4.8 *Let Γ be an infinite, residually finite group. Then the following are equivalent:*

- (i) Γ has property MD.
- (ii) The finite actions of Γ are weakly dense in $A(\Gamma, X, \mu)$.
- (iii) The action $i_\Gamma \times p_\Gamma$ is maximum in the order \prec of weak containment.

Proof. The finite actions are clearly finitely modular, so (ii) \Rightarrow (i).

We next show that (i) \Rightarrow (ii). For that it is enough to show that the finite action are dense in the finitely modular ones. Again for that it is enough to show that if $X = P_1 \sqcup \dots \sqcup P_n$ is a partition of X into finitely many sets of positive measure and a is an action of Γ on (X, μ) that leaves the partition $\mathcal{P} = \{P_1, \dots, P_n\}$ invariant, then there is a finite action b of Γ on (X, μ) that agrees with a on \mathcal{P} , i.e., $\forall \gamma \in \Gamma \forall P \in \mathcal{P} (\gamma^a(P) = \gamma^b(P))$.

It is clear that there is a finite group Δ , an action c of Δ on \mathcal{P} and a surjective homomorphism $\pi: \Gamma \rightarrow \Delta$ such that for each $P \in \mathcal{P}, \gamma \in \Gamma, \gamma^a(P) = \pi(\gamma)^c(P)$. (Simply take Δ to be the group of permutations of \mathcal{P} induced by a .) Next for each $P \in \mathcal{P}, \mu_P = \frac{\mu|_P}{\mu(P)}$ is a non-atomic probability measure on P , so fix a measure preserving bijection $\varphi_P: (P, \mu_P) \rightarrow ([0, 1], \lambda)$ (where λ is the Lebesgue measure). Put for any $P, Q \in \mathcal{P}, \varphi_{P,Q} = \varphi_Q^{-1} \circ \varphi_P$. Then $\varphi_{P,Q}: (P, \mu_P) \rightarrow (Q, \mu_Q)$ is a measure preserving bijection and $\varphi_{P,P} = \text{id}, \varphi_{P,R} = \varphi_{Q,R} \circ \varphi_{P,Q}$. If P, Q are in the same Δ -orbit, then $\mu(P) = \mu(Q)$, thus $\varphi_{P,Q}: (P, \mu|_P) \rightarrow (Q, \mu|_Q)$ is also measure preserving.

We now define an action $d \in A(\Delta, X, \mu)$ as follows: let $x \in X$ and let $P \in \mathcal{P}$ be such that $x \in P$. Let $\delta^c(P) = Q$ and put $\delta^d(x) = \varphi_{P,Q}(x)$. From the preceding discussion, it is clear that d is measure preserving and $\delta^d(P) = \delta^c(P), \forall P \in \mathcal{P}$. Finally, define $b \in A(\Gamma, X, \mu)$ by $\gamma^b = \pi(\gamma)^d$. Then $\gamma^b(P) = \pi(\gamma)^d(P) = \pi(\gamma)^c(P) = \gamma^a(P), \forall P \in \mathcal{P}$, and the proof that (i) \Rightarrow (ii) is complete.

Since the action $i_\Gamma \times p_\Gamma$ is finitely modular, it is clear that (iii) \Rightarrow (i). Finally, we show that (i) \Rightarrow (iii).

It is enough to show that if $a \in A(\Gamma, X, \mu)$ is finitely modular, then $a \prec i_\Gamma \times p_\Gamma$. The action a is isomorphic to a measure preserving action b on the boundary of a tree, thus by §2 each component in the ergodic decomposition of b is isomorphic to an action of Γ on $\partial T(\Gamma, \{\Gamma_n\})$ for some chain $\Gamma = \Gamma_0 \geq \Gamma_1 \geq \dots$. By 2.3, the action of Γ on $\partial T(\Gamma, \{\Gamma_n\})$ is weakly contained in p_Γ , from which it follows that the action b (and thus the action a) is weakly contained in $i_\Gamma \times p_\Gamma$. \dashv

A similar argument as in the proof of (i) \Rightarrow (iii) in Proposition 4.8, together with 4.5, shows that

$$\text{EMD}^* \Rightarrow \text{MD},$$

thus

$$\text{EMD} \Rightarrow \text{EMD}^* \Rightarrow \text{MD}.$$

Moreover, MD implies, for finitely generated Γ , that $C(\Gamma) = C(p_\Gamma) = \text{RG}(\Gamma)$.

Question 4.9 *Does $\text{EMD}^* \Rightarrow \text{EMD}$?*

As we have seen, the answer is positive if Γ does not have property (T), so one has the following further problem:

Question 4.10 *Does EMD^* imply $\neg(\text{T})$?*

Next we have the question:

Question 4.11 *Does $\text{MD} \Rightarrow \text{EMD}^*$?*

A partial answer to these questions is given by the following fact:

Proposition 4.12 *Let Γ be an infinite, finitely generated, residually finite group that does not have property (τ) . Then $\text{EMD} \Leftrightarrow \text{EMD}^* \Leftrightarrow \text{MD}$.*

Proof. It is shown in Abért-Weiss [AW], that if Γ is as in the statement of the proposition, then $p_\Gamma \sim i_\Gamma \times p_\Gamma$. From this it immediately follows that $\text{MD} \Rightarrow \text{EMD}$. \dashv

In view of this, and the fact that $\text{EMD} \Rightarrow \neg(\tau)$, it is clear that the question of whether $\text{EMD} \Leftrightarrow \text{MD}$ is equivalent to

Question 4.13 *Does MD imply $\neg(\tau)$?*

Note that for infinite, finitely generated, residually finite Γ , $\text{MD} \Rightarrow \neg(\tau)$ iff $(\text{MD} \Rightarrow \text{EMD}^*)$ & $(\text{EMD}^* \Rightarrow \neg(\text{T}))$. It is also unknown whether MD implies $\neg(\text{T})$.

In Lubotzky-Shalom [LSh], the authors say that a residually finite group Γ has property FD if the finite unitary representations (i.e., those factoring through a finite quotient of Γ) are dense in the space $\text{Rep}(\Gamma, \mathcal{H})$. Using the fact that any $\pi \in \text{Rep}(\Gamma, \mathcal{H})$ is a subrepresentation of κ_0^a for some $a \in A(\Gamma, X, \mu)$ (see, e.g., Kechris [Ke10], E.1) it follows easily that $\text{MD} \Rightarrow \text{FD}$

but it is unknown whether $\text{FD} \Rightarrow \text{MD}$. In Lubotzky-Zuk [LZ], the question is raised of whether $\text{FD} \Rightarrow \neg(\tau)$. Thus 4.13 is an ergodic theory analog of this problem.

In particular, the various examples of groups that fail property FD given, e.g., in Lubotzky-Zuk [LZ], 9.1, also fail property MD.

On the other hand, we have seen in 3.1 that \mathbb{F}_n ($1 \leq n \leq \infty$) has property MD and in fact the property EMD.

We can also see that every infinite, residually finite, amenable group also has property EMD. To see this let s_Γ be the shift action of Γ on 2^Γ . Then Abért-Weiss [AW] have shown that $s_\Gamma \prec a$ for every free action $a \in A(\Gamma, X, \mu)$. Thus $s_\Gamma \prec p_\Gamma$. But since Γ is amenable, $\forall a \in A(\Gamma, X, \mu)(a \prec s_\Gamma)$ (see Kechris [Ke10], 13.2), so $\forall a \in A(\Gamma, X, \mu)(a \prec p_\Gamma)$. Thus Γ has EMD.

We next note some closure properties of the class MD. First, if Δ is infinite, residually finite with property MD and $\Gamma \leq \Delta$ is infinite, then Γ has property MD. To see this, let $a \in A(\Gamma, X, \mu)$ and let $\text{CInd}_\Gamma^\Delta(a) \in A(\Delta, Y, \nu)$ be the co-induced action (see the Appendix). By assumption $\text{CInd}_\Gamma^\Delta(a) \prec i_\Delta \times p_\Delta$, thus $\text{CInd}_\Gamma^\Delta(a)|\Gamma \prec (i_\Delta \times p_\Delta)|\Gamma = b$. Clearly b is finitely modular. Moreover a is a factor of $\text{CInd}_\Gamma^\Delta(a)|\Gamma$, so $a \prec \text{CInd}_\Gamma^\Delta(a)|\Gamma \prec b$, thus every action of Γ is weakly contained in a finitely modular action, so Γ satisfies also MD. It is not clear if the same fact holds for property EMD.

In the opposite direction, if Γ is infinite, residually finite with property MD and $\Gamma \leq \Delta$ is such that $[\Delta : \Gamma] < \infty$, then Δ has also property MD. To see this let $b \in A(\Delta, X, \mu)$. Let $a = b|_\Gamma$ and let $c = \text{Ind}_\Gamma^\Delta(a) = \text{Ind}_\Gamma^\Delta(b|_\Gamma)$ be the induced action (see, e.g., Zimmer [Z] or Kechris [Ke10], Appendix G). By Zimmer [Z], 4.2.22, $c \cong b \times a_{\Delta/\Gamma}$, where $a_{\Delta/\Gamma}$ is the action of Δ on Δ/Γ . In particular $b \prec c$. By assumption $a \prec i_\Gamma \times p_\Gamma$ and so $b \prec \text{Ind}_\Gamma^\Delta(a) = c \prec \text{Ind}_\Gamma^\Delta(i_\Gamma \times p_\Gamma) = d$. (We are using here that inducing preserves weak containment of actions, which can be verified as in Appendix A.1.) It is easy to see that d is finitely modular and this shows that Δ also has property MD. Since the induced action of an ergodic action is ergodic the same argument works as well for the property EMD. In particular, the group $\text{SL}_2(\mathbb{Z})$ and all groups of the form $A * B$, where A, B are finite non-trivial groups, have EMD.

In Lubotzky-Shalom [LSH] (see also Lubotzky-Zuk [LZ], 9.11 and 9.12), it is shown that, in certain situations, if $\Gamma \triangleleft \Delta$ and Γ has property FD, so does Δ . This is used to provide additional examples of groups with property FD. One would like to prove a similar result for the property MD. The proof of the Lubotzky-Shalom result uses the fact that if $\Gamma \leq \Delta$, the action of

Δ on Δ/Γ is amenable (i.e., admits a finitely additive invariant probability measure) and π is a unitary representation of Δ , then π is weakly contained in the induced representation $\text{Ind}_\Gamma^\Delta(\pi|_\Gamma)$ of the restriction of π to Γ . To prove the analog for the property MD, one would like to have the corresponding result for co-induced actions, but this appears to be unknown - see Problem A.4. Let us state the result about MD that one would like to prove and its implications and then show that an affirmative action to A.4 would indeed provide a proof.

Conjecture 4.14 *Let Γ be an infinite, finitely generated, residually finite group satisfying MD. Let $\Gamma \triangleleft \Delta$ with Δ residually finite. Assume that:*

- (i) *For every $N \triangleleft \Gamma$ with $[\Gamma : N] < \infty$, there is $M \triangleleft \Delta$ such that $M \subseteq N$ and $[\Gamma : M] < \infty$.*
- (ii) *Δ/Γ is a residually finite, amenable group.*

Then Δ satisfies MD.

The result of Lubotzky-Shalom [LSh] is that this holds if MD is replaced by FD.

If the Conjecture 4.14 has a positive answer, it would produce the following additional examples of groups with property MD, since Lubotzky-Shalom [LSh] and Lubotzky-Zuk [LZ], 9.2., verify that the groups below satisfy the conditions of 4.14.

- (a) $H \rtimes \mathbb{F}_n$, where H is a residually finite, amenable group.
- (b) The surface groups

$$T_g = \langle a_1, \dots, a_g, b_1, \dots, b_g \mid \prod_{i=1}^g [a_i, b_i] = 1 \rangle$$

and the groups

$$\text{SL}_2(\mathbb{Z}[i]), \text{SL}_2(\mathbb{Z}[\sqrt{3}i]).$$

Let us now show that an affirmative answer to A.4 gives a proof of 4.14.

We can assume that $[\Delta : \Gamma] = \infty$. Let $a \in A(\Delta, X, \mu)$. Consider the restriction $a|_\Gamma \in A(\Gamma, X, \mu)$ and the co-induced action

$$\text{CInd}_\Gamma^\Delta(a|_\Gamma) \in A(\Delta, X^{\Gamma/\Delta}, \mu^{\Gamma/\Delta}).$$

Let $a_i \in A(\Gamma, X, \mu)$ be finite actions such that $a_i \rightarrow a|_\Gamma$ weakly. Let F_i be a finite group, $b_i \in A(F_i, X, \mu)$ and $\pi_i : \Gamma \rightarrow F_i$ a surjective homomorphism

such that $\gamma^{a_i} = \pi_i(\gamma)^{b_i}$. Let $N_i = \ker(\pi_i)$, so that $N_i \triangleleft \Gamma$, $F_i = \Gamma/N_i$ and $[\Gamma : N_i] < \infty$. By i), there is $M_i \triangleleft \Delta$ such that $M_i \subseteq N_i$ and $[\Gamma : M_i] < \infty$. It follows (by replacing N_i by M_i if necessary) that we can assume that $N_i \triangleleft \Delta$.

Lemma 4.15 *Let F be a finite group, $b \in A(F, X, \mu)$ and let $s_{F,X}$ be the shift action of F on X^F . Then $b \prec s_{F,X}$.*

Proof. Consider $F \times \mathbb{Z}$ and the action $b' \in A(F \times \mathbb{Z}, X, \mu)$ given by $(\gamma, g)^{b'} = \gamma^b$, where $(\gamma, g) \in F \times \mathbb{Z}$. Since $F \times \mathbb{Z}$ is infinite, amenable, we have that if $s_{F \times \mathbb{Z}}$ is the shift action of $F \times \mathbb{Z}$ on $2^{F \times \mathbb{Z}}$, then $b' \prec s_{F \times \mathbb{Z}}$ (see Kechris [Ke10], 13.2). Thus $b = b'|F \prec s_{F \times \mathbb{Z}}|F$. But clearly $s_{F \times \mathbb{Z}}|F$ is isomorphic to the shift action of F on $(\mathbb{Z})^F$ which is isomorphic to the shift action of F on X^F (since (X, μ) is non-atomic) and we are done. \dashv

It follows that $b_i \prec s_{F_i, X}$ and thus $a_i \prec s_{\Gamma, \Gamma/N_i, X}$, where $s_{\Gamma, \Gamma/N_i, X}$ is the (generalized) shift action of Γ on X^{Γ/N_i} , defined by $\delta \cdot p(gN_i) = g(\delta^{-1}gN_i)$.

Now $\text{CInd}_{\Gamma}^{\Delta}(a_i) \rightarrow \text{CInd}_{\Gamma}^{\Delta}(a|\Gamma)$ weakly, and $\text{CInd}_{\Gamma}^{\Delta}(a_i) \prec \text{CInd}_{\Gamma}^{\Delta}(s_{\Gamma, \Gamma/N_i, X})$ (see the Appendix).

By Proposition A.2 in the Appendix, $\text{CInd}_{\Gamma}^{\Delta}(s_{\Gamma, \Gamma/N_i, X}) \cong s_{\Delta, \Delta/N_i, X}$, so $\text{CInd}_{\Gamma}^{\Delta}(a_i) \prec s_{\Delta, \Delta/N_i, X}$. But Δ/N_i is infinite, amenable, and residually finite, by (ii), so $s_{\Delta, \Delta/N_i, X}$ is a weak limit of finite actions of Δ/N_i and thus $s_{\Delta, \Delta/N_i, X}$ is a weak limit of finite actions of Δ . It follows that each $\text{CInd}_{\Gamma}^{\Delta}(a_i)$ is a limit of finite actions of Δ and thus so is $\text{CInd}_{\Gamma}^{\Delta}(a|\Gamma)$. If A.4 in the Appendix has a positive answer, we have that $a \prec \text{CInd}_{\Gamma}^{\Delta}(a|\Gamma)$. It follows that a is also a weak limit of finite actions of Δ and the proof is complete.

5 Maximality of a generalized shift action of

\mathbb{F}_n

We have seen in 3.1 that the profinite action $p_{\mathbb{F}_n}$ ($1 \leq n \leq \infty$) is maximum in the order \prec of weak containment of actions of \mathbb{F}_n . We have also mentioned the result of Abért-Weiss [AW] that, for any infinite Γ , the shift action s_{Γ} of Γ on 2^{Γ} is minimum in the order \prec among free actions. More generally, one can consider an action of Γ on a countable set I and the induced *generalized shift* action $s_{\Gamma, I}$ of Γ on 2^I given by $\gamma \cdot p(i) = p(\gamma^{-1} \cdot i)$. Of special interest are the generalized shifts corresponding to transitive actions of Γ on I . Equivalently these are the generalized shifts induced by the canonical action of Γ on $I = \Gamma/H$ for some $H \leq \Gamma$. We denote this by $s_{\Gamma, \Gamma/H}$. In the case of an arbitrary

action of Γ on I we can decompose $I = \bigsqcup_n I_n$, into the Γ -orbits and then clearly the Γ -shift on $2^I = 2^{\bigsqcup_n I_n}$ is (isomorphic to) the product of the Γ -shifts on each 2^{I_n} . Thus the generalized shifts are just the products of generalized shifts of the form $s_{\Gamma, \Gamma/H}$.

Theorem 5.1 *Let $\Gamma = \mathbb{F}_n$ ($1 \leq n \leq \infty$). Then there is $H \leq \Gamma$ with $[\Gamma : H] = \infty$ such that the generalized shift $s_{\Gamma, \Gamma/H}$ of Γ on $2^{\Gamma/H}$ is maximum in the order \prec of weak containment of actions of Γ .*

Proof. Let $\Gamma = \mathbb{F}_n = \langle \gamma_1, \gamma_2, \dots \rangle$ be free generators. We will consider the space $A(\Gamma, X, \mu)$ of measure preserving action of Γ on (X, μ) with the weak topology. This can be identified with the product space $\text{Aut}(X, \mu)^n$ (where n denotes $\{1, \dots, n\}$, if $n < \infty$, and $\mathbb{N} \setminus \{0\}$, if $n = \infty$), with $\text{Aut}(X, \mu)$ again equipped with the weak topology, by identifying $a \in A(\Gamma, X, \mu)$ with $(\gamma_i^a)_i \in \text{Aut}(X, \mu)^n$.

We say that $a \in A(\Gamma, X, \mu)$ *factors through* a group Δ if there is a surjective homomorphism $\rho: \Gamma \rightarrow \Delta$ and $b \in A(\Delta, X, \mu)$ such that $\forall \gamma \in \Gamma (\gamma^a = \rho(\gamma)^b)$. We will say that $a \in A(\Gamma, X, \mu)$ *factors regularly through* Δ if the above holds but additionally $\rho(\gamma_i)$ has infinite order in $\Delta, \forall i$. In that case we call ρ *regular*.

Lemma 5.2 *The actions of Γ on (X, μ) that factor regularly through an infinite amenable group are weakly dense in $A(\Gamma, X, \mu)$.*

Proof. By §4 the group Γ has property MD, thus the finite actions are weakly dense in $A(\Gamma, X, \mu)$. Consider such an action $a \in A(\Gamma, X, \mu)$, let $\sigma: \Gamma \rightarrow F$ be a surjective homomorphism, where F is finite, and let $c \in A(F, X, \mu)$ be such that $\gamma^a = \sigma(\gamma)^c, \forall \gamma \in \Gamma$. Let $\Delta_1 = F \times \mathbb{Z}$ and let $\pi: \Delta_1 \rightarrow F$ be the projection to the first coordinate. Let $g \neq 0$ be an element of \mathbb{Z} . Let $\tau: \Gamma \rightarrow \Delta_1$ be the homomorphism defined by $\tau(\gamma_i) = (\sigma(\gamma_i), g)$, so that $\tau(\gamma_i)$ has infinite order. Let $\tau(\Gamma) = \Delta \subseteq \Delta_1$. Thus Δ is an infinite amenable group, and if $b \in A(\Delta, X, \mu)$ is defined by $\delta^b = \pi(\delta)^c$, then, as $\sigma = \pi \circ \tau$, we have $\gamma^a = \sigma(\gamma)^c = (\pi(\tau(\gamma)))^c = \tau(\gamma)^b$, i.e., a factors regularly through Δ . \dashv

Suppose now $\rho: \Gamma \rightarrow \Delta$ is a regular surjective homomorphism, where Δ is infinite amenable, $b \in A(\Delta, X, \mu)$ and $a \in A(\Gamma, X, \mu)$ is given by $\gamma^a = \rho(\gamma)^b$. Since Δ is infinite amenable, $b \prec s_\Delta$, where s_Δ is the shift action of Δ on 2^Δ (see Kechris [Ke10], 13.2). Composing with ρ , we see that $a \prec s_{\Gamma, \Gamma/N}$, where $N = \ker(\rho)$.

To simplify notation, we will work from now on with $\Gamma = \mathbb{F}_2 = \langle \gamma_1, \gamma_2 \rangle$. The general case requires only trivial modifications.

Fix a countable open basis $\{U_n\}$ for $A(\Gamma, X, \mu)$, so that every U_n has the following form:

$$U_n = \{a \in A(\Gamma, X, \mu) : \forall i \leq 2\forall j, k \leq m_n |\mu(\gamma_i^a(P_j^{(n)}) \cap P_k^{(n)}) - \mu(\gamma_i^{a_n}(P_j^{(n)}) \cap P_k^{(n)})| < \epsilon_n,$$

where $a_n \in A(\Gamma, X, \mu)$, $\mathcal{P}_n = \{P_1^{(n)}, \dots, P_{m_n}^{(n)}\}$ is a Borel partition of X and $\epsilon_n > 0$ (see Kechris [Ke10], Section 1, (B)).

From 5.2, and the paragraph following it, it follows that for each n , there is an infinite set I_n , a transitive action τ_n of Γ on I_n such that no $\gamma_i^{\tau_n}$ ($i \leq 2$) has an invariant finite set (this is where regularity is used) and is such that an isomorphic copy \bar{a}_n of s_{Γ, I_n} is in U_n , say via the isomorphism $\varphi_n: (2^{I_n}, \nu_n) \rightarrow (X, \mu)$, where ν_n is the usual product measure on 2^{I_n} . Let $\varphi_n^{-1}(\mathcal{P}_n) = \mathcal{R}_n = \{R_1^{(n)}, \dots, R_{m_n}^{(n)}\}$ be the partition of 2^{I_n} in which $R_j^{(n)} = \varphi_n^{-1}(P_j^{(n)})$, so that $\forall i \leq 2\forall j, k \leq m_n, \nu_n(R_j^{(n)}) = \mu(P_j^{(n)})$ and $|\nu_n(\gamma_i^{s_{\Gamma, I_n}}(R_j^{(n)}) \cap R_k^{(n)}) - \mu(\gamma_i^{a_n}(P_j^{(n)}) \cap P_k^{(n)})| < \rho_n < \epsilon_n$.

A basic nbhd of 2^{I_n} is a set of the form $\{p \in 2^{I_n} : p|F = u\}$, where $u \in 2^F$, F finite. A finite union of such basic nbhds, with the same F , will be called a *clopen set supported by F* . Since the clopen sets with finite support are dense in the measure algebra of 2^{I_n} , it follows that for every $\delta > 0$, there is a finite non-empty set $F_{n,\delta}$ and a partition of 2^{I_n} into clopen sets supported by $F_{n,\delta}$, $\mathcal{S}_{n,\delta} = \{S_{1,\delta}^{(n)}, \dots, S_{m_{n,\delta}}^{(n)}\}$, such that $\forall j \leq m_n (\nu_n(R_j^{(n)} \Delta S_{j,\delta}^{(n)}) < \delta)$, thus $\forall i \leq 2\forall j, k \leq m_n (|\nu_n(\gamma_i^{s_{\Gamma, I_n}}(S_{j,\delta}^{(n)}) \cap S_{k,\delta}^{(n)}) - \nu_n(\gamma_i^{s_{\Gamma, I_n}}(R_j^{(n)}) \cap R_k^{(n)})| < 2\delta)$.

Let $\bar{\mathcal{P}}_{n,\delta} = \varphi_n(\mathcal{S}_{n,\delta}) = \{\bar{P}_{1,\delta}^{(n)}, \dots, \bar{P}_{m_{n,\delta}}^{(n)}\}$ be the partition of X in which $\bar{P}_{j,\delta}^{(n)} = \varphi_n(S_{j,\delta}^{(n)})$. Then $\mu(\bar{P}_{j,\delta}^{(n)} \Delta P_j^{(n)}) < \delta, \forall j \leq m_n$, so $\forall i \leq 2\forall j, k \leq m_n$,

$$|\mu(\gamma_i^{\bar{a}_n}(\bar{P}_{j,\delta}^{(n)}) \cap \bar{P}_{k,\delta}^{(n)}) - \mu(\gamma_i^{a_n}(\bar{P}_{j,\delta}^{(n)}) \cap \bar{P}_{k,\delta}^{(n)})| < \rho_n + 4\delta.$$

Fix $\delta = \delta_n$ and ϵ'_n , so that $\rho_n + 4\delta_n < \epsilon'_n < \epsilon'_n + 4\delta_n < \epsilon_n$. Let U'_n be the set of all $\{a \in A(\Gamma, X, \mu)$ such that $\forall i \leq 2\forall j, k \leq m_n$

$$|\mu(\gamma_i^a(\bar{P}_{j,\delta_n}^{(n)}) \cap \bar{P}_{k,\delta_n}^{(n)}) - \mu(\gamma_i^{a_n}(\bar{P}_{j,\delta_n}^{(n)}) \cap \bar{P}_{k,\delta_n}^{(n)})| < \epsilon'_n\}.$$

Then $\bar{a}_n \in U'_n$ and $U'_n \subseteq U_n$. Put $\bar{P}_j^{(n)} = \bar{P}_{j,\delta_n}^{(n)}, \bar{\mathcal{P}}_n = \mathcal{P}_{n,\delta_n}, F_n = F_{n,\delta_n}, \mathcal{S}_n = \mathcal{S}_{n,\delta_n}$. Thus $\varphi_n: 2^{I_n} \rightarrow X$ sends \mathcal{S}_n to $\bar{\mathcal{P}}_n$, \mathcal{S}_n is supported by F_n and φ_n sends

s_{Γ, I_n} to \bar{a}_n , where

$$|\mu(\gamma_i^{\bar{a}_n}(\bar{P}_j^{(n)}) \cap \bar{P}_k^{(n)}) - \mu(\gamma_i^{a_n}(\bar{P}_j^{(n)}) \cap \bar{P}_k^{(n)})| < \epsilon'_n,$$

$\forall i \leq 2\forall j, k \leq m_n$. Also $\mu(\bar{P}_j^{(n)}) = \nu_n(S_j^{(n)})$.

Now consider the product action $s = \prod_n s_{\Gamma, I_n}$, which is the action of Γ on $2^I, I = \bigsqcup_n I_n$, induced by the action τ of Γ on I given by $\tau = \bigsqcup_n \tau_n$. For each n , let $p_n: 2^I \rightarrow 2^{I_n}$ be the projection function $p_n(f) = f|_{I_n}$. Let $\bar{\mathcal{S}}_n = p_n^{-1}(\mathcal{S}_n) = \{p_n^{-1}(S_1^{(n)}), \dots, p_n^{-1}(S_{m_n}^{(n)})\}$. This is a clopen partition of 2^I supported by the finite set $F_n \subseteq I_n \subseteq I$. If $\nu = \prod_n \nu_n$ is the product measure on 2^I , clearly $\forall i \leq 2\forall j, k \leq m_n (\nu(\bar{S}_j^{(n)}) = \nu_n(S_j^{(n)}))$ and $\nu_n(\gamma_i^{s_{\Gamma, I_n}}(S_j^{(n)}) \cap S_k^{(n)}) = \nu(\gamma_i^s(\bar{S}_j^{(n)}) \cap \bar{S}_k^{(n)})$.

Now each $\bar{S}_j^{(n)}$ is a finite union of basic nbhds of the form $N_u = \{f \in 2^I : f|_{F_n} = u\}$, where $u \in 2^{F_n}$. Since $\gamma_i^s(N_u) = N_v$, where $v \in 2^{\gamma_i^{\tau_n}(F_n)}$ is defined by $v(a) = u((\gamma_i^{-1})^{\tau_n}(a))$, it follows that $\gamma_i^s(N_u)$, for each such u , and thus $\gamma_i^s(\bar{S}_j^{(n)}), \forall j \leq m_n$, depends only on $\gamma_i^{\tau}|_{F_n} = \gamma_i^{\tau_n}|_{F_n}$.

So fix any action τ^* of Γ on I such that $\gamma_i^{\tau^*}|_{F_n} = \gamma_i^{\tau}|_{F_n} = \gamma_i^{\tau_n}|_{F_n}, \forall n \forall i \leq 2$, and let s^* be the corresponding shift action on 2^I . Then $\gamma_i^s(\bar{S}_j^{(n)}) = \gamma_i^{s^*}(\bar{S}_j^{(n)}), \forall n \forall i \leq 2 \forall j \leq m_n$.

Lemma 5.3 *For each n there is an isomorphic copy of s^* in U'_n , thus s^* is maximum in \prec .*

Proof. Fix n . We will find an isomorphic copy of s^* in U'_n . Since $\mu(\bar{P}_j^{(n)}) = \nu_n(S_j^{(n)}) = \nu(\bar{S}_j^{(n)})$, let $\psi_n: (2^I, \nu) \rightarrow (X, \mu)$ be an isomorphism such that $\psi_n(\bar{S}_j^{(n)}) = \bar{P}_j^{(n)}, \forall j \leq m_n$. Let $a_n^* \in A(\Gamma, X, \mu)$ be the isomorphic copy of s^* induced by ψ_n . We will check that $a_n^* \in U'_n$. Indeed, $\mu(\gamma_i^{a_n^*}(\bar{P}_j^{(n)}) \cap \bar{P}_k^{(n)}) = \nu(\gamma_i^{s^*}(\bar{S}_j^{(n)}) \cap \bar{S}_k^{(n)}) = \nu(\gamma_i^s(\bar{S}_j^{(n)}) \cap \bar{S}_k^{(n)}) = \nu_n(\gamma_i^{s_{\Gamma, I_n}}(S_j^{(n)}) \cap S_k^{(n)}) = \mu(\gamma_i^{\bar{a}_n}(\bar{P}_j^{(n)}) \cap \bar{P}_k^{(n)})$, and so $|\mu(\gamma_i^{a_n^*}(\bar{P}_j^{(n)}) \cap \bar{P}_k^{(n)}) - \mu(\gamma_i^{a_n}(\bar{P}_j^{(n)}) \cap \bar{P}_k^{(n)})| < \epsilon'_n$, i.e., $a_n^* \in U'_n$. \dashv

To complete the proof, it is enough to show that such a τ^* can be chosen so that τ^* acts transitively on I . Let us recall that $I = \bigsqcup_n I_n, I_n$ infinite, $\emptyset \neq F_n \subseteq I_n$ is finite, $\tau = \bigsqcup_n \tau_n, \gamma_i^{\tau_n}$ does not have any finite (non-empty) invariant sets ($\subseteq I_n$) and we want to find τ^* , a transitive action of Γ on I , such that $\forall i \leq 2 \forall n (\gamma_i^{\tau^*}|_{F_n} = \gamma_i^{\tau}|_{F_n})$. First we take $\gamma_1^{\tau^*} = \gamma_1^{\tau}$. It is then enough to find $\gamma_2^{\tau^*}$, a transitive permutation of I , that satisfies $\gamma_2^{\tau^*}|_{F_n} = \gamma_2^{\tau}|_{F_n}$, for each n . This is possible by the following lemma.

Lemma 5.4 *For each $n \geq 1$, let I_n be an infinite (countable) set, $\emptyset \neq F_n \subseteq I_n$ a finite subset and S_n a permutation of I_n that has no finite (non-empty) invariant sets. Then there is a transitive permutation S of I such that $S|_{F_n} = S_n|_{F_n}, \forall n$.*

Proof. Because S_n has no finite non-empty invariant sets, i.e., no cycles, there is a partition $P_1^{(n)}, \dots, P_{k_n}^{(n)}$ of $F_n \cup S_n(F_n)$, such that each $P_i^{(n)}$ has the form $\{x_i^{(n)}, S_n(x_i^{(n)}), \dots, S_n^{\ell_n}(x_i^{(n)}) = y_i^{(n)}\}$, for some ℓ_n , where all the $S_n^j(x_i^{(n)})$ are distinct and $x_i^{(n)} \notin S_n(F_n), y_i^{(n)} \notin F_n$. Call $x_i^{(n)}, y_i^{(n)}$ the first, resp., last, elements of $P_i^{(n)}$. Clearly $S_n|_{F_n} = \bigcup_{i \leq k_n} S_n|(P_i^{(n)} \setminus \{y_i^{(n)}\})$. Put $P = I \setminus \bigcup_{n, i \leq k_n} P_i^{(n)}$ and enumerate $P = \{p_0, p_1, \dots\}$. Then let S be defined as follows: $S(p_{i+1}) = p_i$ for $i \geq 0$, $S|(P_i^{(n)} \setminus \{y_i^{(n)}\}) = S_n|(P_i^{(n)} \setminus \{y_i^{(n)}\})$ (thus $S|_{F_n} = S_n|_{F_n}$), and finally $S(p_0) = x_1^{(1)}, S(y_1^{(1)}) = x_2^{(1)}, S(y_2^{(1)}) = x_3^{(1)}, \dots, S(y_{k_1}^{(1)}) = x_1^{(2)}, S(y_1^{(2)}) = x_2^{(2)}, \dots$. \dashv

By a similar argument, using direct sums of representations instead of product actions, one can show that for $\Gamma = \mathbb{F}_n$ ($1 \leq n \leq \infty$), there is $H \leq \Gamma$ with $[\Gamma : H] = \infty$ such that the quasi-regular representation $\lambda_{\Gamma/H}$ on $\ell^2(\Gamma/H)$, induced by the canonical action of Γ on Γ/H , is maximum in the order \prec of weak containment of unitary representations of Γ . Monod (private communication) asked whether, for $n \geq 2$, one can also find such an H such that moreover $\lambda_{\Gamma/H}$ is irreducible. (The existence of irreducible representations of \mathbb{F}_n , for $n \geq 2$, that are maximum in the order of weak containment was first proved by Yoshizawa [Y]. For another proof, see Kechris [Ke10], Appendix H, (C)). One can easily modify the preceding argument to show that this is indeed the case. (I would like to thank T. Tsankov for a helpful discussion on this matter.)

Theorem 5.5 *Let $\Gamma = \mathbb{F}_n$ ($2 \leq n \leq \infty$). Then there is $H \leq \Gamma$ with $[\Gamma : H] = \infty$ such that the quasi-regular representation $\lambda_{\Gamma/H}$ is maximum in the order \prec of weak containment of unitary representations of Γ and is moreover irreducible..*

Proof. We again take $\Gamma = \mathbb{F}_2 = \langle \gamma_1, \gamma_2 \rangle$ for notational simplicity. As in the proof of 5.1, we can find a sequence of infinite sets $\{I_n\}$, a transitive action τ_n of Γ on I_n such that no $\gamma_i^{\tau_n}$ ($i \leq 2$) has an invariant finite set, and a sequence $\{F_n\}$ of non-empty finite sets with $F_n \subseteq I_n, \forall n$, and such that if τ^* is any transitive action of Γ on $I = \bigsqcup_n I_n$ with $\forall i \leq 2 \forall n (\gamma_i^{\tau^*}|_{F_n} =$

$\gamma_i^{\tau^n}|F_n)$, then the representation $\lambda_{\Gamma/H}$, where H is the stabilizer in τ^* of some (equivalently any) point of I , is maximum in the order of weak containment of representations of \mathbb{F}_2 . By Mackey's Irreducibility Criterion (see, e.g., Bekka-de la Harpe [BdlH], Example 10), in order to make $\lambda_{\Gamma/H}$ irreducible, it is enough to have that the action of H on Γ/H has infinite orbits except on H itself. In terms of the action τ^* , it is enough to find a point $i_0 \in I$ whose stabilizer H has the property that the orbits of $\tau^*|H$ are infinite on $I \setminus \{i_0\}$.

By a simple modification of the proof of 5.4 (by taking $S_n = \gamma_2^{\tau^n}|I_n)$, we see that we can find finite sets $\tilde{P}_n (= \bigcup_{i \leq k_n} P_i^{(n)})$, in the notation of that proof), with $\tilde{P}_n \subseteq I_n$, and for each $i_0 \in P = I \setminus \bigcup_n \tilde{P}_n$, we can find $\gamma_2^{\tau^*}$ such that $\gamma_2^{\tau^*}|F_n = \gamma_2^{\tau}|F_n$ and $\gamma_2^{\tau^*}$ fixes i_0 and acts transitively on the rest of I . Then the stabilizer H of i_0 contains γ_2 and thus (no matter how we define $\gamma_1^{\tau^*}$) $\tau^*|H$ has a single orbit off i_0 . We only need now to define $\gamma_1^{\tau^*}$ to make sure that the action τ^* is transitive and of course also have $\gamma_1^{\tau^*}|F_n = \gamma_i^{\tau}|F_n$, for every n . But this is clear from 5.4 again. \dashv

Recall that the action of a group Γ on Γ/H is *amenable* if $1_\Gamma \prec \lambda_{\Gamma/H}$. If we now take $\Gamma = \mathbb{F}_n$ and H as in 5.5, then $\pi \prec \lambda_{\Gamma/H}$ for *every* unitary representation π , so, in particular, $1_\Gamma \prec \lambda_{\Gamma/H}$, i.e., the action of Γ on I is amenable. Since we also have that $\lambda_\Gamma \prec \lambda_{\Gamma/H}$, it is easy to check that no $\gamma \neq 1$ fixes every element of Γ/H , i.e., the action of Γ on Γ/H is faithful. Thus we see that the action of Γ on $I = \Gamma/H$, where H comes from 5.5, gives another example of a faithful, transitive, amenable action of \mathbb{F}_n on a countable set I , a result first proved by van Douwen [vD]. Other such examples have been found in Glasner-Monod [GM] and Grigorchuk-Nekrashevych [GN]. However, the example coming from 5.5 has the stronger property that $\forall \pi (\pi \prec \lambda_{\Gamma/H})$ instead of just $1_\Gamma \prec \lambda_{\Gamma/H}$ and is also irreducible.

We finally note that in 5.1, the action $s_{\Gamma/\Gamma/H}$ is weakly mixing (see, e.g., Kechris-Tsankov [KT], 2.1). One can also make it free by using 2.4, (ii) in Kechris-Tsankov [KT] (or else work with $s_{\Gamma/\Gamma/H,X}$, for (X, μ) non-atomic, in which case $s_{\Gamma/\Gamma/H,X}$ is automatically free, as the action of Γ on Γ/H is faithful (see [KT], 2.4, (iii))).

6 Miscellanea

We will consider here some additional density properties in the space of actions $A(\mathbb{F}_n, X, \mu)$ of the free group \mathbb{F}_n and some of its subspaces.

Given a countable, measure preserving equivalence relation E on (X, μ) , we denote by $[E]$ its *full group*,

$$[E] = \{T \in \text{Aut}(X, \mu) : T(x)Ex, \mu\text{-a.e.}(x)\}.$$

For any group Γ , we let $A(\Gamma, [E])$ be the space of actions of Γ “contained” in $[E]$:

$$A(\Gamma, [E]) = \{a \in A(\Gamma, X, \mu) : \forall \gamma \in \Gamma (\gamma^a \in [E])\}.$$

Since $[E]$ is a separable subgroup in the uniform topology of $\text{Aut}(X, \mu)$, it follows that $A(\Gamma, [E])$ is a separable (thus Polish) space in the uniform topology of $A(\Gamma, X, \mu)$. When $\Gamma = \mathbb{F}_n$ ($1 \leq n \leq \infty$), with free generators $\gamma_1, \gamma_2, \dots$, we also let $\text{AP}(\mathbb{F}_n, [E])$ be the uniformly closed subspace of $A(\mathbb{F}_n, [E])$ consisting of all $a \in A(\mathbb{F}_n, [E])$ for which γ_1^a is aperiodic. Denoting by APER the (uniformly closed) set of aperiodic elements of $\text{Aut}(X, \mu)$, we can clearly identify $\text{AP}(\mathbb{F}_n, [E])$ with $(\text{APER} \cap [E]) \times [E]^{n-1}$ (with the product of the uniform topology).

(A) We will first consider any equivalence relation E on (X, μ) of cost $C(E) > 1$. We note that, by the argument in Ioana-Peterson-Popa [IPP], Appendix, and Gaboriau [G], it follows that for any equivalence relation F there is an equivalence relation $F \subseteq E$ with $C(F) > 1$.

Proposition 6.1 *Let E be a countable, measure preserving, ergodic equivalence relation on (X, μ) with $C(E) > 1$. Then for each $2 \leq n \leq \infty$, $\{a \in \text{AP}(\mathbb{F}_n, [E]) : C(a) > 1\}$ is dense and non-meager in the uniform topology of $\text{AP}(\mathbb{F}_n, [E])$.*

Proof. We will consider the case $n = 2$, the case of arbitrary n being similar.

(1) First we will prove density.

Claim. *There is $a \in \text{AP}(\mathbb{F}_2, [E])$ such that E_a is maximal (under a.e. inclusion) with the property that E_a is ergodic and $C(E_a) = 1$.*

Proof. Assume not. Let $a_0 \in \text{AP}(\mathbb{F}_2, [E])$ be such that E_{a_0} is ergodic and $C(E_{a_0}) = 1$ (such exists using, e.g., Kechris [Ke10], 3.5). Then by transfinite induction on $\alpha < \omega_1$ (the first uncountable ordinal), we will find $a_\alpha \in \text{AP}(\mathbb{F}_2, [E])$ such that $C(E_{a_\alpha}) = 1, \alpha < \beta \Rightarrow E_{a_\alpha} \subsetneq E_{a_\beta}$, and $E_{a_\lambda} = \bigcup_{\alpha < \lambda} E_{a_\alpha}$ for λ limit. The successor case is trivial. For the limit case, let first $E_\lambda = \bigcup_{\alpha < \lambda} E_{a_\alpha}$. Then $C(E_\lambda) = 1$ (see Gaboriau [G] or Kechris-Miller [KM],

23.5). It is thus enough to find $a_\lambda \in \text{AP}(\mathbb{F}_2, [E])$ with $E_{a_\lambda} = E_\lambda$. Note that E_λ is ergodic. So let $S_\lambda \in [E_\lambda]$ be ergodic. Since $C(E_\lambda) < 3/2$, by the proof of Kechris-Miller [KM], 27.7, there is $\varphi_\lambda \in [[E_\lambda]]$, such that E_λ is the equivalence relation generated by $S_\lambda, \varphi_\lambda$. If E_{φ_λ} is the equivalence relation generated by φ_λ , then since E_{φ_λ} is a hyperfinite subrelation of E_λ , it is easy to find an aperiodic $T_\lambda \in [E_\lambda]$ such that $E_{\varphi_\lambda} \subseteq E_{T_\lambda}$. So if $\mathbb{F}_2 = \langle \gamma_1, \gamma_2 \rangle$ and we let a_λ be defined by $\gamma_1^{a_\lambda} = S_\lambda, \gamma_2^{a_\lambda} = T_\lambda$, then clearly $a \in \text{AP}(\mathbb{F}_2, [E]), E_{a_\lambda} = E_\lambda$.

The existence of $a_\lambda, \lambda < \omega_1$, clearly violates the countable chain condition in the σ -finite measure space (E, M) , where $M(A) = \int \text{card}(A_x) d\mu(x)$, with $A_x = \{y : (x, y) \in A\}$ for any Borel set $A \subseteq E$. \dashv

So fix $a \in \text{AP}(\mathbb{F}_2, [E])$ as in the claim. Note that $E_a \subsetneq E$, from which it follows that $[E] \setminus [E_a] \neq \emptyset$. Let $\gamma_1^a = S_1, \gamma_2^a = S_2$.

Claim. *If $S'_2 \in [E] \setminus [E_a]$ and a' is defined by $\gamma_1^{a'} = S_1, \gamma_2^{a'} = S'_2$, then $C(a') > 1$.*

Proof. If not, consider $E' = E_a \vee E_{a'}$. Then, since $E_a \cap E_{a'}$ is aperiodic, $C(E') = 1$ (see Gaboriau [G] or Kechris-Miller [KM], 23.4) and as in the proof of the previous claim $E' = E_b$, for some $b \in \text{AP}(\mathbb{F}_2, [E])$, and clearly $E_a \subsetneq E_b$, contradicting the maximality of a . \dashv

We now complete the proof of density. Fix any $a_0 \in \text{AP}(\mathbb{F}_2, [E])$, with $\gamma_1^{a_0} = S_1^0, \gamma_2^{a_0} = S_2^0$. Fix also uniform open nbhds U_1, U_2 of S_1^0, S_2^0 , resp. We will find $c \in \text{AP}(\mathbb{F}_2, [E])$ with $C(c) > 1$ such that $\gamma_1^c \in U_1, \gamma_2^c \in U_2$. Since the conjugates of any aperiodic $S \in [E]$ by elements of $[E]$ are uniformly dense in $\text{APER} \cap [E]$ (see Kechris [Ke10], 3.4), we can assume that $\gamma_1^a = S_1 \in U_1$ (by replacing a by a conjugate action within $[E]$ if necessary). Now $[E_a]$ is a uniformly closed proper subgroup of $[E]$, so it must have empty interior, otherwise it would be open, thus clopen, violating the connectedness of $[E]$ in the uniform topology (see Kechris [Ke10], 3.12). Thus $[E] \setminus [E_a]$ is uniformly dense in $[E]$ and so $U_2 \cap ([E] \setminus [E_a]) \neq \emptyset$. Then pick $T_2 \in U_2 \cap ([E] \setminus [E_a])$. Let $c \in \text{AP}(\mathbb{F}_2, [E])$ be such that $\gamma_1^c = \gamma_1^{a_0} = S_1 \in U_1, \gamma_2^c = T_2 \in U_2$. By the previous claim $C(c) > 1$ and we are done.

(2) Next we prove non-meagerness.

It will be convenient to use the following notation: For any topological space X and $P \subseteq X$, we put

$$\forall^* x \in X P(x) \Leftrightarrow P \text{ is comeager in } X.$$

We also let

$$A = \text{APER} \cap [E],$$

so that $\text{AP}(\mathbb{F}_2, [E])$ can be identified with $A \times [E]$. All these spaces are equipped with the uniform topology.

Assume that $\{a \in \text{AP}(\mathbb{F}_2, [E]) : C(a) = 1\}$ is comeager, towards a contradiction. Then letting for each $V_0, V_1, \dots \in \text{Aut}(X, \mu)$, $E_{V_0, V_1, \dots}$ be the equivalence relation generated by V_0, V_1, \dots , we have

$$\forall^*(S, T) \in A \times [E] (C(E_{S, T}) = 1),$$

so by the Kuratowski-Ulam Theorem

$$\forall^* S \in A \forall^* T \in [E] (C(E_{S, T}) = 1).$$

Claim. For any n ,

$$\forall^* S \in A \forall^* T_0 \in [E] \cdots \forall^* T_n \in [E] (C(E_{S, T_0, T_1, \dots, T_n}) = 1).$$

Proof. By induction on n . This is clear for $n = 0$. Assume it is true for n . Then using this and the $n = 0$ case, we have

$$\begin{aligned} \forall^* S \in A \forall^* T_0 \in [E] \cdots \forall^* T_n \in [E] \forall^* T_{n+1} \in [E] \\ (C(E_{S, T_0, \dots, T_n}) = 1 \wedge C(E_{S, T_{n+1}}) = 1). \end{aligned}$$

For S, T_0, \dots, T_{n+1} as above,

$$E_{S, T_0, \dots, T_n} \cap E_{S, T_{n+1}} \supseteq E_S$$

and

$$E_{S, T_0, \dots, T_n, T_{n+1}} = E_{S, T_0, \dots, T_n} \vee E_{S, T_{n+1}},$$

while E_S is aperiodic. Thus it follows as before that

$$C(E_{S, T_0, \dots, T_{n+1}}) = 1,$$

i.e.,

$$\forall^* S \in A \forall^* T_0 \in [E] \cdots \forall^* T_{n+1} \in [E] (C(E_{S, T_0, \dots, T_{n+1}}) = 1).$$

⊣

Using this claim, we then have

$$\forall^* S \in A \forall^* (T_0, T_1, \dots) \in [E]^{\mathbb{N}} (C(E_{S, T_0, T_1, \dots}) = 1),$$

since $E_{S,T_0,T_1,\dots} = \bigcup_n E_{S,T_0,T_1,\dots,T_n}$ and $\{E_{S,T_0,T_1,\dots,T_n}\}$ is an increasing sequence of cost 1 equivalence relations for comeager many $(S, (T_0, T_1, \dots)) \in A \times [E]^\mathbb{N}$.

On the other hand, we have

$$\forall^* S \in A \forall^* (T_0, T_1, \dots) \in [E]^\mathbb{N} (E_{S,T_0,T_1,\dots} = E),$$

which is a contradiction, since $C(E) > 1$. Indeed, it is enough to verify that

$$\forall^* (T_0, T_1, \dots) \in [E]^\mathbb{N} (\{T_n\} \text{ is dense in } [E]).$$

To see this, let $\{g_n\}$ be dense in $[E]$. Let $\delta_u(S, T) = \mu(\{x : S(x) \neq T(x)\})$ be the uniform metric on $\text{Aut}(X, \mu)$. Then for any $\epsilon > 0, n \in \mathbb{N}$,

$$Y_{\epsilon,n} = \{(T_0, T_1, \dots) \in [E]^\mathbb{N} : \exists i (d_u(T_i, g_n) < \epsilon)\}$$

is open and dense in $[E]^\mathbb{N}$, thus

$$Y = \bigcap_{\epsilon,n} Y_{\epsilon,n}$$

is dense G_δ and clearly

$$(T_0, T_1, \dots) \in Y \Rightarrow \{T_n\} \text{ is dense in } [E].$$

⊣

Corollary 6.2 *Let E be a countable, measure preserving, ergodic equivalence relation on (X, μ) with $C(E) > 1$. Then for each $2 \leq n < \infty$, there is $\epsilon > 0$ such that*

$$\{a \in \text{AP}(\mathbb{F}_n, [E]) : C(a) \geq 1 + \epsilon\}$$

has non-empty interior in $\text{AP}(\mathbb{F}_n, [E])$ with the uniform topology.

Proof. It is shown in Kechris [Ke10], Section 10, Remark following 10.14, that for infinite, finitely generated groups Γ the function $a \mapsto C(a)$ on $A(\Gamma, X, \mu)$ is upper semicontinuous in the uniform topology. It follows that for each δ , $\{a \in A(\Gamma, X, \mu) : C(a) \geq \delta\}$ is uniformly closed. Thus

$$\{a \in \text{AP}(\mathbb{F}_n, [E]) : C(a) > 1\}$$

is non-meager in $\text{AP}(\mathbb{F}_n, [E])$ and the union of the sequence of closed sets

$$\{a \in \text{AP}(\mathbb{F}_n, [E]) : C(a) \geq 1 + \frac{1}{m}\}.$$

Thus for some $\epsilon = \frac{1}{m}$, $\{a \in \text{AP}(\mathbb{F}_n, [E]) : C(a) \geq 1 + \epsilon\}$ has non-empty interior in $\text{AP}(\mathbb{F}_n, [E])$. \dashv

This corollary shows that there are two elements $S, T \in [E]$, S aperiodic, such that $C(E_{S,T}) \geq 1 + \epsilon$ and for any $S', T' \in [E]$, S' aperiodic, which are sufficiently close to S, T in the uniform topology (i.e., differ by S, T on a set of very small measure), we still have $C(E_{S',T'}) \geq 1 + \epsilon$.

In Kechris-Miller [KM], 28.8, it was shown, for E as in 6.2, that there is a free ergodic $a \in A(\mathbb{F}_2, [E])$. Then of course $C(a) = 2$. It would be natural to think that an open nbhd of any such action would be a witness to the conclusion of the previous corollary. However, this is not the case in view of the following example due to Hjorth.

Proposition 6.3 (Hjorth) *Let $\mathbb{F}_n = \langle \gamma_1, \gamma_2, \dots \rangle$ be the free group with free generators $\gamma_1, \gamma_2, \dots$ ($1 \leq n \leq \infty$). Let $a \in A(\mathbb{F}_n, X, \mu)$ be such that γ_1^a is ergodic. Assume that there is an ergodic $U \in \text{Aut}(X, \mu)$ which commutes with a , i.e., U commutes with each γ_i^a . Then there is a sequence $a_m \in A(\mathbb{F}_n, X, \mu)$ with $\gamma_1^{a_m}$ ergodic, $C(a_m) = 1$ and $a_m \rightarrow a$ uniformly.*

Proof. We take $n = 2$ for notational simplicity and let $\gamma_1^a = S, \gamma_2^a = T$.

First we note that $C(E_{S,T,U}) = 1$; this follows from Kechris-Miller [KM], 24.8. It is thus enough to find $S_m, T_m \in \text{Aut}(X, \mu)$ such that S_m is ergodic, $S_m \rightarrow S, T_m \rightarrow T$ uniformly and $E_{S_m, T_m} = E_{S, T, U}$ (then a_m given by $\gamma_1^{a_m} = S_m, \gamma_2^{a_m} = T_m$ works).

Next we note the following fact: if E is a countable, measure preserving equivalence relation, $E_S \subseteq E$ and $A = \{x : xEU(x)\}$ has positive measure, then $E_U \subseteq E$. Indeed, given any $x \in X$, by the ergodicity of S , there is $n \in \mathbb{Z}$ such that $S^n(x) \in A$ and so

$$xE S^n(x) E U(S^n(x)) = S^n(U(x)) E U(x),$$

so $E_U \subseteq E$. Similarly, if $E_U \subseteq E$ and $B = \{x : xET(x)\}$ has positive measure, then $E_T \subseteq E$.

To find S_m, T_m , let first $A_m \subseteq X$ be a Borel set with $\mu(A_m) < 1/m$. Then $\mu(U(A_m)) = \mu(A_m) < 1/m$ and so if $C_m = \{x : T(x) \notin U(A_m)\}$, $\mu(C_m) > 1 - 1/m$ and $T(C_m) \cap U(A_m) = \emptyset$. Let $B_m = C_m \cap (X \setminus A_m)$, so that $\mu(B_m) > 1 - 2/m$. Then, using the ergodicity of U , we can find $T_m \in [E_U]$ such that $T_m|_{B_m} = T|_{B_m}$ (thus T_m, T differ only on a set of measure $< 2/m$), and $T_m|_{A_m} = U|_{A_m}$. Let also $S_m = S$. Clearly $S_m \rightarrow S, T_m \rightarrow T$ uniformly.

Let $E_m = E_{S_m, T_m}$. Then by the preceding fact, applied to E_m , we conclude that $E_m \supseteq E_{S, T, U}$ and since $T_m \in [E_U]$, $E_m = E_{S_m, T_m} \subseteq E_{S, T, U}$, so $E_m = E_{S, T, U}$. \dashv

Take now $n = 2$ in 6.3 and a free, ergodic action $a \in A(\mathbb{F}_2, X, \mu)$ for which there is an ergodic $U \in \text{Aut}(X, \mu)$, which commutes with a , and let a_m be as in the conclusion of 6.3. Let F be the equivalence relation generated by a and $\{a_m\}$, and $E \supseteq F$ be an equivalence relation such that $C(E) > 1$. Then no nbhd of a is a witness to the conclusion of 6.2 for this E .

It is clear from 5.2 that $\{a \in A(\mathbb{F}_n, X, \mu) : E_a \text{ is aperiodic, hyperfinite}\}$ is weakly dense in $A(\mathbb{F}_n, X, \mu)$ and thus so if $\{a \in A(\mathbb{F}_n, X, \mu) : C(a) = 1\}$ ($1 \leq n \leq \infty$). Recall that E_a is *aperiodic* if (almost) all its equivalence classes are infinite. Using the preceding result we can actually prove a stronger statement.

Proposition 6.4 *The set*

$$\{a \in A(\mathbb{F}_n, X, \mu) : E_a \text{ is not hyperfinite \& } C(a) = 1\}$$

is weakly dense in $A(\mathbb{F}_n, X, \mu)$ ($2 \leq n \leq \infty$).

Proof. Again we take $n = 2$ for notational simplicity.

Fix $a_0 \in A(\mathbb{F}_2, X, \mu)$ and a weak nbhd U of a in order to find $a \in U$ with E_a not hyperfinite and $C(a) = 1$.

By Kechris [Ke10], Section 10, **(G)**, we can assume that there is an action $b_0 \in A(\mathbb{F}_2 \times \mathbb{Z}, X, \mu)$ with $a_0 = b_0|_{\mathbb{F}_2}$. Now $\mathbb{F}_2 \times \mathbb{Z}$ has the Haagerup Approximation Property (HAP), so by Hjorth [Hj09] there is a mixing action $b'_0 \in A(\mathbb{F}_2 \times \mathbb{Z}, X, \mu)$ as close as we want to b_0 in the weak topology. Let c_0 be a free, mixing action in $A(\mathbb{F}_2 \times \mathbb{Z}, X, \mu)$ and consider $b'_0 \times c_0$. It is free, mixing and $b'_0 \prec b'_0 \times c_0$, i.e., there is an isomorphic copy of $b'_0 \times c_0$ as close as we want to b'_0 in the weak topology. Thus there is a free mixing action $d_0 \in A(\mathbb{F}_2 \times \mathbb{Z}, X, \mu)$ such that $d_0|_{\mathbb{F}_2} \in U$. Then, by 6.3, $d_0|_{\mathbb{F}_2}$ is the uniform limit of a sequence of actions $a_m \in A(\mathbb{F}_2, X, \mu)$ with $C(a_m) = 1$.

Lemma 6.5 *For any infinite, countable group Γ , the set $\{a \in A(\Gamma, X, \mu) : E_a \text{ is hyperfinite}\}$ is uniformly closed.*

Granting this, since clearly $E_{d_0|_{\mathbb{F}_2}}$ is not hyperfinite, it follows that we can also assume that E_{a_m} is not hyperfinite and the proof is complete.

Proof of 6.5. Let $a_n \in A(\Gamma, X, \mu)$, E_{a_n} hyperfinite and $a_n \rightarrow a \in A(\Gamma, X, \mu)$ uniformly. Let $E_n = E_{a_n}$, $E = E_a$. Clearly for each n , $F_n = \bigcap_{m>n} E_m$ is hyperfinite and $F_0 \subseteq F_1 \subseteq \dots$, so

$$\bigcup_n F_n = \bigcup_n \bigcap_{m>n} E_m$$

is hyperfinite. It is thus enough to check that $E \subseteq \bigcup_n \bigcap_{m>n} E_m$. If not, there is $\gamma \in \Gamma$ and a set of positive measure A such that

$$x \in A \Rightarrow (x, \gamma^a(x)) \notin \bigcup_n \bigcap_{m>n} E_m.$$

i.e., for $x \in A$ and for infinitely many n , $(x, \gamma^a(x)) \notin E_n$. Now $\gamma^{a_n} \rightarrow \gamma^a$ uniformly, so for all large enough n , $\mu(\{x \in A: \gamma^a(x) = \gamma^{a_n}(x)\}) > 0$, a contradiction. \dashv

(B) There have been some very interesting recent results that have the following general form: Let P be a property of countable, measure preserving equivalence relations on (X, μ) , (which we intuitively think as strongly violating hyperfiniteness). Then there exists an ergodic equivalence relation E such that *every* ergodic $F \subseteq E$ either is hyperfinite or else has property P . Chifan-Ioana [CI] show that for P being the property of strong ergodicity (also called E_0 -ergodicity), this holds for the equivalence relation E induced by the shift action of Γ on $[0, 1]^\Gamma$ for any infinite, countable group Γ . Ozawa [O] showed that the same holds when E is the equivalence relation induced by the action of $\text{SL}_2(\mathbb{Z})$ on \mathbb{T}^2 . Finally, Ioana [I09] showed that for P being the property of being rigid, this holds again for the equivalence relation induced by the action of $\text{SL}_2(\mathbb{Z})$ on \mathbb{T}^2 .

Let us note here that for pairs P, E satisfying the above dichotomy, we have the following density result.

Proposition 6.6 *Let P be a property of countable, measure preserving equivalence relations that implies non-hyperfiniteness. Let E be a countable, measure preserving, ergodic equivalence relation which is not hyperfinite such that every ergodic subequivalence relation $F \subseteq E$ is either hyperfinite or has property P . Then $\{a \in A(\mathbb{F}_n, [E]): E_a \text{ has property } P\}$ is uniformly dense in $\text{AP}(\mathbb{F}_n, [E])$, $2 \leq n \leq \infty$.*

Proof. Take again $n = 2$ for notational simplicity. As in the proof of 6.1, let $S_0 \in [E]$ be ergodic such that E_{S_0} is maximal under inclusion. Fix

$(S, T) \in \text{AP}(\mathbb{F}_2, [E])$ and uniform nbhds U, V of S, T , resp. Since we can find $K \in [E]$ such that $KS_0K^{-1} \in U$, we may as well assume that $S_0 \in U$. Now $[E] \setminus [E_{S_0}]$ is uniformly dense in $[E]$, so let $T_0 \in [E] \setminus [E_{S_0}]$ be in V . Then E_{S_0, T_0} is ergodic and non-hyperfinite, by the maximality of E_{S_0} , and thus has property P , which completes the proof. \dashv

Corollary 6.7 *In the context of 6.6, $\{a \in A(\mathbb{F}_n, X, \mu) : E_a \text{ has property } P\}$ is weakly dense in $A(\mathbb{F}_n, X, \mu)$.*

It follows, for example, that the strongly ergodic (resp., rigid) actions are weakly dense in $A(\mathbb{F}_n, X, \mu)$, $2 \leq n \leq \infty$, a fact that can be also proved more directly as pointed out in Ioana [I09], 6.2. Also Abért informed me that he has shown, by a different method, that the actions in $A(\mathbb{F}_n, X, \mu)$ that have spectral gap are weakly dense.

A Appendix. Some facts about co-induced actions

Suppose that a countable group Δ acts on a countable set T and let $\sigma : \Delta \times T \rightarrow \text{Aut}(X, \mu)$ be a cocycle of the action of Δ on T with values in $\text{Aut}(X, \mu)$, i.e., a map satisfying the property

$$\sigma(\delta_1 \delta_2, t) = \sigma(\delta_1, \delta_2 \cdot t) \sigma(\delta_2, t),$$

for $\delta_1, \delta_2 \in \Delta, t \in T$. We define an action of Δ on (Y, ν) , where $Y = X^T, \nu = \mu^T$ (the product measure), by

$$(\delta \cdot f)(t) = \sigma(\delta^{-1}, t)^{-1}(f(\delta^{-1} \cdot t)).$$

It is easily checked that this is a measure preserving action of Δ on (Y, ν) .

If $\tau : \Delta \times T \rightarrow \text{Aut}(X, \mu)$ is another cocycle which is cohomologous to σ , i.e., there is $f : T \rightarrow \text{Aut}(X, \mu)$ such that $\tau(\delta, t) = f(\delta \cdot t) \sigma(\delta, t) f(t)^{-1}$, then the action induced by τ is isomorphic to the action induced by σ via the map $\varphi : Y \rightarrow Y$ given by

$$\varphi(p)(t) = f(t)(p(t)).$$

Let now $\Gamma \leq \Delta$ be a subgroup and let T be a transversal for the left cosets of Γ with $1 \in T$. Let Δ act on T by defining $\delta \cdot t$ to be the unique element of T in $\delta t \Gamma$ and let $\rho : \Delta \times T \rightarrow \Gamma$ be the cocycle defined by

$$(\delta \cdot t) \rho(\delta, t) = \delta t.$$

If $a \in A(\Gamma, X, \mu)$, then ρ gives rise to the cocycle $\sigma: \Delta \times T \rightarrow \text{Aut}(X, \mu)$ defined by $\sigma(\delta, t) = \rho(\delta, t)^a$. We call the action on (Y, ν) given by σ the *co-induced action* of a , in symbols

$$\text{CInd}_\Gamma^\Delta(a).$$

Thus $b = \text{CInd}_\Gamma^\Delta(a)$ is the action of Δ on (X^T, μ^T) given by

$$(\delta \cdot f)(t) = \rho(\delta^{-1}, t)^{-1} \cdot f(\delta^{-1} \cdot t),$$

where the action on the right-hand side is the action a .

Note that T can be identified with Δ/Γ and thus (X^T, μ^T) with the space $(X^{\Delta/\Gamma}, \mu^{\Delta/\Gamma})$. The action of Δ on T becomes then the usual action $\delta \cdot \delta'\Gamma = \delta\delta'\Gamma$ of Δ on Δ/Γ .

Various properties of the co-induced action are given in Ioana [I07] and Kechris [Ke10], Section 10, **(G)**. We record a few that we use in this paper:

- (i) $a \sqsubseteq \text{CInd}_\Gamma^\Delta(a)|\Gamma$, in fact the map $f \mapsto f(1)$ demonstrates that a is a factor of $\text{CInd}_\Gamma^\Delta(a)|\Gamma$.

(ii)

$$\begin{aligned} a &\mapsto \text{CInd}_\Gamma^\Delta(a) \\ A(\Gamma, X, \mu) &\rightarrow A(\Delta, Y\nu) \end{aligned}$$

is continuous in the weak topology.

- (iii) $a \cong b \Rightarrow \text{CInd}_\Gamma^\Delta(a) \cong \text{CInd}_\Gamma^\Delta(b)$.

We prove below some further properties of the co-induced action that we also need in this paper.

Proposition A.1 *Co-inducing preserves weak containment, i.e.,*

$$a \prec b \Rightarrow \text{CInd}_\Gamma^\Delta(a) \prec \text{CInd}_\Gamma^\Delta(b).$$

Proof. Since $a \prec b$, there is a sequence b_n such that $b_n \cong b$ and $b_n \rightarrow a$ weakly. Then $\text{CInd}_\Gamma^\Delta(b_n) \cong \text{CInd}_\Gamma^\Delta(b)$ and by (ii) above $\text{CInd}_\Gamma^\Delta(b_n) \rightarrow \text{CInd}_\Gamma^\Delta(a)$, so $\text{CInd}_\Gamma^\Delta(a) \prec \text{CInd}_\Gamma^\Delta(b)$. \dashv

If $\Gamma \leq \Delta$ and (X, μ) is a measure space (perhaps with atoms), we denote by $s_{\Delta, \Delta/\Gamma, X}$ the shift action of Δ on $X^{\Delta/\Gamma}$ corresponding to the canonical action of Δ on Δ/Γ :

$$(\delta_1 \cdot f)(\delta_2\Gamma) = f(\delta_1^{-1}\delta_2\Gamma).$$

If $X = 2 = \{0, 1\}$ and $\mu(\{0\}) = \mu(\{1\}) = 1/2$, we simply write $s_{\Delta, \Delta/\Gamma}$. If T is a transversal for Δ/Γ with $1 \in T$, then Δ/Γ can be identified with T and $s_{\Delta, \Delta/\Gamma, X}$ is the action

$$(\delta \cdot f)(t) = f(\delta^{-1} \cdot t),$$

where Δ acts on T in the previous sense.

Proposition A.2 *Let $H \leq \Gamma \leq \Delta$ and (X, μ) a measure space (perhaps with atoms). Then*

$$\text{CInd}_{\Gamma}^{\Delta}(s_{\Gamma, \Gamma/H, X}) \cong s_{\Delta, \Delta/H, X}.$$

Proof. Fix a transversal S for the left cosets of H in Γ containing 1 and a transversal T for the left cosets of Γ in Δ containing 1. Then $TS = \{ts : t \in T, s \in S\}$ is a transversal for the left cosets of H in Δ .

The action $s_{\Gamma, \Gamma/H, X}$ is the action of Γ on $X^{\Gamma/H} = X^S$ defined by

$$\gamma \cdot p(s) = p(\gamma^{-1} \cdot s),$$

where Γ acts on S by $\gamma \cdot s =$ (the unique element of S in the coset γsH). Let also $\sigma : \Gamma \times S \rightarrow H$ be the associated cocycle given by $(\gamma \cdot s)\sigma(\gamma, s) = \gamma s$.

The action $\text{CInd}_{\Gamma}^{\Delta}(s_{\Gamma, \Gamma/H, X})$ is the action of Δ on $(X^{\Gamma/H})^{\Delta/\Gamma} = (X^S)^T$, given by

$$\delta \cdot q(t) = \rho(\delta^{-1}, t)^{-1} \cdot q(\delta^{-1} \cdot t),$$

where Δ acts on T in the usual way and $\rho : \Delta \times T \rightarrow \Gamma$ is the associated cocycle given by $(\delta \cdot t)\rho(\delta, t) = \delta t$.

The measure spaces $((X^S)^T, (\mu^S)^T)$ and $(X^{S \times T}, \mu^{S \times T})$ are isomorphic via the map $p \mapsto \varphi(p) = q$, where $q(s, t) = p(t)(s)$. Clearly $(s, t) \mapsto ts$ is a bijection of $S \times T$ with TS and thus $X^{S \times T}$ is identified with $X^{TS} = X^{\Delta/H}$. Therefore $\psi : (X^S)^T \rightarrow X^{TS}$, given by $\psi(p) = q$, where $q(ts) = \varphi(p)(s, t) = p(t)(s)$, is an isomorphism of $((X^S)^T, (\mu^S)^T)$ with $(X^{TS}, \mu^{TS}) = (X^{\Delta/H}, \mu^{\Delta/H})$ and we will show that it sends $\text{CInd}_{\Gamma}^{\Delta}(s_{\Gamma, \Gamma/H, X})$ to $s_{\Delta, \Delta/H, X}$.

We have $(\delta \cdot p)(t) = \rho(\delta^{-1} \cdot t)^{-1} \cdot p(\delta^{-1} \cdot t)$, so

$$\begin{aligned} \psi(\delta \cdot p)(ts) &= (\delta \cdot p)(t)(s) = \rho(\delta^{-1} \cdot t)^{-1} \cdot p(\delta^{-1} \cdot t)(s) \\ &= p(\delta^{-1} \cdot t)(\rho(\delta^{-1} \cdot t) \cdot s) \\ &= \psi(p)((\delta^{-1} \cdot t)(\rho(\delta^{-1}, t) \cdot s)). \end{aligned}$$

On the other hand,

$$(\delta \cdot \psi(p))(ts) = \psi(p)(\delta^{-1} \cdot ts)$$

Now $(\delta^{-1} \cdot t)\rho(\delta^{-1}, t) = \delta^{-1}t$, so $\delta^{-1}ts = (\delta^{-1} \cdot t)\rho(\delta^{-1}, t)s$, thus if we put $\gamma = \rho(\delta^{-1}, t) \in \Gamma$, we have

$$\begin{aligned}\delta^{-1}ts &= (\delta^{-1} \cdot t)\gamma s \\ &= (\delta^{-1} \cdot t)(\gamma \cdot s)\sigma(\gamma, s) \\ &= (\delta^{-1} \cdot t)(\rho(\delta^{-1}, t) \cdot s)\sigma(\gamma, s)\end{aligned}$$

and $\sigma(\gamma, s) \in H$, thus

$$\delta^{-1} \cdot ts = (\delta^{-1} \cdot t)(\rho(\delta^{-1}, t) \cdot s),$$

so

$$\delta \cdot \psi(p)(ts) = \psi(p)((\delta^{-1} \cdot t)(\rho(\delta^{-1}, t) \cdot s)),$$

i.e., $\psi(\delta \cdot p) = \delta \cdot \psi(p)$, and the proof is complete. \dashv

Assume $\Gamma \leq \Delta$ and let now $a \in A(\Delta, X, \mu)$ be an action of the bigger group Δ . One can form the restriction $a|_{\Gamma} \in A(\Gamma, X, \mu)$ and then co-induce that to get $\text{CInd}_{\Gamma}^{\Delta}(a|_{\Gamma})$. We will describe below this action.

Consider first the (diagonal) product action $a^{\Delta/\Gamma}$ on $(X^{\Delta/\Gamma}, \mu^{\Delta/\Gamma}) = (X^T, \mu^T)$:

$$\delta \cdot f = (t \mapsto \delta \cdot f(t)),$$

where the action on the right-hand side is the action a . We also have the shift action $s_{\Delta, \Delta/\Gamma, X}$ on $(X^{\Delta/\Gamma}, \mu^{\Delta/\Gamma})$. Note that these actions commute, i.e., for each $\delta_1, \delta \in \Delta$

$$\delta_1^{a^{\Delta/\Gamma}} \delta_2^{s_{\Delta, \Delta/\Gamma, X}} = \delta_2^{s_{\Delta, \Delta/\Gamma, X}} \delta_1^{a^{\Delta/\Gamma}},$$

so we can define a new action, denoted by $a^{\Delta/\Gamma} \circledast s_{\Delta, \Delta/\Gamma, X}$, which is given by

$$\delta \cdot f(t) = \delta \cdot f(\delta^{-1} \cdot t).$$

Proposition A.3 *For $\Gamma \leq \Delta$ and for each action $a \in A(\Delta, X, \mu)$,*

$$\text{CInd}_{\Gamma}^{\Delta}(a) \cong a^{\Delta/\Gamma} \circledast s_{\Delta, \Delta/\Gamma, X}.$$

Proof. Going back to the beginning of this Appendix, let the cocycle $\sigma_1: \Delta \times T \rightarrow \text{Aut}(X, \mu)$ be given by $\sigma_1(\delta, t) = \delta^a$. Let also $\sigma_2(\delta, t) = \rho(\delta, t)^a$. Then the action of Δ on (X^T, μ^T) corresponding to σ_1 is $a^{\Delta/\Gamma} \circledast s_{\Delta, \Delta/\Gamma, X}$, while the action corresponding to σ_2 is $\text{CInd}_{\Gamma}^{\Delta}(a)$. It is thus enough to show

that σ_1, σ_2 are cohomologous, i.e., there is $f: T \rightarrow \text{Aut}(X, \mu)$ such that $\sigma_2(\delta, t) = f(\delta \cdot t)\sigma_1(\delta, t)f(t)^{-1}$. By definition we have

$$\sigma_2(\delta, t) = \rho(\delta, t)^a,$$

where $\rho(\delta, t) = (\delta \cdot t)^{-1}\delta t$, thus

$$\begin{aligned}\sigma_2(\delta, t) &= ((\delta \cdot t)^{-1})^a \delta^a t^a \\ &= ((\delta \cdot t)^{-1})^a \sigma_1(\delta, t) t^a,\end{aligned}$$

so

$$f(t) = (t^{-1})^a$$

works. ◻

If $\Gamma \leq \Delta$, then the action of Δ on Δ/Γ is *amenable* if it admits a finitely additive invariant probability measure (defined on all subsets of Δ/Γ). For more about these actions, see, e.g., Glasner-Monod [GM] and Kechris-Tsankov [KT]. We conclude this appendix with the following question.

Problem A.4 *Let $\Gamma \leq \Delta$ and assume that the action of Δ on Δ/Γ is amenable. Is it true that for any $a \in A(\Delta, X, \mu)$,*

$$a \prec \text{CInd}_\Gamma^\Delta(a|\Gamma)?$$

Note the assumption that the action of Δ on Δ/Γ is amenable is necessary because if $a = i_\Delta$ is the trivial action of Δ on (X, μ) , then $\text{CInd}_\Gamma^\Delta(a|\Gamma) \cong s_{\Delta, \Delta/\Gamma, X}$, and $i_\Delta \prec s_{\Delta, \Delta/\Gamma, X}$ implies the amenability of the action of Δ on Δ/Γ (see Kechris-Tsankov [KT]).

By extending the arguments in Kechris-Tsankov [KT], we can show that A.4 has a positive answer in certain cases, e.g., when $a = i_\Delta$ or $a = s_{\Delta, \Delta/H, X}$ (for any $H \leq \Delta$), but the general case remains open.

References

- [AN] M. Abért and N. Nikolov, Rank gradient, cost of groups and the rank versus Heegard genus problem, *arXiv:math/0701925*, 2008.
- [AW] M. Abért and B. Weiss, Bernoulli actions are weakly contained in any free action, *forthcoming*.

- [BdlH] M.B. Bekka and P. de la Harpe, Irreducibility of unitary group representations and reproducing kernel Hilbert spaces, *Expo. Math.*, **21** (2003), 115–149.
- [CI] I. Chifan and A. Ioana, Ergodic subequivalence relations induced by a Bernoulli action, *arXiv:0802.2998*, 2008.
- [FM] J. Feldman and C.C. Moore, Ergodic equivalence relations, cohomology and von Neumann algebras, I, *Trans. Amer. Math. Soc.*, **234**(2) (1977), 289–324.
- [G] D. Gaboriau, Coût des relations d’équivalence et des groupes, *Inv. Math.*, **139** (2000), 41–98.
- [GM] Y. Glasner and N. Monod, Amenable actions, free products and a fixed point property, *Bulletin London Math. Soc.*, **39**(1) (2007), 138–150.
- [GTW] E. Glasner, J.-P. Thouvenot and B. Weiss, Every countable group has the weak Rokhlin property, *Bull. London Math. Soc.*, **38**(6) (2006), 932–936.
- [GN] R.I. Grigorchuk and V.V. Nekrashevych, Amenable actions of non-amenable groups, *J. Math. Sci. (N.Y.)*, **140**(3) (2007), 391–397.
- [Hj02] G. Hjorth, A converse to Dye’s Theorem, *Trans. Amer. Math. Soc.*, **357**(8) (2002), 3083–3103.
- [Hj09] G. Hjorth, Mixing actions of groups with the Haagerup approximation property, *Fund. Math.*, **203**(1) (2009), 47–56.
- [I07] A. Ioana, Orbit inequivalent actions for groups containing a copy of \mathbb{F}_2 , *arXiv:math/0701027*, 2007.
- [I09] A. Ioana, Relative property (T) for the subequivalence relations induced by the actions of $\mathrm{SL}_2(\mathbb{Z})$ on \mathbb{T}^2 , *arXiv:0901.1874*, 2009.
- [IPP] A. Ioana, J. Peterson and S. Popa, Amalgamated free products of w -rigid factors and calculation of their symmetry groups, *Acta Math.*, **200**(1) (2008), 85–153.

- [Ke07] A.S. Kechris, Unitary representations and modular actions, *J. Math. Sc.*, **140(3)** (2007), 398–425.
- [Ke10] A.S. Kechris, *Global aspects of ergodic group actions*, Math. Surveys and Monographs, **160**, Amer. Math. Society, 2010.
- [KM] A.S. Kechris and B. Miller, *Topics in Orbit Equivalence*, Lecture Notes in Math. **1852**, Springer, 2004.
- [KT] A.S. Kechris and T. Tsankov, Amenable actions and almost invariant sets, *Proc. Amer. Math. Soc.*, **136(2)** (2007), 687–697.
- [LSh] A. Lubotzky and Y. Shalom, Finite representations in the unitary dual and Ramanujan groups, *Contemp. Math.*, **347** (2004), 173–189.
- [LZ] A. Lubotzky and A. Zuk, *On property (τ)* , preprint, 2003.
(posted in <http://www.ma.huji.ac.il/~alexclub/>, 2003)
- [LS] R.C. Lyndon and P.E. Schupp, *Combinatorial Group Theory*, Springer, 2001.
- [O] N. Ozawa, An example of a solid von Neumann algebra, *Hokkaido Math. J.*, **38(3)** (2009), 557–561.
- [vD] E.K. van Douwen, Measures invariant under actions of F_2 , *Topology Appl.*, **34(1)** (1990), 53–68.
- [Y] H. Yoshizawa, Some remarks on unitary representations of the free group, *Osaka Math. J.*, **3** (1951), 55–63.
- [Z] R. Zimmer, *Ergodic Theory and Semisimple Groups*, Birkhäuser, 1984.

Department of Mathematics
 California Institute of Technology
 Pasadena, CA 91125
 kechris@caltech.edu